

## RESISTANT WEEDS WARNING

GROUP **B** HERBICIDE

ChemAg Amaze Herbicide is a member of the Imidazolinone group of herbicides. ChemAg Amaze has the inhibition of Acetolactate synthase (ALS) mode of action. For weed resistance management, ChemAg Amaze is a Group B Herbicide. Some naturally occurring weed biotypes resistant to Amaze and other Group B herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These weeds will not be controlled by Amaze or other Group B herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, ChemAg Pty Ltd accepts no liability for any losses that may result from failure of AMAZE to control resistant weeds.

### STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point.

If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If not available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should NOT be burnt.

### SAFETY DIRECTIONS

Will damage eyes. Will irritate skin. Avoid contact with eyes and skin. When preparing spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

CONTENTS 2KG

## CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



# ChemAg AMAZE 700 WG HERBICIDE

ACTIVE CONSTITUENT:  
700 g/kg IMAZETHAPYR

GROUP **B** HERBICIDE

For the pre- or post-emergence control of certain weeds in Centrosema (Cavalcade), Chickpeas, faba beans, field peas, Lucerne, mung beans, peanuts, serradella, soybeans and subterranean clover as per the DIRECTIONS FOR USE table.

**DIRECTIONS FOR USE: READ THE ATTACHED BOOKLET BEFORE USING THIS PRODUCT.**

APVMA Approval No: 58746/2/1004



ChemAg

ChemAg Pty Ltd  
ACN 009 234 691  
Suite 11, 11 Preston St, Como, WA 6152  
Tel: (08) 9368 7474 Fax: (08) 9368 7475  
Email: sales@chemag.com.au  
Web: www.chemag.com.au

### FIRST AID

If poisoning occurs, contact a doctor or Poison's Information Centre. Telephone 131 126 Australia wide.

### MSDS

Additional information is listed in the Material Safety Data Sheet.

### CONDITIONS OF SALE

All conditions and warranties rights and remedies implied by law or arising in contact or tort whether due to the negligence of ChemAg Pty Ltd or otherwise are hereby expressly excluded so far as the same may be legally be done provided however that any rights of the Buyer pursuant to non excludable conditions or warranties of the Trade Practices act 1974 or any relevant legislation of any State are expressly excluded. This product must be used or applied strictly in accordance with the instructions appearing hereon. This product is solely sold for use in Australia and must not be exported without the prior written consent of ChemAg Pty Ltd.

THIS PRODUCT IS NOT CONSIDERED TO BE A DANGEROUS GOOD UNDER THE AUSTRALIAN CODE FOR THE TRANSPORT OF DANGEROUS GOODS BY ROAD AND RAIL

FOR SPECIALIST ADVICE IN AN EMERGENCY ONLY  
PHONE 1800 033 111  
TOLL FREE - ALL HOURS - AUSTRALIA WIDE

Batch No:

Date of Manufacture:



**CAUTION**

**KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE  
OPENING OR USING**

**CHEMAG  
AMAZE 700 WG  
HERBICIDE**

**ACTIVE CONSTITUENT: 700 g/kg IMAZETHAPYR**

**GROUP B HERBICIDE**

For the pre- or post-emergence control of certain weeds in Centrosema (Cavalcade), Chickpeas, faba beans, field peas, Lucerne, mung beans, peanuts, serradella, soybeans and subterranean clover as per the DIRECTIONS FOR USE table.

**THIS BOOKLET IS PART OF THE LABEL.**

**APVMA Approval No: 58746/1004**

ChemAg Pty. Ltd. ACN 009 234 691  
Suite 22, 11 Preston Street  
Como Western Australia 6152  
Tel: (08) 9368 7474 Fax: (08) 9368 7475  
Email: sales@chemag.com.au  
Web: www.chemag.com.au

# DIRECTIONS FOR USE:

## ANCILLARY PANEL

### DIRECTIONS FOR USE:

**RESTRAINTS:** Do NOT apply to very wet soils if rain is imminent or to soils prone to waterlogging.  
Do NOT apply to soils of very high organic matter content.  
Do NOT apply to crops or weeds under stress caused by factors such as root or foliar diseases, nutrient deficiencies, or extremes of temperature or moisture.

Crop	Weeds Controlled	State	Rate per HA	Critical Comments
Pre-emergence  Chickpeas	Deadnettle ( <i>Lamium amplexicaule</i> ), Indian hedge mustard ( <i>Sisymbrium orientale</i> ), white ironweed ( <i>Buglossoides arvensis</i> ), wild radish ( <i>Raphanus raphanistrum</i> ), wireweed ( <i>Polygonum aviculare</i> )	Vic, SA only	45 g plus 1.0-1.5L simazine (500 g/L)	<p>Apply to moist, well prepared, clod and weed-free soil after planting and before crop emergence. Sufficient rainfall is required after application and prior to weed emergence to wet soil to a depth of 5cm.</p> <p>Use the higher rate of simazine on heavier soils, or where higher weed pressure is expected, or where wireweed is a problem.</p> <p>Under adverse conditions, weeds may not be totally controlled but populations will be significantly reduced and surviving plants will generally be severely retarded. Good crop growth will aid weed control.</p> <p>Transient yellowing or reddening of the crop may occur. The risk of crop injury may be increased under adverse growing conditions.</p> <p>Do NOT use this mixture on soils, and in areas, ill suited to growing chickpeas as crop injury will be increased.</p>

Crop	Weeds Controlled	State	Rate per HA	Critical Comments
Pre-emergence  Faba beans, Field peas	Indian hedge mustard ( <i>Sisymbrium orientale</i> ), Shepherd's purse ( <i>Capsella bursa-pastoris</i> ), stinging nettle ( <i>Urtica urens</i> ) (NSW, Vic only), toad rush ( <i>Juncus bufonius</i> ), turnip weed ( <i>Rapistrum rugosum</i> ), * deadnettle ( <i>Lamium amplexicaule</i> ), * doublegee ( <i>Emex australis</i> ), * Paterson's curse ( <i>Echium plantagineum</i> ), * wireweed ( <i>Polygonum aviculare</i> ), ** wile oats ( <i>Avena fatua</i> )	NSW, Vic, SA, WA only	70 g	Apply to moist, well prepared, clod and weed-free soil after planting and before crop emergence. Sufficient rainfall is required after application and prior to weed emergence to wet soil to a depth of 5cm.  Under adverse conditions, weeds may not be totally controlled but populations will be significantly reduced and surviving plants will generally be severely retarded. Good crop growth will aid weed control.  * Surviving plants will generally be retarded and will not compete with the crop. A follow-up spray with another product may be required for control of wild radish under high weed pressure or high rainfall conditions.
	* Wild radish ( <i>Raphanus raphanistrum</i> ), ** annual ryegrass ( <i>Lolium rigidum</i> ), ** barley grass ( <i>Hordeum leporinum</i> )	NSW, WA only		** Competition from grass weeds will be significantly reduced for at least 8 weeks. A post-emergence grass herbicide application may be required. (Refer to COMPATIBILITY section of label).
	Storksbill ( <i>Erodium spp.</i> )	SA, WA only		*** Populations will not be reduced but plants will generally be significantly stunted. Seed set will also be reduced. Use the higher rates under anticipated high weed pressure and in high winter rainfall areas.
	Mouse-ear chickweed ( <i>Cerastium glomeratum</i> ) * capeweed ( <i>Arctotheca calendula</i> ), * prickly lettuce ( <i>Lactuca serriola</i> ), * yellow burweed ( <i>Amsinckia intermedia</i> )	NSW only	70 to 100g	

Crop	Weeds Controlled	State	Rate per HA	Critical Comments
Pre-emergence  Faba beans, Field peas	* Ivy leaf speedwell ( <i>Veronica hederifolia</i> ), *** Bifora ( <i>Bifora testiculata</i> )	SA only	70 to 100g	Do NOT use on faba beans on light, sandy soils.  On alkaline soils, the risk of crop damage to faba beans may be increased under adverse growth conditions.  Refer to the Crop Safety and Follow Crop sections of this label regarding varietal selectivity and follow crops.
	* Threehorn bedstraw ( <i>Galium tricornutum</i> ), * wild radish ( <i>Raphanus raphanistrum</i> ), * yellow burweed ( <i>Amsinckia intermedia</i> )	Vic, SA only		
	* Lesser loosestrife ( <i>Lythrum hyssopifolia</i> ), * prickly lettuce ( <i>Lactuca serriola</i> ), ** annual ryegrass ( <i>Lolium rigidum</i> ), ** barley grass ( <i>Hordeum leporinum</i> )		100 g	
Post-emergence  Field peas (Following varieties only: Alma, Dun, Dundale, Early Dun, Wirrega)	Deadnettle ( <i>Lamium amplexicaule</i> ), Hare's ear ( <i>Conringia orientalis</i> ), Indian hedge mustard ( <i>Sisymbrium orientale</i> ), Toadrush ( <i>Juncus bufonius</i> ), * wireweed ( <i>Polygonum aviculare</i> )	NSW, Vic, SA only	70 g plus a non-ionic surfactant at 200mL /100L	Apply to actively growing weeds in the cotyledon to 3 leaf stage.  Weeds may not be totally controlled but populations will be significantly reduced and surviving plants will generally be severely retarded. Good crop growth will aid weed control.  * Surviving plants will generally be retarded and will not compete with the crop.
	* Threehorn bedstraw ( <i>Galium tricornatum</i> )	Vic, SA only	70 to 100 g plus a non-ionic surfactant at 200mL /100L	Use the higher rate under high weed pressure.  Refer to the Follow Crop section of this label regarding follow crops.

Crop	Weeds Controlled	State	Rate per HA	Critical Comments
<p><b>Pre-emergence</b></p> <p>Mungbeans, peanuts, soybeans.</p>	<p>Common sida (<i>Sida rhombifolia</i>), Fat hen (<i>Chenopodium album</i>), Green amaranth (<i>Amaranthus viridis</i>), Pigweed (<i>Portulaca oleracea</i>), Redroot amaranth (<i>Amaranthus retroflexus</i>), Wild gooseberry (<i>Physalis minima</i>), *Anoda weed (<i>Anoda cristate</i>), * bellvine (<i>Ipomoes plebeia</i>), * common thornapple (<i>Datura stramonium</i>), * deadnettle (<i>Lamium amplexicaule</i>), * jute (<i>Corchorus olitorius</i>), * mintweed (<i>Salvia reflexa</i>) * starburr (<i>Acanthospermum hispidum</i>) * wild radish (<i>Raphanus raphanistrum</i>)</p>	<p>Qld, NSW, Vic, NT only</p>	<p>100 g</p>	<p>Apply to moist, well prepared, clod and weed-free soil after planting and before crop emergence. Sufficient rainfall or irrigation is required after application and prior to weed emergence to wet soil to a depth of 5 cm.</p> <p>Where soil crusting is likely, apply post-emergence. (Not mung beans).</p> <p>Under adverse conditions, weeds may not be totally controlled but populations will be significantly reduced and surviving plants will generally be severely retarded. Good crop growth will aid weed control.</p> <p>* Surviving plants will generally be retarded and will not compete with the crop.</p>
	<p>Above weeds plus Bladder ketmia (<i>Hibiscus trionum</i>), * awnless banyard grass (<i>Echinochloa colona</i>), * Apple of Peru (<i>Nicandra physalodes</i>), * Noogoora burr (<i>Xanthium pungens</i>), * Nutgrass (<i>Cyperus rotundus</i>)</p>		<p>140 g</p>	

Crop	Weeds Controlled	State	Rate per HA	Critical Comments
<b>Post-emergence</b>  Peanuts, Soybeans	Common sida ( <i>Sida rhombifolia</i> ), Wild gooseberry ( <i>Physalis minima</i> ), *Anoda weed ( <i>Anoda cristate</i> ), *Bathurst burr ( <i>Xanthium spinosum</i> ), *bellvine ( <i>Ipomoea plebeia</i> ), *common thornapple ( <i>Datura stramonium</i> ), *deadnettle ( <i>Lamium amplexicaule</i> ), *jute ( <i>Corchorus olitorius</i> ), *fierce thornapple ( <i>Datura ferox</i> )	Qld, NSW, Vic, NT only	100 g plus a non-ionic surfactant at 200mL/100L or Hasten or Kwickin at 500mL/100L	Apply to actively growing weeds in the 2 to 4 leaf stage.  Weeds may not be totally controlled but populations will be significantly reduced and surviving plants will generally be severely retarded. Good crop growth will aid weed control.  *Surviving plants will generally be retarded and will not compete with the crop. For grass weeds, a follow-up spray with a selective grass herbicide may be required.
	Above weeds plus Apple of Peru ( <i>Nicandra physalodes</i> ), Fat hen ( <i>Chenopodium album</i> ), *awnless barnyard grass ( <i>Echinochloa colona</i> ), *barnyard grass ( <i>Echinochloa crus-galli</i> ), *nutgrass ( <i>Cyperus rotundus</i> )		140 g plus a non-ionic surfactant at 200mL/100L or Hasten or Kwickin at 500mL/100L	
<b>Pre-emergence to weeds</b>  Lucerne (established), serradella (established)	As for pre-emergence use in faba beans and field peas (winter weeds) and in soybeans (summer weeds).	Qld, NSW, Vic, SA, WA only	70 to 140 g	Use pre-emergence to weeds in <b>established</b> Lucerne only.  Apply following cutting or grazing, if necessary in mixtures with registered knock-down products.  Apply at rates as per pre-emergence use in faba beans and field peas (winter weeds) and soybeans (summer weeds).  Note <b>CRITICAL COMMENTS</b> applying to weed control in those crops.
		NT only		
Centrosema (Cavalcade) (pre-emergence to crop)				

Crop	Weeds Controlled	State	Rate per HA	Critical Comments
<p><b>Post-emergence</b></p> <p>Lucerne, serradella</p>	<p>As for post-emergence use in field peas (winter weeds) and in soybeans (summer weeds)</p>	<p>Qld, NSW, Vic, SA, WA only</p>	<p>70 to 140 g plus a non-ionic surfactant at 200mL /100L</p>	<p>Apply to actively growing weeds in the cotyledon to 3 leaf stage (winter weeds) and 2-4 leaf stage (summer weeds).</p> <p>Apply at rates as per post-emergence use in field peas (winter weeds) and soybeans (summer weeds).</p> <p>Note <b>CRITICAL COMMENTS</b> applying to weed control in those crops.</p> <p><u>Seedling:</u> Apply when crop is at the first trifoliolate leaf stage or later (Spring sown) and the 2 trifoliolate leaf or later (Autumn sown). Serradella varieties on which ChemAg Amaze has been tested and found to be selective are: Avila, Elgara, Tauro</p> <p><u>Established:</u> Apply as above following cutting or grazing.</p>
<p>Subterranean clover (Trifolium subterraneum)</p>	<p>Doublegee (Emex Australis)</p>	<p>WA only</p>	<p>50 g plus 300mL diuron (500 g/L) plus BOOST* at 2 L per 100 L water</p>	<p>Apply to actively growing weeds up to the 4 leaf stage and when the sub-clover is at the 3 leaf stage or later.</p> <p>This treatment should only be used in the first year of a pasture phase to aid sub-clover establishment.</p> <p>Weed numbers will be reduced and survivors will be stunted. Seed set will also be reduced.</p> <p>Sub-clover biomass may be reduced by this treatment. Varieties on which ChemAg Amaze has been tested are Dalkeith and Nungarin.</p> <p>Other weeds may also be affected, thereby reducing the total pasture biomass.</p>

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIOD:**

DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION.

HARVEST: NOT REQUIRED WHEN USED AS DIRECTED.

**GENERAL INSTRUCTIONS**

This product can be used for either pre- or post-emergence weed control depending on the crop and weeds to be controlled. When the product is applied pre-emergence, susceptible weeds may emerge but growth will be retarded and weeds will either die or remain stunted and will not compete with the crop.

A non-ionic surfactant, or an adjuvant as specified in the DIRECTIONS OF USE, must be added to ChemAg Amaze for post-emergence weed control. The addition of liquid ammonium sulphate at a rate of 2 L/100 L water may assist in post-emergence control of summer growing weeds. Adequate soil moisture is important for optimum activity. When applied post-emergence, weeds will either die or remain stunted and will not compete with the crop.

**MIXING**

This product mixes readily with both hard and soft water. Fill the spray tank one half to three quarters full with clean water and then, with agitator running, add the required amount of this product and then fill the tank with water. When tank mixing this product with other recommended compatible products, first add the other product to the tank and mix thoroughly before adding this product.

**APPLICATION**

ChemAg Amaze should not be applied for a minimum of two hours before rainfall or irrigation.

Ground Application: Avoid overlaps when spraying. Apply in 50 to 100 L/ha water using flat fan nozzles. This product may be applied in a band over the row in row crops. This will assist in minimizing soil residue carry-over.

Aerial Application (Pre-emergence only): Avoid overlaps when spraying. Apply in a minimum swath width of 18 m using a droplet VMD of 230-280 microns.

Do NOT apply ChemAg Amaze under conditions (e.g. dead calm, excessive wind and/or small droplets) likely to cause spray drift onto wetlands or waterways, natural vegetation, crops other than specified on label or land to be planted with susceptible crops.

**EQUIPMENT CLEAN-UP**

Thoroughly flush all spray equipment with water following use of ChemAg Amaze and before use with other products.

Rinse water should NOT be discharged where it will reach streams, water bodies or natural vegetation.

## COMPATIBILITY

This product is compatible with Cyclone, Dictate Duo, trifluralin, metribuzin, cyanazine, diuron, glyphosate, paraquat plus diquat, simazine, diflufenican, dimethoate, omethoate, and endosulfan.

Do NOT tank mix with selective post-emergence grass herbicides. Do NOT apply these herbicides following use of ChemAg Amaze until grasses have resumed active growth.

## RESISTANT WEEDS WARNING

**GROUP B HERBICIDE**

ChemAg Amaze Herbicide is a member of the Imidazolinone group of herbicides. ChemAg Amaze has the inhibition of acetolactate synthase (ALS) mode of action. For weed resistance management, ChemAg Amaze is a Group B Herbicide. Some naturally-occurring weed biotypes resistant to ChemAg Amaze and other Group B herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These weeds will not be controlled by ChemAg Amaze or other Group B herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, ChemAg Pty Ltd accepts no liability for any losses that may result from failure of ChemAg Amaze to control resistant weeds.

## PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Do NOT spray drift to contact desirable vegetation or crops other than specified on label.

Do NOT spray within 50 m of wetlands or waterways.

**Crop Safety:** This product may cause slight shortening of plant internodes and may in some circumstances lead to transient crop yellowing but plants soon recover and yield is unaffected. This effect may be more pronounced when the product is used post-emergence or under poor growth conditions.

Do NOT use this product pre-emergence on Collegian and Cressy Blue field pea varieties.

Do NOT use this product post-emergence on field pea varieties other than Alma, Dun, Dundale, Early Dun and Wirrega.

Do NOT use pre-emergence on serradella, seeding Lucerne or subterranean clover.

Do NOT use post-emergence on chickpeas, faba beans or mung beans.

Should re-sowing of chickpeas, faba beans, field peas, Lucerne, mung beans, peanuts, serradella, soybeans or subterranean clover be necessary, do NOT reapply ChemAg Amaze.

**Follow Crops:** Under conditions which do not favour breakdown of this product, carry-over soil residues can affect susceptible follow crops. As environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product, rotational crop injury is always possible.

The following minimum re-cropping intervals (months after application) should be observed.

## Following use in winter crops:-

MONTHS AFTER APPLICATION			
0	10	22	34
Maize varieties with CLEARFIELD Technology <b>ONLY</b> ; -Pacific Hycorn 62IT -Pacific Hycorn 53IT -Pioneer 3395IR;	Lucern; Lupins; Pasture legumes; Vetch; *triticale *Barley;	Oats; Safflower	All other crops including canola (Except varieties with CLEARFIELD Technology; see 0 months
Wheat varieties with CLEARFIELD Technology <b>ONLY</b> ; Canola varieties with CLEARFIELD Technology <b>ONLY</b> ; Faba beans; Field peas; Chickpeas	*Wheat (except varieties with CLEARFIELD Technology; see 0 months)		

\*The following additional requirements apply if it is intended to sow WHEAT (except varieties with CLEARFIELD Technology), BARLEY or TRITICALE during the next winter season.

- Do NOT apply ChemAg Amaze pre-emergence later than the end of June and post-emergence later than the end of July.
- Do NOT use ChemAg Amaze in areas where rainfall from spraying to sowing of cereals is expected to be below 300mm.

### Furthermore:

- In SA and WA do NOT use on soils of pH 5.5 (Ca Cl<sub>2</sub>) or less in areas where rainfall from spraying to sowing of cereals is expected to be below 400mm.
- In NSW, Vic and SA do NOT use the 100g/ha rate in areas where rainfall from spraying to sowing of cereals is expected to be below 400mm.

If expected rainfall is not received following the use of ChemAg Amaze, consult your local ChemAg representative before planting wheat, barley or triticale. (In calculating rainfall actually received, exclude single isolated heavy summer and autumn falls about 100mm.)

## Following use in summer crops:-

### Irrigated only:

MONTHS AFTER APPLICATION			
0	5	10	18
Maize varieties with CLEARFIELD Technology <b>ONLY</b> : -Pacific Hycorn 62IT -Pacific Hycorn 53IT -Pioneer 33951R; mung beans; peanuts; soybeans	Chickpeas; Lucerne; Lupins; Pasture legumes; * barley; * wheat (except for wheat varieties with CLEARFIELD Technology); canola varieties with CLEARFIELD Technology <b>ONLY</b>	** maize (except for varieties with CLEARFIELD Technology; see 0 months); ** sorghum	All other crops (providing rainfall and irrigation exceeds 2000mm)

\* Do NOT plant these crops unless interim moisture (rainfall plus irrigation) from application to sowing is at least 500mm.

\*\* Do NOT plant these crops unless interim moisture (rainfall plus irrigation) from application to sowing is at least 800mm. Do NOT plant sorghum if ChemAg Amaze rates higher than 100 g/ha were used in the previous crop.

### Dry land only:

Do NOT use the 140g rate in dryland soybeans, mung beans or peanuts unless it is intended to recrop with a leguminous crop or crop varieties with CLEARFIELD Technology.

MONTHS AFTER APPLICATION				
0	10	15	22	27
Maize varieties with CLEARFIELD Technology <b>ONLY</b> : -Pacific Hycorn 62IT -Pacific Hycorn 53IT -Pioneer 33951R; mung beans; peanuts; soybeans	** maize (except varieties with CLEARFIELD Technology); ** sorghum	Chickpeas; Lucerne; Lupins; Pasture legumes; * barley; * wheat (except for wheat varieties with CLEARFIELD Technology); canola varieties with CLEARFIELD Technology <b>ONLY</b>	Cotton; Maize (see also 0 months); Sorghum; sunflower	All other crops (providing rainfall exceeds 2000mm)

\* Do NOT plant these crops unless interim rainfall from application to sowing is at least 500mm.

\*\* Do NOT plant these crops unless interim rainfall from application to sowing is at least 800mm.

Do NOT plant sorghum if Amaze rates higher than 100 g/ha were used in the previous crop.

## **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

Do NOT contaminate dams, waterways or drains with this product or used containers.

## **STORAGE AND DISPOSAL**

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point.

If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If not available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should NOT be burnt.

## **SAFETY DIRECTIONS**

Will damage eyes. Will irritate skin. Avoid contact with eyes and skin. When preparing spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

## **FIRST AID**

If poisoning occurs, contact a doctor or Poison's Information Centre. Telephone 13 11 26 Australia wide.

## **MSDS**

Additional information is listed in the Material Safety Data Sheet.

## **CONDITIONS OF SALE**

All conditions and warranties rights and remedies implied by law or arising in contract or tort whether due to the negligence of ChemAg Pty Ltd or otherwise are hereby expressly excluded so far as the same may be legally be done provided however that any rights of the Buyer pursuant to non excludable conditions or warranties of the Trade Practices act 1974 or any relevant legislation of any State are expressly preserved but the liability of ChemAg Pty Ltd or any intermediate Seller pursuant thereto shall be limited if so permitted by the said legislation to the replacement of goods sold or the supply of equivalent goods and all liability for indirect or consequential loss or damage of whatsoever nature is expressly excluded. This product must be used or applied strictly in accordance with the instructions appearing hereon. This product is solely sold for use in Australia and must not be exported without the prior written consent of ChemAg Pty Ltd.

FOR SPECIALISTS ADVICE IN AN **EMERGENCY ONLY PHONE 1800 033 111**TOLL FREE ALL HOURS AUSTRALIA WIDE

APVMA Approval Number: 58746/1004