


STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsing to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Harmful if swallowed. Will damage eyes. Will irritate the skin. Avoid contact with eyes and skin. Do not inhale vapour. When preparing spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist and washable hat, elbow-length PVC gloves, face shield and goggles. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.



CAUTION

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

MADDOG SELECTIVE HERBICIDE

ACTIVE CONSTITUENT:

110 g/L FENOXAPROP-P-ETHYL

CROP SAFENER: 30 g/L MEFENPYR-DIETHYL

SOLVENT: 594 g/L HYDROCARBON LIQUID

GROUP A HERBICIDE

For the selective control of wild oats and annual phalaris in wheat, triticale and cereal rye crops.

IMPORTANT:

READ THE LEAFLET THOROUGHLY BEFORE USE.

Batch No:

Date of Manufacture:

Imtrade Australia Pty Ltd ABN 13 090 151 134

17 Ocean Street, Kwinana WA 6167

Tel: (08) 9419 0333 Fax: (08) 9419 7516

www.imtrade.com.au



FIRST AID

IF poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26). If swallowed **DO NOT** induce vomiting. Give a glass of water.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet, which can be obtained from Imtrade Australia Pty Ltd. www.imtrade.com.au

CONDITIONS OF SALE

This product must be used strictly as directed and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so Imtrade Australia Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such direction and instructions.

**In a Transport Emergency Dial 000
Police or Fire Brigade**

CONTENTS 1L
APVMA Approval No: 61502/1/0108

CONTENTS 5L
APVMA Approval No: 61502/5/0108

GENERAL INSTRUCTIONS

Results are best under good growing conditions. Application to weeds or crop under stress (eg. Due to continuous severe frosts, nutrient deficient, and diseased, dry or waterlogged conditions) should be avoided. Generally the best yield advantage will come from controlling weeds as early as possible.

RESISTANT WEEDS WARNING

GROUP **A** HERBICIDE

ChemAg Maddog Selective Herbicide is a member of the aryloxyphenoxypropionate group of herbicides. ChemAg Maddog Selective Herbicide is a herbicide, which inhibits production of acetyl-coA carboxylase. For weed resistance management ChemAg Maddog Selective Herbicide is a Group **A** herbicide. Some naturally occurring weed biotypes resistant to ChemAg Maddog Selective Herbicide, and other Group **A** herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by ChemAg Maddog Selective Herbicide or other Group **A** herbicides.

Since occurrence of resistant weeds is difficult to detect prior to use. Imtrade Australia accepts no liability for any losses that may result from the failure of ChemAg Maddog Selective Herbicide to control resistant weeds.

DO NOT make more than one application of a Group **A** Herbicide to a crop per season. For further information on herbicide resistance management consult an Imtrade Australia Representative, trained agricultural consultant or local department agronomist.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to the aquatic environment. **DO NOT** contaminate streams, rivers, or waterways with this product or the used container.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply in weather conditions or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

DO NOT plant susceptible crops (eg corn, sorghum, oats or barley) for weeks after application.

Under sown Clovers and Medics ChemAg Maddog Selective Herbicide does not affect under sown clovers or medics or other broadleaf crops.

Imtrade Australia Pty Ltd
ABN 13 090 151 134
17 Ocean Street,
Kwinana WA 6167
Tel: (08) 9419 0333
Fax: (08) 9419 7516
www.imtrade.com.au

CAUTION

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**



MADDOG SELECTIVE HERBICIDE

ACTIVE CONSTITUENT:

110 g/L FENOXAPROP-P-ETHYL

CROP SAFENER:

30 g/L MEFENPYR-DIETHYL

SOLVENT:

594 g/L HYDROCARBON LIQUID

GROUP **A** HERBICIDE

**For the selective control of wild oats
and annual phalaris in wheat, triticale
and cereal rye crops.**

**IMPORTANT: READ THE LEAFLET
THOROUGHLY BEFORE USE.**

Batch No:

Date of Manufacture:



CONTENTS 10L
APVMA Approval No: 61502/10/0108

CONTENTS 20L
APVMA Approval No: 61502/20/0108

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsing to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Harmful if swallowed. Will damage eyes. Will irritate the skin. Avoid contact with eyes and skin. Do not inhale vapour. When preparing spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist and washable hat, elbow-length PVC gloves, face shield and goggles. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

IF poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26). If swallowed **DO NOT** induce vomiting. Give a glass of water.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet, which can be obtained from Imtrade Australia Pty Ltd. www.imtrade.com.au

CONDITIONS OF SALE

This product must be used strictly as directed and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so Imtrade Australia Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such direction and instructions.

In a Transport Emergency Dial

000

Police or Fire Brigade

CAUTION

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

ChemAg MADDOG SELECTIVE HERBICIDE

ACTIVE CONSTITUENT:

110 g/L FENOXAPROP-P-ETHYL

CROP SAFENER: 30 g/L MEFENPYR-DIETHYL

SOLVENT: 594 g/L HYDROCARBON LIQUID

GROUP	A	HERBICIDE
--------------	----------	------------------

For the selective control of wild oats and annual phalaris in wheat, triticale and cereal rye crops.

**IMPORTANT:
READ THIS LEAFLET THOROUGHLY BEFORE USE.**

APVMA Approval No: 61502/0108

Imtrade Australia Pty Ltd
ABN 13 090 151 134
17 Ocean Street, Kwinana WA 6167
Tel: (08) 9419 0333 Fax: (08) 9419 7516
www.imtrade.com.au

DIRECTIONS FOR USE**RESTRAINTS**

DO NOT apply if rainfall is expected in 4 hours.

DO NOT apply to weeds or crop under stress due to, for example, very dry, very wet, nutrient deficient, and frosty or diseased conditions.

CROP	WEED	RATE (mL/ha)	CRITICAL COMMENTS
Wheat, Triticale, Cereal Rye.	Wild Oats (<i>Avena spp.</i>)	300	Apply 3 to 4 weeks after sowing when cereal has 2 to 5 true leaves (Z12 to Z15). Do not use this rate when wild oats are past the four-leaf stage (Z14) or when there are more than 200 plants per square metre. Add BS 1000® or an equivalent adjuvant at 250 mL/100 L water.
		350	Apply generally 3 to 6 weeks after sowing when cereal is usually 2 leaves to early tillering (Z12 – Z22). Aim to apply actively growing young weeds up to 5 total leaves. (1 Tiller) (Z14, Z21). Add BS 1000® or an equivalent adjuvant at 250 mL/100 L water.
		400	Apply up to approximately 10 weeks after sowing to actively growing weeds up to mid tillering (Z22 – Z24). Add BS 1000® or an equivalent adjuvant at 250 mL/100 L water.
	Annual phalaris (<i>Phalaris minor, P. paradoxa</i>)	400	Apply generally 3 to 4 weeks after sowing when cereal usually has 2 to 5 true leaves (Z12 – Z15). Apply only to weed densities below 200 plants per square metre. Aim to apply actively growing young weeds up to 4 total leaves (Z14). Add BS 1000® or an equivalent adjuvant at 250 mL/100 L water.
		500	Apply generally 4 to 6 weeks after sowing when cereal has usually reached early tillering (Z21 – Z22). Aim to apply to actively growing weeds up to 5 total leaves (1 tiller) (Z14, Z21). Add BS 1000® or an equivalent adjuvant at 250 mL/100 L water.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

Harvest - DO NOT HARVEST FOR 10 WEEKS AFTER APPLICATION

Grazing - DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 3 WEEKS AFTER APPLICATION.

GENERAL INSTRUCTIONS

Results are best under good growing conditions. Application to weeds or crop under stress (eg. Due to continuous severe frosts, nutrient deficient, and diseased, dry or waterlogged conditions) should be avoided. Generally the best yield advantage will come from controlling weeds as early as possible.

RESISTANT WEEDS WARNING

GROUP

A

HERBICIDE

ChemAg Maddog Selective Herbicide is a member of the aryloxyphenoxypropionate group of herbicides. ChemAg Maddog Selective Herbicide is a herbicide, which inhibits production of acetyl-coA carboxylase. For weed resistance management ChemAg Maddog Selective Herbicide is a Group **A** herbicide. Some naturally occurring weed biotypes resistant to ChemAg Maddog Selective Herbicide, and other Group **A** herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by ChemAg Maddog Selective Herbicide or other Group **A** herbicides.

Since occurrence of resistant weeds is difficult to detect prior to use, Imtrade Australia accepts no liability for any losses that may result from the failure of ChemAg Maddog Selective Herbicide to control resistant weeds.

DO NOT make more than one application of a Group **A** Herbicide to a crop per season. For further information on herbicide resistance management consult an Imtrade Australia Representative, trained agricultural consultant or local department agronomist.

Export of Treated Produce

Growers should note that suitable MRLs or import tolerance might not be established in all markets for produce treated with ChemAg Maddog Selective Herbicide. IF you are growing produce for export, please check with Imtrade Australia Pty Ltd for the latest information on MRLs and import tolerance BEFORE using ChemAg Maddog Selective Herbicide.

Application

Aim for complete coverage of weeds with ChemAg Maddog Selective Herbicide. Follow directions under "Equipment".

Mixing

Add the required quantity of ChemAg Maddog Selective Herbicide directly to the spray tank when about two-thirds full. Top up with water to the required volume, with agitators in motion, after adding broadleaf herbicides if required.

Equipment

Ground Sprayers – Standard boom sprayers only are recommended and must be fitted with by-pass or mechanical agitation. It is recommended to use approximately 50 to 100 L of water/ha and a droplet size of 200 to 300 microns. **DO NOT** apply with boom less jets or misters.

Aircraft – ChemAg Maddog Selective Herbicide may be applied by aircraft. Best results are obtained using 25 to 30 L of water/ha with steady crosswind. Do not exaggerate the swath width. Aim for a droplet size of 200 to 300 microns.

Compatibility

Mixtures of ChemAg Maddog Selective Herbicide with broadleaf herbicides may lead to a reduction in grass weed control. This reduction can be minimised by closely following all use recommendations and restrictions. Use wetting agent in any of the tank mixtures described below at the rate prescribed in the Directions for Use on this label. The following table summarises the current knowledge on compatibility:

	Wild Oats	Annual Phalaris	
Hoegrass#	Green	Green	
Tigrex# up to 0.8 L/ha	Yellow	Red	
Jaguar# up to 0.75 L/ha	Yellow	Red	
Eclipse# up to 7 g/ha	Green	Yellow	
Broadstrike# up to 25 g/ha	Green	Yellow	
MCPA LVE (500 g/L) up to 700 mL/ha	Yellow	Red	
Bromoxynil + MCPA up to 1.4 L/ha	Red	Red	
Glean# up to 15 g/ha	Yellow	Red	
Lontrel# up to 150 mL/ha	Green	Red	
Ally#	Yellow	Red	
Bromoxynil	Red	Red	
Harmony# M up to 30 g/ha	Green	White	
Eclipse up to 5 g/ha	+ Lontrel up to 100 mL/ha	Green	White
	+ Bromoxynil + MCPA up to 350 mL/ha	Yellow	White
	+ Bromoxynil + MCPA up to 350 mL/ha + Lontrel up to 100 mL/ha	Yellow	White
Tordon# 242 up to 1 L/ha	Yellow	White	

Green Trials indicate no reduction in grass weed control when used as recommended.

Results at below recommended rates may indicate some reduction in some cases.

These mixtures can be used with minimal loss of activity under good growing conditions with all label recommendations followed.

Yellow Trials indicate some reduction in grass weed control at recommended rates, even under good growing conditions.

These mixtures will usually result in reduced grass weed control and cannot be recommended unless such efficacy loss is accepted. These mixtures have demonstrated transient crop yellowing which does not significantly reduce yield.

Red Trails at recommended rates indicate a severe reduction in grass weed control and/or significant crop injury.

White No data available.

Allow at least 10 days between any application of ChemAg Maddog Selective Herbicide and herbicides containing 2,4-D, MCPA, Dicamba or other similar sprays, which should preferably follow ChemAg Maddog Selective Herbicide.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to the aquatic environment. **DO NOT** contaminate streams, rivers, or waterways with this product or the used container.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply in weather conditions or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

DO NOT plant susceptible crops (eg corn, sorghum, oats or barley) for weeks after application.

Under sown Clovers and Medics – ChemAg Maddog Selective Herbicide does not affect under sown clovers or medics or other broadleaf crops.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsing to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury

empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Harmful if swallowed. Will damage eyes. Will irritate the skin. Avoid contact with eyes and skin. Do not inhale vapour. When preparing spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist and washable hat, elbow-length PVC gloves, face shield and goggles. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

IF poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26). If swallowed **DO NOT** induce vomiting. Give a glass of water.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet, which can be obtained from Imtrade Australia Pty Ltd. www.imtrade.com.au

CONDITIONS OF SALE

This product must be used strictly as directed and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so Imtrade Australia Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such direction and instructions.

Other Trademarked products

**In a Transport Emergency Dial
000
Police or Fire Brigade**