

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CHEMAG
MCPA 500
SELECTIVE HERBICIDE

Active Constituent: 500g/L MCPA
PRESENT AS THE DIMETHYLAMINE SALT

GROUP I HERBICIDE

For the selective control of broadleaf weeds in
cereals, linseed, pastures, sugar cane and turf
as per Directions For Use table.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT.

Contents:5/20/200/500/1000 Litres

ChemAg Pty. Ltd. ACN 009 234 691
Suite 12, 11 Preston Street Como Western Australia 6152
Tel: (08) 9367 1799 Fax: (08) 9368 6144

PROTECTION OF CROPS, NATIVE AND OTHER NON TARGET PLANTS:

DO NOT spray in high winds. DO NOT spray crops or weeds outside the stages indicated in "Critical Comments" as damage, loss of yield or inadequate weed control may result.

Drift Warning: DO NOT apply under meteorological conditions or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. Avoid spray drift onto susceptible crops such as cotton, tobacco, tomatoes, vines, lupins, fruit trees, ornamentals, and susceptible trees (eg. Kurrajongs, Belahs).

Legume Tolerance: The tolerance of pasture legumes to this product can vary with species, variety, growing conditions, stage of growth and companion crop cover. While MCPA is preferable to 2,4-D in most situations, this product will cause greater damage than MCPA amine formulations. Lucerne and medics should not be sprayed under any circumstances with this product, as severe damage will result.

NB: The tolerance of pasture legume to applied product can vary with species, variety, growing conditions, stage of growth and companion crop cover. DO NOT spray cereal crops undersown with lucerne as damage may result.

PROTECTION OF LIVESTOCK:

Low hazard to bees. May be applied at any time as recommended in the Directions For Use.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIROMENT:

DO NOT contaminate streams, rivers of water with the chemicals or used container.

STORAGE AND DISPOSAL:

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling break, crush, or puncture and bury containers at a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

For REFILLABLE containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS:

Harmful if swallowed. Attaches eyes. Will irritate the skin. When opening the container, preparing the spray and using product wear elbow-length PVC gloves and goggles, cotton overalls buttoned to the neck and wrist and a washable hat. If product on skin immediately wash area with soap and water. If in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing

FIRST AID:

If poisoning occurs, contact a Doctor or Poisons Information Center (Phone 131126). If swallowed, and if more than 15minutes from a hospital induce vomiting, preferably using Ipecac Syrup APF.

MATERIAL SAFETY DATA SHEET:

Additional information is listed in the Material Safety Data Sheet (MSDS).

GENERAL INSTRUCTIONS:

Weeds should be sprayed while actively growing and at their most susceptible stage. As a general guide, annuals should be sprayed when young and perennials just prior to appearance of bud stage. Extremes of cold or drought are unfavorable. The best conditions are when the soil is moist, weather fine and rain unlikely within 6 hours of application.

RESISTANT WEED WARNING: CHEMAG MCPA 500 SELECTIVE HERBICIDE (MCPA 500) is a member of the Phenoxys group of herbicides. MCPA 500 has the Disrupters of plant cell growth mode of action. For weed resistance management MCPA 500 is a Group I herbicide.

Some naturally - occurring weed biotypes resistant to MCPA 500 and other disrupters of plant cell growth herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by MCPA 500 or other disrupters of plant cell growth herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, ChemAg Pty Ltd. accepts no liability for any losses that may result from the failure of MCPA 500 to control resistant weeds.

APPLICATION/MIXING INFORMATION: This product mixes readily with water. Half fill the spray tank with clean water and add the required amount of product. Agitate thoroughly before adding remainder of water.

BOOM SPRAYING - Use 30-120L/ha of water.

-Use minimum of 170 litres water per hectare for LINSEED

AERIAL SPRAYING - Use 10-90L/ha of water.

EQUIPMENT MAINTENANCE & USAGE: Equipment that has been used for this chemical should not be used for the application of other materials to sensitive plants, unless it has been cleaned with Spraymate Tank & Equipment Cleaner or well washed out with a 2% solution of cloudy ammonia followed by several clear water rinses

COMPATIBILITY/TANK MIXES: This product can be tank mixed with Ally ®, Banvel ®, Bromicide ®, Flowable Diuron, Flowable Terbutryne, Lontrel ®, Sertin ®, Sprayseed ®, Tordon ® and Tribunil®. The effectiveness of Imidan ®, Le Mat®, Metasystox® and Dimethoate may be reduced by alkaline hydrolysis if tank mixed with this product. For these situations the addition of Spraymate LI-700 is recommended. When tank mixing add the required amount of compatible product (and surfactant if recommended) when the spray tank is nearly full and while continuously agitating. If left overnight, agitate thoroughly before spraying.

CONDITIONS OF SALE:

“ChemAg Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on ChemAg’s skill of judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of ChemAg has any authority to add to or alter these conditions”.

NRA Approval No. 51691/0700

In case of emergency: Phone 000 Police or Fire Brigade

CHEMAG
MCPA 500
SELECTIVE HERBICIDE

Active Constituent: 500g/L MCPA
PRESENT AS THE DIMETHYLAMINE SALT

GROUP I HERBICIDE

For the selective control of
broadleaf weeds in cereals, linseed,
pastures, sugar cane and turf
as per Directions For Use table.

READ THOROUGHLY BEFORE USING

DIRECTIONS FOR USE:**DO NOT apply if crop or weeds are stressed due to dry or excessively moist conditions.****DO NOT apply if rain is likely within 6 hours.****1.FIELD CROPS**

Situation & Crop	Weeds Controlled	State	Rate/ha	Critical Comments
Wheat, Barley, Oats Triticale	Refer to Weed Table	S.A, Tas. only	700ml to 2.1L	Apply through the tillering stage after the crop plants have 5 leaves and before they commence jointing.
		Qld. only		Apply from early tillering (4 to 5 fully emerged main stem leaves plus one or more tillers) until before the boot stage (visible swelling of the head at the top of the main stem).
		N.S.W. only		Apply 700ml rate after crop has reached 5 leaf stage and weeds marked E in weed table have 3-6 leaves. Higher rates after the first node can be felt in a tiller.
		Vic. Only	700ml to 1.5L	Apply 700ml rate after crop has reached 2-3 leaf stage and weeds marked E in weed table have 2-4 leaves. Higher rate after the crop plants have 5 leaves to fully tillered.
		W.A. only	900ml to 2L	Apply to crop from 5 leaf to flag emergence (Z 15-37)
Wheat, Barley, Triticale, Cereal, Rye	Capeweed, Doublegee, Erodium, Mustard, Rapistrum, Wild Turnip, Wild Radish	W.A. only	400ml plus 350ml Nufarm Flowable Diuron	CROP: 3-4 leaf stage (Z 13-14)
			500ml plus 400ml Nufarm Flowable Diuron	CROP: 4-5 leaf stage (Z 14-15)
				Apply by aircraft or boom with 20-100L/ha water, when most weeds have germinated and are in 2-5 leaf stage. Crop and weeds should be dry at the time of application. Some temporary yellowing of crop may occur after application. Undersown sub-clovers may be slightly retarded. DO NOT apply to undersown medics.
	Amsinckia(Yellow Burr), Capeweed, Charlock, Deadnettle, Doublegee (Spiny Emex, Three corner Jack), Erodium sp., Mustard, Turnip Weed, White Ironweed (Corn Gromwell, Sheepweed), Wild Radish, Wild Turnip.	N.S.W., Vic. Only	500ml plus 500ml Nufarm Flowable Diuron	Apply by boom spray or aircraft when the crop is at the 3-5 leaf stage (N.S.W and S.A. only) and 5 leaf stage (Vic only). Weeds must be in the 2-5 leaf stage. Do not Apply to undersown medic or lucerne. Do not use on Kamilaroi, Kitr, Olympic or Shortim Wheat.
Capeweed, Mustard, Three Cornered Jack, Turnip Weed, Wild Radish, Wild Turnip, Wards Weed, Common Ice Plant, Pimpernel	S.A. only	350ml plus 500ml Nufarm Flowable Diuron	Do not use on very light sandy soils.	

FIELD CROPS Continued

Situation & Crop	Weeds Controlled	State	Rate/ha	Critical Comments
Wheat, Barley, Oats, Triticale	Soldier Thistle	S.A. only	700ml plus 200ml Lontrel	Apply to crop at tillering stage after the crop has 5 leaves and before the crop commences jointing.
	Skeleton Weed		700ml plus 500ml Lontrel	Apply to rosettes from 5-15cm diameter when the crop is between the 5 leaf and late tillering growth stages. It is essential that all the skeleton weed is emerged and minimum of 5cm diameter, which is usually not before August. This rate will provide control of skeleton weed until harvest and will substantially reduce seed numbers the following season
Linseed	Refer to Weed Table	Qld. only N.S.W., Vic., S.A., Tas. Only	700ml to 2.1L 700ml to 1 L	Apply when crop is 10-15cm tall with at least 170L/ha of water. Do not spray after buds appear. Some wilting effect may be seen following application but crop usually recovers quickly. Do not apply by aircraft.
Sugar Cane	Blue Top, Chinese Burr, Flannel Weed, Gambia Pea, Bell Vine, Streaked Rattle Pod, Bindweed, Pink Convolvulus, Cupids Flower, Merremia Vine, Morning Glory.	Qld. only	1.4L	POST - Emergent: Apply as directed inter-row spray. Do not exceed rates with phenoxy sensitive varieties.
	As above plus Fat Hen, Noogoora Burr		2.1L	
	All of the above		10L	
Flax	Refer to Weed Table	Vic., Tas. Only	490mL	Apply to crop when 8-20cm high.

2. PASTURES, SEED CROPS & TURF

Situation & Crop	Weeds Controlled	State	Rate/ha	Critical Comments
Grass Pastures	Refer to Weed Table	All States	700ml to 4L	Apply to established grass based pastures. Damage may occur to Legumes if present. See "Legumes Tolerance" below. NOTE: Capeweed, Paterson's Curse and Variegated Thistle may be poisonous to stock after spraying
	Salvation Jane	S.A. only	1L plus 400ml Flowable Diuron	Use this mixture only if spraying is delayed until early winter. Rosettes should not exceed 20cm in diameter.
	Soldier Thistle	S.A. only	700ml to 2.1L	Use low rate on seedlings with no more than 2 true leaves. Use 100 litres of water for best results. If clover present do not exceed 600ml

2. PASTURES, SEED CROPS & TURF Continued

Situation & Crop	Weeds Controlled	State	Rate/ha	Critical Comments
Grass Pastures	Seedling and established Horehound	N.S.W. only	3L	Apply in autumn when horehound plants are at the seedling stage. If sub clover or medic plants are present they will be damaged by this application.
Grass Seed Crops	Refer to Weed Table	Qld., S.A., Tas., Vic., W.A. only	700ml - 4L	Apply to established grass seed crops.
Sub-clover Seed Crops	Refer to Weed Table	Vic., S.A. only	700ml - 1.5L	Do not exceed rate specified. Spray in Autumn & early Winter.
Pastures - Spray/Graze Technique	Caltrop, Capeweed, Charlock, Mustards, Paterson's Curse, Shepherd's Purse, Saffron, Slender, Spear or Variegated Thistle, Turnip Weed, Wild Radish, Wild Turnip	N.S.W., Vic only Tas. Only	350ml to 1.4L 700ml	Spray actively growing 6-8 weeks old weeds. Introduce stock 7-10 days after spraying, preferably sheep as cattle are less effective. Use stocking rate at least 5 times heavier than normal until weeds have been reduced but before survival of desirable pasture species is threatened. Lucerne and medics may be damaged and should be grazed short before spraying. Refer "Legume Tolerance" Precaution: An increased quantity of poisonous plants may be eaten by stock using Spray/Graze, e.g. Caltrop, Capeweed, Paterson's Curse and Variegated Thistle and deaths could result from causes such as nitrate poisoning. With Paterson's Curse preferably graze stock soon destined for slaughter and avoid extended periods of grazing. Avoid grazing with young or breeding stock. Do not graze horses or pigs on Paterson's Curse. Observe Withholding Period of at least 7 days after spraying.
	Refer to Weed Table	W.A. only S.A. only	925ml to 1.85L 500ml	
	Capeweed, Dandelion, Seedling Dock, Three Cornered Jack, Geranium, Erodium, Mustard, Annual Thistles, Turnip Weed, Wild Radish, Wild Turnip			
	Soldiers Thistle		600ml	
Turf	Refer to Weed Table	All States	500ml or 5ml per 100sq. meters	Newly laid - Delay until well established or at least two months old. Apply in high volume to actively growing weeds. See "Legume Tolerance". Buffalo grass is more easily damaged than other grasses
			1.4 to 2.8L or 14 to 28ml per 100 sq. meters	Established - Apply in high volume to actively growing weeds. Do not mow for 2 days before application. Some transitory damage may occur to fine turf grasses and strawberry clover. See "Legume Tolerance". Buffalo grass is more easily damaged than other grasses.

3. SPOT SPRAYING

Situation & Crop	Weeds Controlled	State	Mixing / Critical Comments
High Volume Spraying	Refer to Weed Table	All States	Add 1/10 th of rate on weed table to 150litres of water. Each 150 litres of mix will cover 1000m ² (1/10 th Ha). E.g. If rate in weed table is 1.4L use 140ml/150L water
Knapsack Application			Add 1/100 th of rate on weed table to 10 litres of water. Each 10 litres of mix will cover 100m ² (1/100 th Ha) e.g. If rate in weed table is 1.4L use 14ml/10L water

4. NON CROP AREAS, FALLOW LAND, INDUSTRIAL & COMMERCIAL AREAS, RIGHTS-OF-WAY

Situation & Crop	Weeds Controlled	State	Rate/ha	Critical Comments
Non Crop Areas, Fallow Land, Industrial and Commercial Areas, Rights-of-Way	Refer to Weed Table	All States	Refer to Weed Table	Apply at rates shown in Weed Table when weeds only are present and no crop or pasture is involved.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD: CEREALS, PASTURE - DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION

WEED TABLE:

Where weeds are to be sprayed in a crop or pasture, use only the rate given for particular crop or situation indicated under the Directions For Use table. In most cases this will give control, however some hard to kill weeds or those in advanced stages of growth may only be suppressed.

The rates listed in the Weed Table below are for use where weeds only are present and no crop or pasture is involved. Use at these rates in a crop or pasture other than for spot spraying may cause damage.

This product should only be used in those States where a rate or range of rates is indicated for the particular weeds listed.

APPLICATION RATE PER HECTARE							
WEEDS CONTROLLED	VIC	N.S.W	S.A.	Tas.	Qld.	W.A.	CRITICAL COMMENTS
Ball Mustard			1.7L				Apply at rosette stage
Bathurst Burr	2L	1-2L	2L	2L	2L	2L	Spray young seedlings only
Black Bindweed		1.4-2L			1.4-2L		Apply at young stage prior to flowering
California Burr		1-2L		2.2L			Spray young seedlings only
Cape Tulip						2L	Apply just prior to flowering. Gives mild suppression only.
Capeweed		2.1L	2.1L	2.1L	2.1L		Apply at seedling stage
Carrot Weed	2L				2L		Apply at seedling stage
Charlock	700ml	1L	700ml	1-2.2L	700ml		Spray up to rosette stage
Cutleaf Mignonette			2.1L				Apply at young rosette stage
Dandelion	2L	2L	2L	2L	2L	2L	Apply at young rosette stage
Deadnettle		2.1L			2.1L		Moderately susceptible. Apply at seedling stage
Docks	1.5L		2.1L	2.1L			Apply at rosette stage and before flower heads emerge. Only the top growth of old established plants will be affected.
Fat hen	2L	1-2L		2L	2L		Spray up to rosette stage
Fennel	2.7L	2.7L	2.7L	2.7L	2.7L		Apply at young stage
Field Bindweed	2L		2L	2L	2L	2L	Apply at bud stage, repeating over several seasons
Field Cress				2.2L			Apply at young stage
Flatweed or Catsear				2.2L		1.4L	Apply when weeds are 10-15 cm high
Fumitory (Red)	1.4L	1.4L	1.4L		1.4L		Apply at young stage prior to flowering
Hoary Cress	4.5L		2.1L		4.5L		Apply before flowering
Horehound		3L	2.1L				Apply at rosette stage
Hedge Mustard	1.1L		700ml	700ml	700ml	1.25L	Apply at rosette stage
Hexham Scent or Melilotus	1.5L	1.2L	1.4L		1.4L		Apply at young seedling stage
Lesser Swinecress				2.2L			Apply at young stage

WEED TABLE CONT...

APPLICATION RATE PER HECTARE							
WEEDS CONTROLLED	VIC	N.S.W	S.A.	Tas.	Qld.	W.A.	CRITICAL COMMENTS
Lincoln Weed			700ml				Apply at rosette stage
London Rocket						1.25L	
Lupins		700ml-1.5L				1.25L	Spray up to 10cm high
Mintweed	2L	2L			2L		Spray young seedlings only
Mustards	1.1L	1L	1.1L	1.1L		1.25L	Spray up to rosette stage
Noogoora Burr	2L	1-2L	2L		2L	2L	Spray young seedlings only
Opium Poppy				2.2L			Apply at young stage
Paterson's Curse		1-1.5L			4L		Apply early rosette stage
Pimpernel			700ml				Apply at rosette stage
Plantains		2L		2L	2L		Apply at rosette stage
Rapeseed		1-1.5L					Spray up to rosette stage
Rough Poppy		700ml-1.5L					Spray up to rosette stage
Safflower		700ml-1L					Spray up to rosette stage
Shepherd's Purse				2.2L			Apply at rosette stage
Skeleton Weed		1.5-2L					Spray rosettes before aerial growth commences
Sorrel			2.1L				Apply to rosette stage
Stinkwort	2L	1-1.5L	2L			2L	Apply when weeds are 10-15cm high
Sunflower (seedlings)		1-1.5L					Spray up to rosette stage
Thistles							
- Artichoke	1.6L		2.1L		2L	700ml	Apply at rosette stage
- Nodding		2.5L					Spray early rosettes. Repeat spray required in following spring
- Saffron	1.5-2.5L	1-2L	1.4-2.5L	1.4-2.5L	1.4-2.5L	1.5-2.5L	Spray up to rosette stage (September WA)
- Scotch or Cotton	2.2L		700ml-1.4L	2.2L			Apply at seedling stage
- Slender	3L	1-2L	1L	3L		3L	Spray young seedlings only
- Soldier	1.1L		1.1L				Spray young seedlings. Use higher rate for larger weeds
- Spear		1.5-2L		2.2L			Apply at rosette stage
- Star			1 - 1.4L				Apply at seedling stage. Use higher rate for larger weeds
- Variegated	1.5L	1-2L	700ml-1.4L	1.5L	1.5L	1.5L	Spray young seedlings only. Use 2L/ha for well developed weeds.

WEED TABLE CONT...

WEEDS CONTROLLED	VIC	N.S.W	S.A.	Tas.	Qld.	W.A.	CRITICAL COMMENTS
Turnip Weed	1.1L	1L	1L				Spray up to rosette stage
Wild Artichoke			2.1L				Apply at rosette stage
Wild Radish	1.4L	1L	1.4L	2L	700ml-1.1L	1.2L	Spray up to rosette stage
Wild Turnip	1.1L	1L	700ml	2L	700ml-1.1L	900ml	Spray up to rosette stage

GENERAL INSTRUCTIONS:

Weeds should be sprayed while actively growing and at their most susceptible stage. As a general guide, annuals should be sprayed when young and perennials just prior to appearance of bud stage. Extremes of cold or drought are unfavorable. The best conditions are when the soil is moist, weather fine and rain unlikely within 6 hours of application.

RESISTANT WEED WARNING: CHEMAG MCPA 500 SELECTIVE HERBICIDE (MCPA 500) is a member of the Phenoxy group of herbicides. MCPA 500 has the Disrupters of plant cell growth mode of action. For weed resistance management MCPA 500 is a Group I herbicide.

Some naturally - occurring weed biotypes resistant to MCPA 500 and other disrupters of plant cell growth herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by MCPA 500 or other disrupters of plant cell growth herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, ChemAg Pty Ltd. accepts no liability for any losses that may result from the failure of MCPA 500 to control resistant weeds.

APPLICATION/MIXING INFORMATION: This product mixes readily with water. Half fill the spray tank with clean water and add the required amount of product. Agitate thoroughly before adding remainder of water.

BOOM SPRAYING - Use 30-120L/ha of water.

-Use minimum of 170 litres water per hectare for LINSEED

AERIAL SPRAYING - Use 10-90L/ha of water.

EQUIPMENT MAINTENANCE & USAGE: Equipment that has been used for this chemical should not be used for the application of other materials to sensitive plants, unless it has been cleaned with Spraymate Tank & Equipment Cleaner or well washed out with a 2% solution of cloudy ammonia followed by several clear water rinses

COMPATIBILITY/TANK MIXES: This product can be tank mixed with Ally®, Banvel®, Bromicide®, Flowable Diuron, Flowable Terbutryne, Lontrel®, Sertin®, Sprayseed®, Tordon® and Tribunil®. The effectiveness of Imidan®, Le Mat®, Metasystox® and Dimethoate may be reduced by alkaline hydrolysis if tank mixed with this product. For these situations the addition of Spraymate LI-700 is recommended. When tank mixing add the required amount of

compatible product (and surfactant if recommended) when the spray tank is nearly full and while continuously agitating. If left overnight, agitate thoroughly before spraying.

PROTECTION OF CROPS, NATIVE AND OTHER NON TARGET PLANTS: DO NOT spray in high winds. DO NOT spray crops or weeds outside the stage indicated in 'Critical Comments' as damage, loss of yield or inadequate weed control may result.

Drift warning: DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. Avoid spray drift and Vapor movement on susceptible crops such as cotton, vines, tomatoes, vegetables, fruit trees, Lucerne, legume crops and pastures, oilseed crops or other susceptible crops or trees (e.g. Kurrajongs, Belahs, Eucalypts).

Legume Tolerance: The use of MCPA is preferable to 2,4-D Amine (Amicide) or Ester (Estercide) in situations where legumes are undersown in the crop. N.B.: The tolerance of pasture legumes to applied product can vary with species, variety, growing conditions, stage of growth and companion crop cover. DO NOT spray cereal crops undersown with lucernes as damage may result.

PROTECTION OF LIVESTOCK: Low hazard to bees. May be applied at any time as recommended in the Directions For Use.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIROMENT: DO NOT contaminate streams, rivers of water with the chemicals or used container.

STORAGE AND DISPOSAL: Store in the closed, original container in a dry, cool, well ventilated area away from children, animals, food, foodstuff, seeds and fertilizers. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse container before disposal. Add rinsing to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

Returnable containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS: Harmful if swallowed. Attaches eyes. Will irritate the skin. When opening the container, preparing the spray and using product wear elbow-length PVC gloves and goggles, cotton overalls buttoned to the neck and wrist and a washable hat. If product on skin immediately wash area with soap and water. If in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID: If poisoning occurs, contact a Doctor or Poisons Information Centre (Phone 131126). If swallowed, and if more than 15minutes from a hospital induce vomiting, preferably using Ipecac Syrup APF.

For further information refer to the Material Safety Data Sheet (MSDS)

CONDITIONS OF SALE: “ChemAg Limited (ChemAg) shall not be liable for any loss injury or damage whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on ChemAg’s skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of ChemAg has any authority to add to or alter these conditions.”

NRA Approval No: 51691/0700