

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Will damage eyes. Will irritate skin. Avoid contact with eyes and skin. When preparing spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (*Phone Australia 13 11 26*). If in eyes, hold eyes open, flood with water for at least 15 minutes and see doctor.

Refer to attached Leaflet for additional GHS
Hazard & Precautionary Statements

IMTRADE

AMAZE 700 WG

HERBICIDE

ACTIVE CONSTITUENT: 700 g/kg IMAZETHAPYR

GROUP 2 HERBICIDE

For the pre- or post-emergence control of certain weeds in Centrosema (Cavalcade), Chickpeas, Faba Beans, Field Peas, Lucerne, Mung Beans, Peanuts, Serradella, Soybeans and Subterranean Clover as per the Directions for Use table.

IMPORTANT: Read the attached Leaflet before use.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from your supplier or Imtrade Australia Pty Ltd website at www.imtrade.com.au

CONDITIONS OF SALE

Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Imtrade Australia Pty Ltd has any authority to add to or alter these conditions.

In a Transport Emergency Dial 000
Police or Fire Brigade

APVMA Approval No: 63390/102504

GROUP
2

FORM
WG

CONTENTS
250g-8kg



Batch No.
DOM:

IMTRADE
CropScience

17 Ocean St, Kwinana Beach WA 6167
08 9419 0333 | www.imtrade.com.au

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



IMTRADE AMAZE 700 WG HERBICIDE

ACTIVE CONSTITUENT: 700 g/kg IMAZETHAPYR

GROUP	2	HERBICIDE
--------------	----------	------------------

For the pre- or post-emergence control of certain weeds in Centrosema (Cavalcade), Chickpeas, Faba Beans, Field Peas, Lucerne, Mung Beans, Peanuts, Serradella, Soybeans and Subterranean Clover as per the Directions for Use table

**IMPORTANT: READ THIS LEAFLET THOROUGHLY
BEFORE OPENING OR USING THIS PRODUCT**

APVMA Approval No: 63390/102504

Imtrade Australia Pty Ltd
ABN 13 090 151 134
17 Ocean Street, Kwinana WA 6167
Tel: 1800 171 799 Fax: 1800 171 788
Web: www.imtrade.com.au

RESTRAINTS

DO NOT apply to very wet soils if rain is imminent or to soils prone to waterlogging.

DO NOT apply to soils of very high organic matter content.

DO NOT apply to crops or weeds under stress caused by factors such as root or foliar diseases, nutrient deficiencies, or extremes of temperature or moisture.

DIRECTIONS FOR USE

HIGH VOLUME SPRAYING

Crop	Weeds Controlled	State	Rate/ha	Critical Comments
Pre-emergence Chickpeas	Deadnettle (<i>Lamium amplexicaule</i>), Indian hedge mustard (<i>Sisymbrium orientale</i>), White ironweed (<i>Buglossoides arvensis</i>), Wild radish (<i>Raphanus raphanistrum</i>), Wireweed (<i>Polygonum aviculare</i>)	Vic, SA only	45g plus 1.0-1.5L Flowable Simazine (500 g/L)	Apply to moist, well prepared, clod and weedfree soil after planting and before crop emergence. Sufficient rainfall is required after application and prior to weed emergence to wet soil to a depth of 5cm. Use the higher rate of simazine on heavier soils, or where higher weed pressure is expected, or where wireweed is a problem. Under adverse conditions, weeds may not be totally controlled but populations will be significantly reduced and surviving plants will generally be severely retarded. Good crop growth will aid weed control. Transient yellowing or reddening of the crop may occur. The risk of crop injury may be increased under adverse growing conditions. DO NOT use this mixture on soils, and in areas, ill- suited to growing chickpeas as crop injury will be increased.

Crop	Weeds Controlled	State	Rate/ha	Critical Comments
Pre-emergence Faba beans, Field peas	Indian hedge mustard (<i>Sisymbrium orientale</i>), Shepherds purse (<i>Capsella bursa-pastoris</i>), Stinging nettle (<i>Urtica urens</i>) (NSW, Vic only), Toad rush (<i>Juncus bufonius</i>), Turnip weed (<i>Rapistrum rugosum</i>), * Deadnettle (<i>Lamium amplexicaule</i>), * Doublegee (<i>Emex australis</i>), * Patersons curse (<i>Echium plantagineum</i>), * Wireweed (<i>Polygonum aviculare</i>), ** Wild oats (<i>Avena fatua</i>)	NSW, Vic, SA, WA only	70g	<p>Apply to moist, well prepared, clod and weed-free soil after planting and before crop emergence. Sufficient rainfall is required after application and prior to weed emergence to wet soil to a depth of 5cm.</p> <p>Under adverse conditions, weeds may not be totally controlled but populations will be significantly reduced and surviving plants will generally be severely retarded. Good crop growth will aid weed control.</p> <p>* Surviving plants will generally be retarded and will not compete with the crop. A follow-up spray with another product may be required for control of wild radish under high weed pressure or high rainfall conditions.</p> <p>** Competition from grass weeds will be significantly reduced for at least 8 weeks. A post-emergence grass herbicide application may be required. (Refer to COMPATIBILITY section of label).</p>
	* Wild radish (<i>Raphanus raphanistrum</i>), ** Annual ryegrass (<i>Lolium rigidum</i>), ** Barley grass (<i>Hordeum leporinum</i>)	NSW, WA only		

continued next page ...

Crop	Weeds Controlled	State	Rate/ha	Critical Comments
Pre-emergence Faba beans, Field peas (contd)	Storksbill (<i>Erodium</i> spp.)	SA, WA only	70g	... continued from previous page *** Populations will not be reduced but plants will generally be significantly stunted. Seed set will also be reduced. Use the higher rates under anticipated high weed pressure and in high winter rainfall areas.
	Mouse-ear chickweed (<i>Cerastium glomeratum</i>) * Capeweed (<i>Arctotheca calendula</i>), * Prickly lettuce (<i>Lactuca serriola</i>), * Yellow burrweed (<i>Amsinckia intermedia</i>)	NSW only	70 to 100g	DO NOT use on faba beans on light, sandy soils. On alkaline soils, the risk of crop damage to faba beans may be increased under adverse growth conditions.
	* Ivy leaf speedwell (<i>Veronica hederifolia</i>), *** Bifora (<i>Bifora testiculata</i>)	SA only		Refer to the Crop Safety and Follow Crop sections of this label regarding varietal selectivity and follow crops.
	* Threehorn bedstraw (<i>Galium tricornutum</i>), * Wild radish (<i>Raphanus raphanistrum</i>), * Yellow burrweed (<i>Amsinckia intermedia</i>)	Vic, SA only		
	* Lesser loosestrife (<i>Lythrum hyssopifolia</i>), * Prickly lettuce (<i>Lactuca serriola</i>), ** Annual ryegrass (<i>Lolium rigidum</i>), ** Barley grass (<i>Hordeum leporinum</i>)			100g

Crop	Weeds Controlled	State	Rate/ha	Critical Comments
Post-emergence Field peas (Following varieties only: Alma, Dun, Dundale, Early Dun, Wirrega)	Deadnettle (<i>Lamium amplexicaule</i>), Hares ear (<i>Conringia orientalis</i>), Indian hedge mustard (<i>Sisymbrium orientale</i>), Toadrush (<i>Juncus bufonius</i>), * Wireweed (<i>Polygonum aviculare</i>)	NSW, Vic, SA only	70g plus a non-ionic surfactant at 200mL/100L	Apply to actively growing weeds in the cotyledon to 3 leaf stage. Weeds may not be totally controlled but populations will be significantly reduced and surviving plants will generally be severely retarded. Good crop growth will aid weed control. * Surviving plants will generally be retarded and will not compete with the crop.
	* Threehorn bedstraw (<i>Galium tricornatum</i>)	Vic, SA only	70 to 100g plus a non-ionic surfactant at 200mL/100L	Use the higher rate under high weed pressure. Refer to the Follow Crop section of this label regarding follow crops.

Crop	Weeds Controlled	State	Rate/ha	Critical Comments
Pre-emergence Mungbeans, Peanuts, Soybeans	Common sida (<i>Sida rhombifolia</i>), Fat hen (<i>Chenopodium album</i>), Green amaranth (<i>Amaranthus viridis</i>), Pigweed (<i>Portulaca oleracea</i>), Redroot amaranth (<i>Amaranthus retroflexus</i>), Wild gooseberry (<i>Physalis minima</i>), * Anoda weed (<i>Anoda cristate</i>), * Bellvine (<i>Ipomoes plebeia</i>), * Common thornapple (<i>Datura stramonium</i>), * Deadnettle (<i>Lamium amplexicaule</i>), * Jute (<i>Corchorus olitorius</i>), * Mintweed (<i>Salvia reflexa</i>) * Starburr (<i>Acanthospermum hispidum</i>) * Wild radish (<i>Raphanus raphanistrum</i>)	Qld, NSW, Vic, NT only	100g	Apply to moist, well prepared, clod and weed-free soil after planting and before crop emergence. Sufficient rainfall or irrigation is required after application and prior to weed emergence to wet soil to a depth of 5cm. Where soil crusting is likely, apply post-emergence. (Not mungbeans). Under adverse conditions, weeds may not be totally controlled but populations will be significantly reduced and surviving plants will generally be severely retarded. Good crop growth will aid weed control. * Surviving plants will generally be retarded and will not compete with the crop.
	Above weeds plus Bladder ketmia (<i>Hibiscus trionum</i>), * Awnless barnyard grass (<i>Echinochloa colona</i>), * Apple of Peru (<i>Nicandra physalodes</i>), * Noogoora burr (<i>Xanthium pungens</i>), * Nutgrass (<i>Cyperus rotundus</i>)		140g	

Crop	Weeds Controlled	State	Rate/ha	Critical Comments
Post-emergence Peanuts, Soybeans	Common sida (<i>Sida rhombifolia</i>), Wild gooseberry (<i>Physalis minima</i>), * Anoda weed (<i>Anoda cristate</i>), * Bathurst burr (<i>Xanthium spinosum</i>), * Bellvine (<i>Ipomoea plebeia</i>), * Common thornapple (<i>Datura stramonium</i>), * Deadnettle (<i>Lamium amplexicaule</i>), * Jute (<i>Corchorus olitorius</i>), * Fierce thornapple (<i>Datura ferox</i>)	Qld, NSW, Vic, NT only	100g plus a non-ionic surfactant at 200mL/100L or Hasten® or Kwickin® at 500mL/ 100L	Apply to actively growing weeds in the 2 to 4 leaf stage. Weeds may not be totally controlled but populations will be significantly reduced and surviving plants will generally be severely retarded. Good crop growth will aid weed control. *Surviving plants will generally be retarded and will not compete with the crop. For grass weeds, a follow-up spray with a selective grass herbicide may be required.
	Above weeds plus Apple of Peru (<i>Nicandra physalodes</i>), Fat hen (<i>Chenopodium album</i>), * Awnless barnyard grass (<i>Echinochloa colona</i>), * Barnyard grass (<i>Echinochloa crusgalli</i>), * Nutgrass (<i>Cyperus rotundus</i>)		140g plus a non-ionic surfactant at 200mL/100L or Hasten® or Kwickin® at 500mL/ 100L	

Crop	Weeds Controlled	State	Rate/ha	Critical Comments
<p>Pre-emergence to weeds</p> <p>Lucerne (established), Serradella (established)</p>	<p>As for pre-emergence use in faba beans and field peas (winter weeds) and in soybeans (summer weeds).</p>	<p>Qld, NSW, Vic, SA, WA only</p>	<p>70 to 140g</p>	<p>Use pre-emergence to weeds in established Lucerne only.</p> <p>Apply following cutting or grazing, if necessary in mixtures with registered knock-down products.</p> <p>Apply at rates as per preemergence use in faba beans and field peas (winter weeds) and soybeans (summer weeds).</p> <p>Note CRITICAL COMMENTS applying to weed control in those crops.</p>
<p><i>Centrosema</i> (Cavalcade) (pre-emergence to crop)</p>		<p>NT only</p>		

Crop	Weeds Controlled	State	Rate/ha	Critical Comments
<p>Post-emergence</p> <p>Lucerne, Serradella</p>	<p>As for post-emergence use in field peas (winter weeds) and in soybeans (summer weeds)</p>	<p>Qld, NSW, Vic, SA, WA only</p>	<p>70 to 140g plus a non-ionic surfactant at 200ml/100L</p>	<p>Apply to actively growing weeds in the cotyledon to 3 leaf stage (winter weeds) and 2-4 leaf stage (summer weeds).</p> <p>Apply at rates as per post-emergence use in field peas (winter weeds) and soybeans (summer weeds).</p> <p>Note CRITICAL COMMENTS applying to weed control in those crops.</p> <p>Seedling: Apply when crop is at the first trifoliate leaf stage or later (Spring sown) and the 2 trifoliate leaf or later (Autumn sown). Serradella varieties on which Imtrade Amaze 700 WG Herbicide has been tested and found to be selective are: Avila, Elgara, Tauro</p> <p>Established: Apply as above following cutting or grazing.</p>

Crop	Weeds Controlled	State	Rate/ha	Critical Comments
Subterranean clover (<i>Trifolium subterraneum</i>)	Doublegee (<i>Emex australis</i>)	WA only	50g+ 300mL diuron (500 g/L) + BOOST at 2L/100L water	<p>Apply to actively growing weeds up to the 4-leaf stage and when the sub-clover is at the 3-leaf stage or later.</p> <p>This treatment should only be used in the first year of a pasture phase to aid sub-clover establishment.</p> <p>Weed numbers will be reduced and survivors will be stunted. Seed sat will also be reduced.</p> <p>Sub-clover biomass may be reduced by this treatment. Varieties on which Imtrade Amaze 700 WG Herbicide has been tested are Dalkeith and Nungarin.</p> <p>Other weeds may also be affected, thereby reducing the total pasture biomass.</p>

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD:

GRAZING: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION.

HARVEST: NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS

This product can be used for either pre- or post-emergence weed control depending on the crop and weeds to be controlled. When the product is applied preemergence, susceptible weeds may emerge but growth will be retarded and weeds will either die or remain stunted and will not compete with the crop.

A non-ionic surfactant, or an adjuvant as specified in the **DIRECTIONS OF USE**, must be added to Imtrade Amaze 700 WG Herbicide for post-emergence weed control. The addition of liquid ammonium sulphate at a rate of 2L/100L water may assist in post-emergence control of summer growing weeds. Adequate soil moisture is important for optimum activity. When applied post-emergence, weeds will either die or remain stunted and will not compete with the crop.

MIXING

This product mixes readily with both hard and soft water. Fill the spray tank one half to three quarters full with clean water and then, with agitator running, add the required amount of this product and then fill the tank with water. When tank mixing this product with other recommended compatible products, first add the other product to the tank and mix thoroughly before adding this product.

APPLICATION

Imtrade Amaze 700 WG Herbicide should not be applied for a minimum of two hours before rainfall or irrigation.

Ground Application: Avoid overlaps when spraying. Apply in 50 to 100L/ha water using flat fan nozzles. This product may be applied in a band over the row in row crops. This will assist in minimizing soil residue carry-over.

Aerial Application (Pre-emergence only): Avoid overlaps when spraying. Apply in a minimum of 20L/ha water with a maximum swath width of 18m using a droplet VMD or 230-280 microns.

DO NOT apply Imtrade Amaze 700 WG Herbicide under conditions (e.g. dead calm, excessive wind and/or small droplets) likely to cause spray drift onto wetlands or waterways, natural vegetation, crops other than specified on label or land to be planted with susceptible crops.

EQUIPMENT CLEAN-UP

Thoroughly flush all spray equipment with water following use of Imtrade Amaze 700 WG Herbicide and before use with other products.

Rinse water should **NOT** be discharged where it will reach streams, water bodies or natural vegetation.

COMPATIBILITY

This product is compatible with alpha-cypermethrin duo, pendimethalin, trifluralin, metribuzin, cyanazine, diuron, glyphosate, paraquat plus diquat, diquat, simazine, diflufenican, dimethoate and omethoate.

DO NOT tank mix with selective post-emergence grass herbicides. **DO NOT** apply these herbicides following use of Imtrade Amaze 700 WG Herbicide until grasses have resumed active growth.

RESISTANT WEEDS WARNING

GROUP 2 HERBICIDE

Imtrade Amaze 700 WG Herbicide is a member of the Imidazolinone group of herbicides. Imtrade Amaze 700 WG Herbicide has the inhibition of acetolactate synthase (ALS) mode of action. For weed resistance management, Imtrade Amaze 700 WG Herbicide is a Group 2 Herbicide. Some naturally occurring weed biotypes resistant to Imtrade Amaze 700 WG Herbicide and other Group 2 Herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These weeds will not be controlled by Imtrade Amaze 700 WG Herbicide or other Group 2 Herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Imtrade Australia Pty Ltd accepts no liability for any losses that may result from failure of Imtrade Amaze 700 WG Herbicide to control resistant weeds.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

DO NOT spray within 50m of wetlands or waterways.

Crop Safety: This product may cause slight shortening of plant internodes and may in some circumstances lead to transient crop yellowing but plants soon recover and yield is unaffected. This effect may be more pronounced when the product is used post-emergence or under poor growth conditions.

DO NOT use this product pre-emergence on Collegian and Cressy Blue field pea varieties.

DO NOT use this product post-emergence on field pea varieties other than Alma, Dun, Dundale, Early Dun and Wirrega.

DO NOT use pre-emergence on serradella, seedling Lucerne or subterranean clover.

DO NOT use post-emergence on chickpeas, faba beans or mung beans.

Should re-sowing of chickpeas, faba beans, field peas, Lucerne, mung beans, peanuts, serradella, soybeans or subterranean clover be necessary, **DO NOT** reapply Imtrade Imazethapyr 700 WG Herbicide.

Follow Crops: Under conditions which do not favour breakdown of this product, carry-over soil residues can affect susceptible follow crops. As environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product, rotational crop injury is always possible.

The following minimum re-cropping intervals (months after application) should be observed.

Following use in Winter Crops:

MONTHS AFTER APPLICATION			
0	10	22	34
Maize varieties with CLEARFIELD Technology ONLY : -Pacific Hycom 621T -Pacific Hycom 531T -Pioneer 33951R; Wheat varieties with CLEARFIELD Technology ONLY ; Canola varieties with CLEARFIELD Technology ONLY ; Faba beans; Field peas; Chickpeas	Lucerne; Lupins; Pasture legumes; Vetch; *Triticale *Barley; *Wheat (except varieties with CLEARFIELD Technology; see 0 months)	Oats; Safflower	All other crops including canola (except varieties with CLEARFIELD Technology; see 0 months)

* The following additional requirements apply if it is intended to sow WHEAT (except varieties with CLEARFIELD Technology), BARLEY or TRITICALE during the next winter season.

- **DO NOT** apply Imtrade Amaze 700 WG Herbicide pre-emergence later than the end of June and post-emergence later than the end of July.
- **DO NOT** use Imtrade Amaze 700 WG Herbicide in areas where rainfall from spraying to sowing of cereals is expected to be below 300mm.
- In SA and WA **DO NOT** use on soils of pH 5.5 (Ca Cl₂) or less in areas where rainfall from spraying to sowing of cereals is expected to be below 400mm.
- In NSW, Vic and SA **DO NOT** use the 100g/ha rate in areas where rainfall from spraying to sowing of cereals is expected to be below 400mm.

If expected rainfall is not received following the use of Imtrade Amaze 700 WG Herbicide, consult your local Imtrade representative before planting wheat, barley or triticale. (In calculating rainfall actually received, exclude single isolated heavy summer and autumn falls above 100mm.)

**Following use in summer crops:
Irrigated only:**

MONTHS AFTER APPLICATION			
0	5	10	18
Maize varieties with CLEARFIELD Technology ONLY : -Pacific Hycorn 621T -Pacific Hycorn 531T -Pioneer 33951R; mung beans; peanuts; soybeans	Chickpeas; Lucerne; Lupins; Pasture legumes; * barley; * wheat (except for varieties with CLEARFIELD Technology); canola varieties with CLEARFIELD Technology ONLY	** maize (except for varieties with CLEARFIELD Technology; see 0 months); ** sorghum	All other crops (providing rainfall and irrigation exceeds 200mm)

* **DO NOT** plant these crops unless interim moisture (rainfall plus irrigation) from application to sowing is at least 500mm.

** **DO NOT** plant these crops unless interim moisture (rainfall plus irrigation) from application to sowing is at least 800mm. **DO NOT** plant sorghum if Imtrade Amaze 700 WG Herbicide rates higher than 100 g/ha were used in the previous crop.

Dry land only:

MONTHS AFTER APPLICATION				
0	10	15	22	22
Maize varieties with CLEARFIELD Technology ONLY : -Pacific Hycorn 621T -Pacific Hycorn 531T -Pioneer 33951R; mung beans; peanuts; soybeans	** maize (except varieties with CLEARFIELD Technology; see 0 months); ** sorghum	Chickpeas; Lucerne; Lupins; Pasture legumes; * barley; * wheat (except varieties with CLEARFIELD Technology); Canola varieties with CLEARFIELD Technology ONLY	Cotton; Maize (see also 0 months); Sorghum; sunflower	All other crops (providing rainfall exceeds 200mm)

* **DO NOT** plant these crops unless interim rainfall from application to sowing is at least 500mm.

** **DO NOT** plant these crops unless interim rainfall from application to sowing is at least 800mm. **DO NOT** plant sorghum if Imazethapyr rates higher than 100 g/ha were used in the previous crop.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate dams, waterways or drains with this product or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Empty containers and product should **NOT** be burnt.

SAFETY DIRECTIONS

Will damage eyes. Will irritate skin. Avoid contact with eyes and skin. When preparing spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. If product is in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26). If in eyes, hold eyes open, flood with water for at least 15 minutes and see doctor.

Additional GHS Hazard & Precautionary Statements

- **DO NOT** get on clothing;
- **IF SWALLOWED:** Rinse mouth. Do **NOT** induce vomiting;
- **IF ON SKIN:** Wash with plenty of soap and water;
- If skin irritation occurs: Get medical advice;
- If eye irritation persists: Get medical advice;
- In case of fire, use carbon dioxide, dry chemical, foam, water fog;
- Store in a dry place.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from your supplier or Imtrade Australia Pty Ltd website at www.imtrade.com.au

CONDITIONS OF SALE

Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Imtrade Australia Pty Ltd has any authority to add to or alter these conditions.

**In a Transport Emergency
Dial 000
Police or Fire Brigade**