

#### STORAGE AND DISPOSAL (10L & 20L only)

Store product in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling break, crush, or puncture and bury containers at a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burnt.

#### STORAGE AND DISPOSAL (60L only)

Store product in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. If tamper evident seals are broken prior to initial use then the integrity of the contents cannot be assured. Empty container by pumping through dry-break connection system. **DO NOT** attempt to breach the valve system or the filling point, or contaminate the container with water or other products. Ensure that the coupler, pump, meter and hoses are disconnected, triple rinsed and drained after each use. When empty, or contents no longer required, return the 60L container only to the point of purchase. This container remains the property of Imtrade Australia Pty Ltd.

#### SAFETY DIRECTIONS

Harmful if swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. Repeated exposure may cause allergic disorders. When preparing product for use, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

#### FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Phone Australia 13 11 26).

## CAUTION

**KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**



# IMTRADE ANNIHILATE 200 SC ALL PURPOSE INSECTICIDE

**ACTIVE CONSTITUENT: 200 g/L IMIDACLOPRID**

**GROUP 4A HERBICIDE**

**For the control of:**

- 1) Insect pests on cotton, fruit, vegetables, ornamentals, shrub trees in home and commercial use.**
- 2) Larvae of African Black Beetle, Argentinian Scarab, Pruinose Scarab and Bill Bug in Turfgrass as specified in the Directions for Use table**

**IMPORTANT: READ THIS LABEL AND THE ATTACHED LEAFLET  
THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT**

**NET CONTENTS:  
200mL-60L**

Imtrade Australia Pty Ltd  
ABN 13 090 151 134  
17 Ocean Street, Kwinana WA 6167  
Tel: 1800 171 799 Fax: 1800 171 788  
Web: [www.imtrade.com.au](http://www.imtrade.com.au)

#### Additional GHS Hazard & Precautionary Statements

•Avoid breathing fumes, mists vapours or spray; **•DO NOT** get on clothing; •Contaminated work clothing should not be allowed out of the workplace; **•IF SWALLOWED:** Rinse mouth. **DO NOT** induce vomiting; **•IF ON SKIN:** Wash with plenty of soap and water; **•IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing; •If skin irritation occurs: Get medical advice; •If eye irritation persists: Get medical advice; •Not combustible. Use extinguishing media suited to burning materials; •Store in a dry place.

#### SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet which can be obtained from your supplier or Imtrade Australia Pty Ltd website at [www.imtrade.com.au](http://www.imtrade.com.au)

#### CONDITIONS OF SALE

Imtrade Australia Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgement in purchasing or using same and every person dealing with this product does so at his/hers own risk. No representative of Imtrade has any authority to add to or alter these conditions.

**APVMA Approval No. 60465/106758**

Batch No:

Date of Manufacture:

# **CAUTION**

**KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**



## **IMTRADE ANNIHILATE 200 SC ALL PURPOSE INSECTICIDE**

**ACTIVE CONSTITUENT: 200 g/L IMIDACLOPRID**

<b>GROUP</b>	<b>4A</b>	<b>INSECTICIDE</b>
--------------	-----------	--------------------

**For the control of:**

- 1) Insect pests on cotton, fruit, vegetables, ornamentals, shrub trees in home and commercial use.**
- 2) Larvae of African Black Beetle, Argentinian Scarab, Pruinose Scarab and Bill Bug in Turfgrass as specified in the DIRECTIONS FOR USE table.**

**IMPORTANT: READ THIS LEAFLET THOROUGHLY  
BEFORE OPENING OR USING THIS PRODUCT**

**APVMA Approval No. 60465/106758**

Imtrade Australia Pty Ltd  
ABN 13 090 151 134  
17 Ocean Street, Kwinana WA 6157  
**Tel: 1800 171 799 Fax: 1800 171 788**  
Web: [www.imtrade.com.au](http://www.imtrade.com.au)

# **DIRECTIONS FOR USE**

## **FOLIAR SPRAY APPLICATIONS**

<b>CROP</b>	<b>PEST</b>	<b>RATE</b>	<b>WHP</b>	<b>CRITICAL COMMENTS</b>
Cotton	Aphids	<p>250 mL/ha + Pulse™ Penetrate at 0.2% v/v (2 mL/L water) or equivalent organosilicone surfactant* (e.g. Freeway Gold™)</p> <p>* Check with Imtrade for recommended organosilicone surfactants</p>	13 weeks	<p>The addition of Pulse™ Penetrate or equivalent is critical for the performance of Imtrade Annihilate 200 SC All Purpose Insecticide. Apply early in the establishment of an aphid infestation when numbers are low (ie no more than 1 or 2 leaves per plant with honeydew present). Applications made later than this may result in reduced control.</p> <p>Shorter residual control may be evident and a repeat application of a registered aphicide (follow the Cotton Insecticide Resistance Management Strategy for cotton aphid) may be required to achieve complete control:</p> <ul style="list-style-type: none"> <li>• if applications of Imtrade Annihilate 200 SC All Purpose Insecticide plus Pulse™ or equivalent are timed too late (see above); or</li> <li>• if existing high density aphid colonies (hotspots) are present; or</li> <li>• if aphids have established throughout the plant canopy (especially lower in the canopy); or</li> <li>• if there is high reinfestation pressure; or</li> <li>• if there is rapid crop growth; or</li> <li>• if Imtrade Annihilate 200 SC All Purpose Insecticide plus Pulse™ or equivalent is used following a spray-failure (eg resistance to organophosphate or carbamate insecticides).</li> </ul> <p><i>Continued next page</i></p>

## FOLIAR SPRAY APPLICATIONS (continued)

CROP	PEST	RATE	WHP	CRITICAL COMMENTS
Cotton	Aphids	250 mL/ha + Pulse™ Penetrate at 0.2% v/v (2 mL/L water) or equivalent organosilicone surfactant* (e.g. Freeway Gold™)  * Check with Imtrade for recommended organosilicone surfactants	13 weeks	<i>Continued from previous page</i>  Note: Where resistance to carbamates or organophosphates is suspected, Imtrade Annihilate 200 SC All Purpose Insecticide plus Pulse" or equivalent should be used first so as not to delay control of the aphids present.  Aphids treated with Imtrade Annihilate 200 SC All Purpose Insecticide plus Pulse™ or equivalent may still be present on the plant but will not be feeding. Control of aphids should initially be assessed by a reduction in fresh honeydew and not on the presence of aphids on the plant. After ingesting Imtrade Annihilate 200 SC All Purpose Insecticide, aphids may take up to 5 days to die.
	Mirids, Brown Flea Beetle			Apply when pest number reach treatment threshold levels as determined by field checks.
Stone Fruit	Green Peach Aphid, Black Peach Aphid	<b>Dilute spraying</b> 25 mL/100L <b>Concentrate spraying</b> Refer to the Mixing/ Application section	21 days	Apply at first sign of aphid infestation. Apply as a full cover spray, ensuring thorough coverage. Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. <b>DO NOT</b> use in equipment that requires rates greater than 125 mL/100L of water (i.e. greater than 5 x concentrate).

## FOLIAR SPRAY APPLICATIONS (continued)

CROP	PEST	RATE	WHP	CRITICAL COMMENTS
Cucurbits	Green Peach Aphid	25 mL/100L or 300 mL/ha	1 day	Apply at first sign of aphid infestation.
Capsicum, Egg Plant, Potato			7 days	
Tomato			3 days	
Brassicas	Grey Cabbage Aphid, Turnip Aphid		7 days	Apply at first sign of aphid infestation. Add a wetting agent.
Sweet Potato	Silverleaf Whitefly including Type B	25 mL/100L or 250 mL/ha	7 days	Apply at first sign of whitefly or melon thrips infestation. Apply dilute sprays (25 mL/100L) to run off. Ensure thorough coverage of underside of leaves. Use of droppers will improve coverage of underside of leaves.
Cucumber			1 day	
Egg Plant	Melon Thrips		7 days	
Roses	Aphids	25 mL/100L	-	Apply as a thorough cover spray at first sign of insect infestation.
Ornamental Plants	Aphids, Azalea Lace Bug, Bronze Orange Bug, Harlequin Bug, Citrus Mealybug, Greenhouse Thrips, Fullers Rose Weevil			

## FOLIAR SPRAY APPLICATIONS (continued)

CROP	PEST	RATE	WHP	CRITICAL COMMENTS
Ornamental Plants	Hibiscus Flower Beetle	50 mL/100L		Spray buds and flowers as needed.
	Longtailed Mealybug	50 mL/100L + surfactant		Apply 3 sprays 2 weeks apart. Use a non-ionic surfactant at label rate.
	Psyllids	25 mL/100L		Spray at first sign and then a week later.
	Soft Scales	25 mL/100L		Spray in late spring or when small scales are first seen. Apply 3 sprays 2 weeks apart. Add a wetting agent.
Duboisia	Green Peach Aphid	25 mL/100L		Apply when aphid numbers reach spray threshold levels as determined by regular monitoring. Ensure thorough coverage of all leaves.
Pandanus Trees	Flatid ( <i>Jamella australiae</i> )	<b>Spot spray</b> 875 mL/ 100L of water  <b>Stem injection</b> 1.75L/1L of water		<b>Spot spray:</b> Spray 100 mL of mixture directly into the leafy throat of each head. <b>Stem injection:</b> Drill holes 0.5 to 1cm in diameter and 10cm deep at an angle of 30°, 1 to 1.5m above ground level. Drill one hole per limb (or trunk in single trunked trees). Apply 5 mL of mixture in each hole and seal the hole. <b>DO NOT</b> re-apply in the same holes. Uptake of Imtrade Annihilate 200 SC All Purpose Insecticide, and therefore control of the pest in heavily infested heads already showing severe damage, will be slow and may be incomplete.

## SOIL DRENCH APPLICATION

CROP	PEST	RATE	CRITICAL COMMENTS
Elm	Elm Leaf Beetle	7 mL/25mm of tree diameter at breast height	Mix the required dose in sufficient water to adequately treat each tree. Use at least 50L of mix per tree up to a tree diameter of 400-500mm and then 100L per tree for larger trees. Inject mix to a depth of 20-30cm in a minimum of 4 injection sites per tree, 0.75 to 1.5m apart, arranged in an evenly spaced grid to just beyond the drip line. Ensure root zone is adequately moist with active root growth. Keep treated area moist for 7 – 10 days after treatment. Treat at least 6-10 weeks prior to pest attack in late winter or early spring when roots are active. <b>DO NOT</b> treat if soil is waterlogged.
Azaleas in Pots	Azalea Lace Bug	3.5 mL/250 mL water/pot	Use as a soil drench for pots up to 20L capacity. Prior to application remove mulch and dead vegetation, and moisten the soil surface. Apply the Imtrade Annihilate 200 SC All Purpose Insecticide mixture, and then water it in well immediately after application.
Ornamentals in Pots	Scarab Beetle Larvae	3.5 mL/5L water	Use as a soil drench 5L of mixture will treat twenty 6L pots. Prior to application remove mulch and dead vegetation, and moisten the soil surface. Apply the Imtrade Annihilate 200 SC All Purpose Insecticide mixture, and then water it in well immediately after application.

## SOIL DRENCH APPLICATION (continued)

CROP	PEST	RATE	CRITICAL COMMENTS
Roses	Aphids	3.5 mL/2L water/plant	Use as a soil drench by pouring mixture evenly around drip zone. Use this rate for plants up to 1m high. For each additional metre of plant height, add 2 mL extra of Imtrade Annihilate 200 SC All Purpose Insecticide to the 2L of water. Prior to application remove mulch and dead vegetation, and moisten the soil surface. Apply the Imtrade Annihilate 200 SC All Purpose Insecticide mixture, and then water it in well immediately after application.
Seedling eucalypts (to 1 m high) in pots	Chrysomelid Beetle Larvae, Psyllids	2.5 mL/plant	Mix in water up to 0.5L per 3L pot and apply to soil. Use less water for smaller pots. <b>DO NOT</b> dilute to the point where mix runs out the bottom of pots.
Turfgrass	First instar larvae of: African Black Beetle, Argentinian Scarab, Pruinose Scarab	2.5L/ha or 25 mL/100 m <sup>2</sup> Spray with at least 400L water per hectare to ensure even coverage. Preferably spray onto wet or dewy grass. Irrigate with 12mm of water commencing within one hour of application.	Apply at peak egg hatch, that is mid-spring to mid-summer depending on species.
	Larvae of Billbug		Monitor adult activity through late spring and early summer. Spray when numbers peak, or when small larvae (4mm) are found in the thatch or soil surface. Early application is essential to minimize grass damage due to feeding.



**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL  
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

### **WITHHOLDING PERIODS**

**Apples:** NOT REQUIRED WHEN USED AS DIRECTED

**Cotton:** DO NOT HARVEST FOR 13 WEEKS AFTER APPLICATION  
DO NOT GRAZE OR CUT FOR STOCK FOOD

**Cucurbits:** DO NOT HARVEST FOR 1 DAY AFTER APPLICATION

**Tomatoes:** DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION

**Brassicas, Capsicum, Eggplant, Potatoes, Sweet Potatoes:**  
DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION

**Stone fruit:** DO NOT HARVEST FOR 21 DAYS AFTER APPLICATION

**DO NOT graze any treated area, or cut for stock food.**

**DO NOT feed produce harvested from treated area to animals, including poultry.**

### **GENERAL INSTRUCTIONS**

#### **INSECTICIDE RESISTANCE WARNING**

#### **GROUP 4A INSECTICIDE**

For insecticide resistance management, Imtrade Annihilate 200 SC All Purpose Insecticide is a Group **4A** Insecticide. Some naturally occurring insect biotypes resistant to Imtrade Annihilate 200 SC All Purpose Insecticide and other Group **4A** Insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Imtrade Annihilate 200 SC All Purpose Insecticide and other Group **4A** Insecticides are used repeatedly. The effectiveness of Imtrade Annihilate 200 SC All Purpose Insecticide on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Imtrade accepts no liability for any losses that may result from the failure of Imtrade Annihilate 200 SC All Purpose Insecticide to control resistant insects. Imtrade Annihilate 200 SC All Purpose Insecticide may be subject to specific resistance management strategies. For further information contact your local supplier, Imtrade representative or local agricultural department agronomist.

#### **RESISTANCE MANAGEMENT STRATEGY**

##### **Cotton Aphid in cotton**

Observe the cotton industry Insecticide Resistance Management Strategy (IRMS).

### **Aphids, Whitefly and Melon Thrips in various crops**

**DO NOT** apply Imtrade Annihilate 200 SC All Purpose Insecticide (or other Group **4A** Insecticide) in consecutive sprays within and between seasons. Rotate with registered insecticides from other mode of action groups.

#### Confined environments such as glasshouses:

**Annuals** – **DO NOT** apply more than one spray of Imtrade Annihilate 200 SC All Purpose Insecticide (or other Group **4A** Insecticide) to any one crop.

**Perennials** – Rotate with registered insecticides from other groups. Use a maximum of three Imtrade Annihilate 200 SC All Purpose Insecticide (or other Group **4A** Insecticide) sprays in any 12 months period.

### **MIXING/APPLICATION**

Prior to pouring, shake container vigorously, then add the required quantity of Imtrade Annihilate 200 SC All Purpose Insecticide to water in the spray vat while stirring or with agitators in motion.

#### **Special Instructions for Stone Fruit**

##### **Dilute Spraying**

- Use a sprayer designed to apply high volumes of water up to the point of runoff and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of run-off. Avoid excessive run-off.
- The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice.
- Add the amount of product specified in the Directions for Use table for each 100L of water. Spray to the point of run-off.
- The required dilute spray volume will change and the sprayer set up and operation may also need to be changed, as the crop grows.

##### **Concentrate Spraying**

- Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume.
- Determine an appropriate dilute spray volume (See Dilute Spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate.

- The mixing rate for concentrate spraying can then be calculated in the following way:  
EXAMPLE ONLY
  1. Dilute spray volumes as determined above: For example 1500L/ha
  2. Your chosen concentrate spray volume: For example 500L/ha
  3. The concentration factor in this example is:  $3 \times$  (ie 1500L, 500L = 3)
  4. As the dilute label rate is 25 mL/100L for stone fruit, then the concentrate rate becomes  $3 \times 25$ , that is 75 mL/100L of concentrate spray.
- The chosen spray volume, amount of product per 100L of water, and the sprayer set up and operation may need to be changed as the crop grows.
- **DO NOT** use a concentrate rate higher than that specified in the Critical Comments.
- For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

### Application (Cotton)

Thorough coverage of cotton plants is essential to achieve maximum performance from Imtrade Annihilate 200 SC All Purpose Insecticide plus Pulse™. Equipment should be calibrated to achieve a **minimum of 60 droplets/cm<sup>2</sup> on the target foliage**. A droplet **Volume Median Diameter (VMD)** for optimum performance from Imtrade Annihilate 200 SC All Purpose Insecticide plus Pulse™ is dependent on equipment and is defined below. Do not apply when unfavorable environmental conditions may reduce the quality of spray coverage.

### **Ground Application (Cotton)**

Application using ground equipment should be made using **hollow cone nozzles** with a **minimum spray volume of 100L/ha**. Hollow cone nozzles are recommended but if flat fan nozzles are used, higher water volumes will be required and nozzles should be configured to ensure thorough coverage. A droplet **VMD of 150–180 microns** must be used. Where multiple nozzles per row are used, they should be of the same specification to ensure that each nozzle contributes an equal proportion of the required dose. Where multiple nozzles per row are used (particularly for banded applications) ensure the correct nozzle overlap pattern is achieved on the target foliage. **Banded applications less than 100% are not recommended beyond the 15 node crop stage.**

### **Aerial Application (Cotton)**

Apply in a minimum **spray volume of 25L/ha**. A droplet **VMD of 120–150 microns** must be used. **DO NOT** exaggerate swath width or exceed a swath width of 20 to 22m. **DO NOT** apply Imtrade Annihilate 200 SC All Purpose Insecticide plus Pulse™ using Ultra Low Volume (ULV) methods. The use of large droplet placement equipment is not recommended.

### **COMPATIBILITY**

Imtrade Annihilate 200 SC All Purpose Insecticide is compatible with Propineb, Bitertanol, Methamidophos. **DO NOT** mix concentrates together but add each to the spray tank separately. As formulations of other manufacturers' products are beyond the control of Imtrade, all mixtures should be tested prior to mixing commercial quantities. As changes in climatic conditions can alter the sensitivity of plants to mixtures of sprays, Imtrade cannot be responsible for the behavior of such mixtures.

## **EXPORT OF TREATED PRODUCE**

Growers should note that MRLs or import tolerances do not exist in all markets for edible produce treated with Imtrade Annihilate 200 SC All Purpose Insecticide. If you are growing edible produce for export, please check with Imtrade for the latest information on MRLs and import tolerances before using Imtrade Annihilate 200 SC All Purpose Insecticide.

### **Note on Ornamentals**

Imtrade Annihilate 200 SC All Purpose Insecticide has been used on a wide range of ornamental plant species without damage. However, some species and varieties are particularly sensitive to chemical sprays and as this is often related to local conditions it is advisable to treat only a small number of plants first, in order to ascertain their reaction before treating the whole crop.

## **PROTECTION OF LIVESTOCK**

Dangerous to bees. **DO NOT** spray any plants in flower while bees are foraging.

**DO NOT** graze any treated area, or cut for stock food.

**DO NOT** feed produce harvested from treated area to animals, including poultry.

## **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

**DO NOT** contaminate streams, rivers or waterways with the chemical or used containers. A spraydrift minimization strategy should be employed at all times when aerially applying sprays. The strategy envisaged is exemplified by the cotton industry's Best Management Practices Manual.

### **STORAGE AND DISPOSAL (200mL, 1L, 2.5L only)**

Store product in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to the spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling break, crush, or puncture and bury containers at a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burnt.

### **STORAGE AND DISPOSAL (5L, 10L & 20L only)**

Store product in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling break, crush, or puncture and bury containers at a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burnt.

## **STORAGE AND DISPOSAL (60L only)**

Store product in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. If tamper evident seals are broken prior to initial use then the integrity of the contents cannot be assured. Empty container by pumping through dry-break connection system. **DO NOT** attempt to breach the valve system or the filling point, or contaminate the container with water or other products. Ensure that the coupler, pump, meter and hoses are disconnected, triple rinsed and drained after each use. When empty, or contents no longer required, return the 60L container only to the point of purchase. This container remains the property of Imtrade Australia Pty Ltd.

## **SAFETY DIRECTIONS**

Harmful if swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. Repeated exposure may cause allergic disorders. When preparing product for use, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

## **FIRST AID**

If poisoning occurs contact a doctor or Poisons Information Centre (Phone Australia 13 11 26).

### **Additional GHS Hazard & Precautionary Statements**

- Avoid breathing fumes, mists vapours or spray;
- **DO NOT** get on clothing;
- Contaminated work clothing should not be allowed out of the workplace;
- **IF SWALLOWED:** Rinse mouth. **DO NOT** induce vomiting;
- **IF ON SKIN:** Wash with plenty of soap and water;
- **IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing;
- If skin irritation occurs: Get medical advice;
- If eye irritation persists: Get medical advice;
- Not combustible. Use extinguishing media suited to burning materials;
- Store in a dry place.

## **SAFETY DATA SHEET**

Additional information is listed in the Safety Data Sheet which can be obtained from your supplier or Imtrade Australia Pty Ltd website at [www.imtrade.com.au](http://www.imtrade.com.au)

## **CONDITIONS OF SALE**

Imtrade Australia Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgement in purchasing or using same and every person dealing with this product does so at his/hers own risk. No representative of Imtrade has any authority to add to or alter these conditions.

™ Other Registered Trademarks

**APVMA Approval No. 60465/106758**