

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

STORAGE AND DISPOSAL

Store in the closed original container in a dry, cool, well-ventilated area away from children, animals, food, feedstuffs, herbicides, seed and fertilizers. **DO NOT** store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemical on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers and product should not be burnt. **DO NOT** re-use empty container for any other purpose.

For refillable containers: Empty contents fully into application equipment. Close all valves and return to the point of supply for refill or storage.

SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with eyes and skin. **DO NOT** inhale spray mist. When opening container and preparing spray wear elbow-length PVC gloves and face-shield or goggles. If product on skin, immediately wash area with soap and water after use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves and face shield or goggles and contaminated clothing.

IMTRADE

DICAMBA 500

SELECTIVE HERBICIDE

ACTIVE CONSTITUENTS: 500 g/L DICAMBA (present as the dimethylamine salt)

GROUP 4 HERBICIDE

For the control of certain broadleaf weeds in winter cereals, pastures, conservation tillage, sugar cane, turf, rice, grain sorghum, pine plantations and non-crop areas as specified in the Directions for Use Table.

IMPORTANT: Read the attached Leaflet before use.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre. (Phone Australia 13 11 26; New Zealand 0800 764 766).

Refer to attached Leaflet for additional GHS Hazard & Precautionary Statements.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from your supplier or from Imtrade Australia Pty Ltd website at www.imtrade.com.au

CONDITIONS OF SALE

Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Imtrade Australia Pty Ltd has any authority to add to or alter these conditions.

In a Transport Emergency Dial 000
Police or Fire Brigade

APVMA Approval No. 54658/122047

GROUP

4

FORM

SL

CONTENTS

5L-1000L



Batch No.
DOM:



17 Ocean St, Kwinana Beach WA 6167
08 9419 0333 | www.imtrade.com.au

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



IMTRADE DICAMBA 500 SELECTIVE HERBICIDE

ACTIVE CONSTITUENT: 500 g/L DICAMBA
(present as the dimethylamine salt)

GROUP	4	HERBICIDE
-------	----------	-----------

For the control of certain broadleaf weeds in winter cereals, pastures, conservation tillage, sugar cane, turf, rice, grain sorghum, pine plantations and non-crop areas as specified in the Directions for Use Table

**IMPORTANT: READ THIS LEAFLET THOROUGHLY BEFORE
OPENING OR USING THIS PRODUCT**

APVMA Approval No. 54658/122047

Imtrade Australia Pty Ltd
ABN 13 090 151 134
17 Ocean Street, Kwinana Western Australia 6167
Tel: 1800 171 799 Fax: 1800 17 1788
Web: www.imtrade.com.au

RESTRAINTS:**DO NOT** spray when rain seems likely to occur within 4 hours.**DO NOT** spray when weeds are wet with dew or rain or under stress from drought, low soil fertility, extreme cold or water logging.**DO NOT** spray outside recommended crop growth as crop damage may result.**DO NOT** apply to crops or pastures with clover, lucerne or medics.**DIRECTIONS FOR USE**

Crop	Weeds Controlled	State	Rates/ha	Critical Comments
Wheat, Oats, Barley, Triticale, Commercial Rye. From 5 leaf to early tillering (Zadoks 15 - 22)	Doublegee (Three-cornered Jack, Spiny Emex), Wireweed (Hogweed), Seedling Docks, Tares, Hexham Scent (<i>Melilotus</i>), Wild Turnip (<i>Brassica tournefortii</i>), Soldier Thistle, Mustards, Wild Radish, Capeweed, Common Ice-Plant, Charlock, Ward's Weed, Vetch, Turnip Weed (<i>Rapistrum rugosum</i>)	All States	160 mL plus 700 mL of MCPA Amine 500g/L	Spray when most weeds have germinated and are in the 2 to 4 leaf stage. In some instances suppression only of wireweed (Hogweed) and Capeweed will occur.
Wheat, Oats, Barley, Triticale, Commercial Rye. From 5 leaf early tillering stage until the fully tillered stage and before jointing occurs (Zadocks 15, 21-30)	Doublegee (Three-cornered Jack, Spiny Emex), Wireweed, Vetch, Variegated Thistle, Tares, Sunflower, Hexham Scent (<i>Melilotus</i>), Docks.	SA	280 mL	Spray when most weeds have germinated and are still in the young rosette stage.
	Doublegee (Three-cornered Jack, Spiny Emex), Climbing Buckwheat, (Black Bindweed), Clover, Hexham Scent (<i>Melilotus</i>), Docks	WA		When weeds do not form a rosette, spray when they have not more than 8 true leaves. In some instances suppression only of wireweed (Hogweed) and Climbing Buckwheat (Black Bindweed) will occur.
	New Zealand Spinach, Climbing Buckwheat (Black Bindweed), Doublegee Three-cornered Jack, Spiny Emex), Wireweed (Hogweed), Hexham Scent (<i>Melilotus</i>), Docks, Clover, Fat-Hen, Variegated Thistle, Scotch (Cotton or Woolly) Thistle, Sunflowers, Vetch, Tree Hogweed	Qld, NSW, Vic		

Crop	Weeds Controlled	State	Rates/ha	Critical Comments
Wheat, Oats, Barley, Triticale, Commercial Rye. From 5 leaf early tillering stage until the fully tillered stage and before jointing occurs (Zadocks 15, 21-30) (Cont'd)	Wild Turnip (<i>Brassica tournefortii</i>), Radish, Mustards, Tares, Charlock, Capeweed, Sorrel, Saffron Thistle, Chickweed, Clover, Common Ice-Plant, Hoary Cress, Ward's Weed, Turnip Weed (<i>Rapistrum rugosum</i>)	SA	280 mL plus 1L of MCPA Amine 500g/L or 700 mL of 2,4-D Amine 500g/L	DO NOT use Imtrade Dicamba 500 Selective Herbicide plus 2,4-D in Oats in SA. If any of these weeds are found in these cereal crops as well as the above weeds, the addition of MCPA or 2,4-D is essential. Spray when most weeds have germinated and are still in the young rosette stage.
	Common Ice Plant	WA	320 mL plus 800 mL of 2,4-D Amine 500g/L	
	Chickweed, Sorrel, Wireweed	WA	400mL or 280mL plus 800mL of 2,4D Amine 500g/L or 280mL plus 1.2L MCPA Amine 500g/L	In some instances, suppression only of Wireweed (Hogweed), Capeweed and Hoary Cress will occur.
	Mintweed	NSW, Qld	280 mL plus 1L of MCPA Amine 500g/L or 700 mL of 2,4-D Amine 500g/L	A mixture with 2,4-D Amine must not be used in Tasmania.
	Wild Turnip (<i>Brassica tournefortii</i>), Radish, Mustards, Tares, Charlock, Capeweed, Sorrel, Saffron Thistle, Chickweed, Turnip Weed (<i>Rapistrum rugosum</i>)	Qld, NSW, Vic, Tas		

Crop	Weeds Controlled	State	Rates/ha	Critical Comments
Wheat, Barley, Triticale, Commercial Rye From early tillering (4 to 5 fully emerged main stem leaves, plus one or more tillers) until fully tillered and before the start of jointing (Zadocks 21-30)	Climbing Buckwheat (Black Bindweed), Deadnettle, Turnip Weed (<i>Rapistrum rugosum</i>), Wireweed, Wild Turnip (<i>Brassica tournefortii</i>)	NSW, Qld	200 – 280 mL plus 5 - 7g Metsulfuron	Spray when weeds have no more than 8 true leaves. Use the higher rate on larger weeds. Tank mixes with Imtrade Dicamba 500 Selective Herbicide and Metsulfuron may result in reduced control of Wireweed in some situations.
Barley, Triticale, Commercial Rye. From 5 leaf until fully tillered and before jointing occurs (Zadocks 21-30)	Volunteer Peas	SA	60 mL plus 1L KAMBA M*	Apply when most peas are 5 - 7cm high.
Wheat, Oats From 5 leaf until fully tillered and before jointing occurs (Zadocks 15-30)			120 mL plus 1L KAMBA M*	
Grain Sorghum From 3 leaf stage until 6 leaf stage Maize Post emergence until maize is approx 90cm high or until 15 days prior to tassle emergence.	Caltrop (Yellow Vine), Bathurst & Noogoora Burrs, Partenium Weed, Mintweed, Annual Gooseberry, Blackberry Nightshade, Fat-Hen, Sunflower, Seedling Perennial Gooseberry, Seedling Khaki Weed, Redshank, Dwarf Amaranth, Redroot Amaranth, Green Amaranth, Cobbler's Pegs, Field Bindweed, Thornapples (<i>Datura</i>), Bellvine, Lucerne, Annual Ground Cherry, Wireweed, Climbing Buckwheat (Black Bindweed), Pigweed, Common Sowthistle, New Zealand Spinach, Doublegee, (Three-cornered Jack, Spiny Emex).	All States (NOT Central Qld)	320 – 560 mL	Apply when weeds have 3 to 5 true leaves. DO NOT treat weeds which are beyond the rosette stage. Method of Application: By boom spray as an overall directed spray, applying 100 to 250L of water per hectare or by aircraft applying 45L per hectare. Use the higher rate on larger weeds and where Amaranthus, Mintweed and Blackberry Nightshade are the major problems or where Annual and Perennial Gooseberry are present. Certain maize varieties may develop clubfoot of the prop roots.

Crop	Weeds Controlled	State	Rates/ha	Critical Comments
Grain Sorghum From 3 leaf stage until 6 leaf stage	Thornapple (Datura), Mintweed, Amaranthus, Black (Giant) Pigweed, Caltrop (Yellowvine), Bladder Ketmia, Noogoora Burr.	NSW, Qld, NT	280 mL plus 1.8L Atrazine 500g/L or 320 mL plus 1.5L Atrazine 500g/L	Apply when weeds have 3 to 5 true leaves. DO NOT apply in hot dry conditions. Apply by boom spray as an overall spray applying 50 to 250L of water per hectare. Use the higher rate of Imtrade Dicamba 500 Selective Herbicide where Caltrop or Bladder Ketmia is present.
Rice Pre-Sowing Post-Sowing either before permanent water or until rice is at the mid-tillering stage.	Dirty Dora, Docks, Sedges, Starfruit	All States	400 – 560 mL 400 mL (seedling Docks) 560 mL (mature Docks)	Pre-sowing: Apply before the end of August and do not cultivate or graze for 14 days after spraying. Post-sowing: Preferably apply before the application of permanent water. If permanent water has been applied Imtrade Dicamba 500 Selective can be applied up to the mid-tillering stage, provided the Docks are exposed. Use 100L/ha for boom application.
Pinus Radiata Plantations At least 10 months prior to planting pines	Sorrel (in pastures to be planted to pines)	All States	1.1L or 560 mL plus 1.4L of 2,4-D Amine 500g/L	Apply in September / October period before the sorrel flowers. Add Agral 60*, ChemAg Wetter 1000 or another non-ionic wetting agent to the mixture.
	Naturally regenerated pines	SA, Vic, NSW	2 – 8L	Use the high rate where total control is essential. Apply in Spring.

Crop	Weeds Controlled	State	Rates/ha	Critical Comments
Pine Plantations	Weaken trees sufficiently to encourage attack by Sirex Wood Wasp (<i>Sirex noctilio</i>) to enable detection, monitoring or control.	NSW, Qld, NT	Prepare diluent at the ratio of 1L of Imtrade Dicamba Selective Herbicide to 1.5L water. Apply 1 mL of prepared diluent per 10cm of tree circumference.	Apply by drill injection, basal spear incision, axe incision or chainsaw cuts. Drill injection is the preferred option. Inject diluent into sapwood, not bark. Apply Dicamba treatment to selected trees prior to the peak Sirex flight season.
Sugarcane Post-emergence	Bathurst Burr, Bellvine, Blackberry Nightshade, Calopo, Caltrop, Cleome, Cobbler's Pegs, Common Sow Thistle, Convolvulus, Fat-Hen, Green Amaranth, Khaki Weed, Noogoora Burr, Parthenium Weed, Perennial Gooseberry, Pigweed, Pink Burr, Redroot Amaranth, Sensitive Plant, Sida-retusa, Tar Vine, Thickhead, Thornapple.	NSW, Qld, WA, NT	560 mL	Spray when most weeds have germinated and are in the young rosette stage or when they have not more than 8 true leaves. Method of Application: By boom spray as an overall directed spray, applying 100 to 250L of water per hectare or by aircraft applying 45L per hectare.
	Amaranthus, Black (Giant) Pigweed, Caltrop, Mintweed, Noogoora Burr, Pigweed, Thornapple (<i>Datura</i>).	NSW, Qld, NT	280 mL plus 1.8L Atrazine 500g/L or 320 mL plus 1.5L Atrazine 500g/L	Apply when weeds have 3 to 5 true leaves. DO NOT apply in hot dry conditions. Apply by boom spray as an overall spray applying 50 to 250L of water per hectare. Use the higher rate of Imtrade Dicamba 500 Selective Herbicide where Caltrop is present.

Crop	Weeds Controlled	State	Rates/ha	Critical Comments
Potatoes	Clover, Docks, Wireweed (Hogweed), Thistles, Chickweed, Climbing Buckwheat (Black Bindweed), Black Nightshade, Fat-Hen.	Tas	500 mL – 1.2L	Apply after haulm senescence in sufficient water to give even coverage. May be mixed with Amitrole or 2,4-D to improve spectrum of weeds controlled. Refer to respective labels for application rates. Add suitable surfactant for improved coverage. NOTE: DO NOT apply when potato haulms are still green.

Crop	Weeds Controlled	State	Rates/ha			Critical Comments
			/ha	High Vol /100L	Per 15L Knapsack	
Grass Pastures and Perennial Grass Seed Crops	Cotton (Woolly or True Scotch) Thistle, Variegated Thistle, Star Thistle, Black or Spear Thistle, Illryian Thistle, Doublegee (Three-cornered Jack, Spiny Emex), Parthenium Weed, Smart weed (Water Pepper), Hexham Scent (Melilotus), Stemless Thistle, Caltrop (Yellow Vine), Horehound and Rumex spp., including Curled and Broadleaf Docks	All States	280 - 560 mL for seedlings	40 mL for seedlings	10 mL for seedlings	When applying by boom spray use 110 to 280L of mixture per hectare.
	560 mL - 1L for young mature plants	76 mL for young mature plants	18 mL for young mature plants			
	Common Ice-Plant, Hoary Cress, Sorrel and Saffron Thistle.	All States	280 mL plus 800 mL of 2,4-D Amine 500g/L	20 mL plus 60 mL of 2,4-D Amine 500g/L	6 mL plus 15 mL of 2,4-D Amine 500g/L	Add Agral 60* or Nufarm Surfactant 600* wetting agents to spray mix.

Crop	Weeds Controlled	State	Rates/ha			Critical Comments
			/ha	High Vol /100L	Per 15L Knapsack	
Non-Crop Situation	Annual Ragweed, Blue Heliotrope, Burr Ragweed, Galvanised Burr, Ground Cherry, Ox-eye Daisy, Perennial Ragweed, Russian Knapweed, Skeleton Weed, Wild (Field or Crow) Garlic	All States	8.8L	600 mL	130 mL	Use a minimum of 1500L/ha water carrier. Add Agral 60* or Nufarm Surfactant 600* wetting agents. Spray Russian Knapweed at flowering. Spray Angled Onion Weed, Bladder Champion, Ox-Eye Daisy and Paterson's Curse before flowering. Spray Wild Garlic during Spring before bulbs start to form and re-spray later if dormant bulbs shoot. Spray prior to flowering.
	California (Perennial or Canada) Thistle, Field Bindweed, Paterson's Curse, Prairie Ground Cherry, Ragwort, Stinking Mayweed (Fetid Chamomile)		4L	280 mL	60 mL	
	Afghan Thistle, Artichoke Thistle, Star Thistle		1.6L	100 mL	24 mL	
	Horehound, Angled Onion Weed (Three-cornered Garlic), Bladder Champion, Cotton (Woolly or True Scotch) Thistle, Docks, Sorrel, Stemless Thistle, Variegated (or Spotted) Thistle		1.2L	80 mL	18 mL	
	Illryian Thistle, Parthenium Weed, Smart-Weed		600 mL	40 mL	10 mL	
	Poverty Weed	SA		2L	440 mL	
	Mist Flower	Qld, NT	12L	800 mL	180 mL	

Crop	Weeds Controlled	State	Rates/ha			Critical Comments
			/ha	High Vol /100L	Per 15L Knapsack	
Non-Crop Situation (Cont'd)	Blackberry Knightshade	WA	320 - 560 mL		5-9 mL	Blackberry Knightshade at seeding stage. For Creeping Knotweed apply in early Summer to actively growing plants. For Slender Ice-Plant and Khaki Weed spray when actively growing at young rosette stage. For Rubber Vine apply in Autumn during active growth.
	Creeping Knot Weed	WA	1.2 – 2.4L		18 – 36 mL	
	Slender Ice-Plant	WA	320 mL + 800 mL 2,4-D 500g/L		6 mL	
	Khaki Weed	WA	600 mL		9 mL	
	Rubber Vine	WA	1.4L		22 mL	
Grass Pastures and Non-Crop	Rubber Vine	Qld, NT	2-4L plus 2.5L of 2,4-D Ester 800g/L	140 - 280 mL plus 175 mL of 2,4-D Ester 800g/L	20 - 40 mL plus 25 mL of 2,4-D Ester 800g/L	Apply during April or May. For application by misting machines, add 600 mL Imtrade Dicamba 500 Selective Herbicide to 10L of water.
Turf	Catsear, Cudweeds, Pennyworth, Lambs Tongue, Fleabanes, Fumitory, Wireweed, Chickweed, Cotula, Sorrel, Dock, Clovers, Capeweed, Toad Rush, Peppergrass, Dandelion, Scarlet Pimpernel, Creeping Oxalis, Pearlwort, Trefoil, Yarrow, Bindy-Eye, Jo-Jo (Onehunga)	All States	1.2L plus 4L of 2,4-D Amine 500g/L	80 mL plus 250 mL of 2,4-D Amine 500g/L	18 mL plus 60 mL of 2,4-D Amine 500g/L	Use a minimum of 1000L/ha water carrier. DO NOT spray on Buffalo Grass, Bent Grass, Lippia or Strawberry Clover. Spray when weeds are young and actively growing.
	Fat-Hen, Red lowered Mallow, Purple-Top, Stagger Weed, Swine Cress	NSW, NT, SA, Qld, WA, Tas				

Crop	Weeds Controlled	State	Rates/ha			Critical Comments
			/ha	High Vol /100L	Per 15L Knapsack	
Woody Species	Moreton Bay Ash (Carbeen), Blue Gum, Bitter Bark, Brisbane Black Wattle	All States	By partial drill or tree injection technique mix 1 part Imtrade Dicamba 500 Selective Herbicide 500 with 12.5 parts of water and apply 2mL of the mixture to cuts in the softwood at not more than 6cm spacings around the circumference of each tree.			
	Mimosa (<i>Minosa pigra</i>)	NT	2 – 2.4L	400mL	60mL	Apply to actively growing plants. Preferably apply during wet season. Retreatment may be necessary to control seeding regrowth (for application see General Instructions)

ADDITIONAL RESTRAINTS FOR CONSERVATION TILLAGE

- When grass and broadleaf weeds are present, use a mixture of Imtrade Dicamba 500 Selective Herbicide and Glyphosate 450.
- **DO NOT** disturb treated weeds by cultivation, sowing or grazing for 1 day after treatment of annual weeds and 7 days for perennial weeds.
- **DO NOT** treat weeds heavily covered with dust or silt.
- **DO NOT** apply if rainfall is likely within 6 hours of application.
- Observe plant back periods listed under General Instructions.
- When applying to dry soil surfaces, at least 15mm of rainfall is required before the plant back period begins.

Crop	Weeds Controlled	State	Rates/ha	Critical Comments
Seed Bed Salvage, Conservation Tillage, Direct Drilling For weed control in fallow, stubble or pasture prior to sowing with a full disturbance implement.	Amaranthus, (Red Shank), Native Amaranthus, Sow Thistle, Volunteer Cereals, Turnip Weed, Wireweed (Hogweed), Medics, Variegated Thistle, Wild Oats, Wild Mustard, Docks, Paterson's Curse	Qld, NSW, Vic, SA, WA & NT	160 - 240 mL plus Glyphosate 450g/L at 400 - 600 mL or Glyphosate 360g/L at 500 - 700 mL	If excessively wet conditions occur shortly after application or sowing, some retardation of crop establishment may occur in certain broadleaf crops. Spray when most weeds have germinated and are still in the young rosette or pre-tillering stage. Use the lower rate when weeds are prior to tillering or seedlings. Increase to the higher rate as weeds reach tillering or are 10-15cm diameter/high. In grazed situations if heavy grazing has occurred allow re-growth to 6-8cm high before spraying. Visible symptoms of weed control may not be apparent for 10-21 days.
	Barnyard Grass, Liverseed Grass	NSW		
	Apple of Peru, Mintweed, Noogoora Burr, Yellow Vine (Caltrop), Climbing Buckwheat (Black Bindweed), Barnyard Grass, Canary Grass (Annual Phalaris), Thorn Apple, Wild or Prickly Lettuce	Qld, NSW, NT		
	Annual Ryegrass, Volunteer Field Peas, Tree Hogweed, Capeweed	NSW, Vic, SA, WA		
	Barnyard Grass, Liverseed Grass	Qld, NT	160 - 240 mL plus Glyphosate 450g/L at 800 mL - 1.2L	
	Sorrel	WA, SA, Vic, NSW	160 - 240 mL plus Glyphosate 450g/L at 650 - 800 mL or Glyphosate 360g/L at 800 mL - 1L	

Crop	Weeds Controlled	State	Rates/ha	Critical Comments
Direct Drilling, Conservation Tillage, No-Till Fallow, Minimum Tillage and all between cropping applications	Clover	All States	200 mL	Add to Glyphosate or Sprayseed as required (mixtures of Sprayseed plus Imtrade Dicamba 500 Selective Herbicide must not be used in Tas). Follow label recommended rates.
	Doublegee (Three-cornered Jack, Spiny Emex), Wireweed, Vetch, Docks, Tares, Sunflower, Hexham Scent (Melilotus), Variegated Thistle	SA	280 mL	Spray when most weeds have germinated and are still in the young rosette stage, when they have not more than 8 true leaves.
	Doublegee (Three-cornered Jack, Spiny Emex), Climbing Buckwheat (Black Bindweed), Hexham Scent (Melilotus), Docks	WA		
	New Zealand Spinach	Qld, NSW, NT		
	Climbing Buckwheat (Black Bindweed), Doublegee (Three-cornered Jack, Spiny Emex), Wireweed (Hogweed), Hexham Scent (Melilotus), Docks, Fat-Hen, Variegated Thistle, Scotch (Cotton or Woolly) Thistle, Sunflowers, Vetch, Tree Hogweed	Qld, NSW, Tas, Vic, NT		
	Chickweed, Sorrel, Wireweed	WA	400 mL	Spray when most weeds have germinated and are still in the young rosette stage, when they have not more than 8 true leaves.
	Common Ice-Plant, Hoary Cress, Sorrel, Saffron Thistle	All States	280 mL plus 800 mL of 2,4-D Amine 500g/L	

Crop	Weeds Controlled	State	Rates/ha	Critical Comments
Direct Drilling, Conservation Tillage, No-Till Fallow, Minimum Tillage and all between cropping applications (Cont'd)	Caltrop (Yellow Vine), Datura, Bathurst and Noogoora Burrs, Parthenium Weed, Amaranthus, Mintweed, Annual Gooseberry, Blackberry Nightshade, Fat-Hen, Sunflower, Seedling Perennial Gooseberry, Seedling Khaki Weed, Bellvine, Cobbler's Pegs, Star Thistle, Black or Spear Thistle, Illyrian Thistle, Doublegee (Three-cornered Jack, Spiny Emex), Smart-Weed (Water Pepper), Stemless Thistle, Horehound and Rumex spp. including Curled and Broadleaf Docks	All States	320 – 560 mL for seedlings	Apply when weeds have at least 3 to 5 true leaves and are actively growing. DO NOT treat weeds which are beyond the rosette stage. Use the higher rate on large weeds.
	Convolvulus, Sida-retusa, Sensitive Plant, Pink Burr, Cleome, Centro, Thickhead, Tar-Vine.	Qld, NSW, NT, Vic, Tas, WA		

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHOLDING PERIOD:

DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.

***Rice:* DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.**

***Pine Plantations:* NOT REQUIRED WHEN USED AS DIRECTED.**

GENERAL INSTRUCTIONS

Before opening, carefully read Directions for Use, Precautionary Statements, Safety Directions and First Aid Instructions.

MIXING

Add the required amount of the product directly to the tank and mix well.

APPLICATION

For Boom Application: Apply by a properly calibrated boom spray using not less than 50L of water per hectare unless indicated otherwise in Critical Comments column.

For Aerial application: Apply at least 25L total spray volume per hectare. The aircraft should fly as low as practical under the prevailing conditions to minimise drift.

For High Volume Application: Apply minimum 1500L per hectare.

For Knapsack Application: A 15L knapsack treats 150 square metres.

MIMOSA CONTROL

Broadacre application – Apply 2-2.4L/ha in 60L water per hectare. Helicopter application preferred.

High Volume application – Apply 400 mL in 110L water using ground equipment. Thoroughly wet all leaves and stems of the plant.

Basal Bark application – Apply 60 mL in 15L water to plants less than 5cm basal diameter. Spray liberally all around the base and 25-40cm up the stem.

Cut Stump application – Apply 60 mL in 15L water on plants up to 10cm basal diameter. Cut trees as close to ground level as is possible and spray the freshly cut surface before the sap dries.

Note: A temporary wilting may be evident in crops after application. If seasonal conditions are late or dry seek further advice before spraying as damage may result from spraying crops under these conditions. Growers should seek advice before spraying recently released cereal varieties. Certain Maize varieties may develop clubfoot of the prop roots.

PLANT BACK PERIODS for Conservation Tillage

Wait for the following periods before planting the following crops into soils sprayed with the product. If applied to dry soil, at least 15mm of rainfall is required before the Plant Back Period begins.

Note: if water-logging occurs shortly after sowing, the product may cause some reduction in crop vigour. Use as directed only in the States indicated.

NSW, VIC, SA, WA, TAS

Rate per Hectare	PLANT BACK PERIOD (DAYS)				
	Wheat	Barley	Oats	Triticale	Rye
200 mL	1	1	1	1	1
280 mL	7	7	7	7	7
560 mL	14	14	14	14	14

NSW, VIC, SA, WA, TAS

Rate per Hectare	PLANT BACK PERIOD (DAYS)				
	Lupins	Clover/Medics	Mung Beans	Sunflower	Safflower
200 mL	7	7	5	1	14
280 mL	14	14	5	7	21
560 mL	21	21	10	14	28

NSW, VIC, SA, WA, TAS

Rate per Hectare	PLANT BACK PERIOD (DAYS)				
	Sorghum	Maize	Rapeseed	Field Peas	Chick Peas
200 mL	1	1	7	N.D.*	N.D.*
280 mL	3	3	10	14	21
560 mL	7	7	14	21	28

* Not Determined

NSW, VIC, SA, WA, TAS

Rate per Hectare	PLANT BACK PERIOD (DAYS)			
	Pigeon Peas	Soybeans	Millet	Cotton
200 mL	5	5	1	7
280 mL	5	5	3	7
560 mL	10	10	7	14

QLD, NT only

RATE	CROP	PLANT BACK PERIOD
160 – 240 mL / Ha	Wheat, Barley, Oats, Triticale, Commercial Rye, Sorghum, Maize, Millet, Sunflowers	1 day
	Soybeans, Mung Beans, Pigeon Peas	5 days
	Cotton, Rapeseed	7 days

NOTE: FOR QLD, NT ONLY: FOR ALL CROPS AND RATES ABOVE 600ML/HA – PLANT BACK PERIOD 21 DAYS. FOLLOWING POTATO HAULM DESTRUCTION IN TAS: FOR RATES AND CROPS NOT INCLUDED ABOVE, A PLANT BACK PERIOD OF 6 WEEKS APPLIES GENERALLY AND 8 WEEKS FOR SEED CROPS

COMPATIBILITY

Imtrade Dicamba 500 Selective Herbicide is compatible with Flowable Igran, Glyphosate 360/450, Chlorsulfuron/Glean*, Metsulfuron/Ally*, Smash 500/2-4D Amine, 2-4D Ester 800, Diuron, Ordram (Molinate). Also compatible with Atrazines, but when used in combination with these products some temporary visible effects in the crop growth may be observed.

RESISTANT WEEDS WARNING

GROUP 4 INSECTICIDE

For herbicide resistance management Imtrade Dicamba 500 Selective Herbicide is a Group 4 Herbicide. Some naturally occurring weed biotypes resistant to Imtrade Dicamba 500 Selective Herbicide and other Group 1 Herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if Imtrade Dicamba 500 Selective Herbicide or other Group 4 Herbicides are used repeatedly. These resistant weeds will not be controlled by Imtrade Dicamba 500 Selective Herbicide or other Group 4 Herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Imtrade Australia Pty Ltd accepts no liability for any losses that may result from the failure of Imtrade Dicamba 500 Selective Herbicide to control resistant weeds.

Spray when weeds are young and actively growing, and unless stated otherwise, before the 3-5 true leaf stage.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. Spray equipment must be calibrated accurately prior to working on crops. Keep container sealed (airtight) when not in use.

To prevent damage, **DO NOT** permit drift or vapour or spray mist to come into contact with fruit and ornamental trees, vines, vegetables, flowers, tobacco or any other field crop. Minimise spray drift by using low pressure and nozzles, which **DO NOT** have a fine droplet size.

DO NOT apply this product on or near desirable trees or plants or in locations where the chemical may be washed or moved into contact with their roots.

DO NOT use in high wind. All spray equipment must be thoroughly washed out after use and must not be used for spraying horticultural crops.

DO NOT use at higher rate than recommended.

PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT use container for any other purpose.

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed original container in a dry, cool, well-ventilated area away from children, animals, food, feedstuffs, herbicides, seed and fertilizers. **DO NOT** store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemical on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers and product. **DO NOT** re-use empty container for any other purpose.

For refillable containers: Empty contents fully into application equipment. Close all valves and return to the point of supply for refill or storage.

SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with eyes and skin. **DO NOT** inhale spray mist. When opening container and preparing spray wear elbow-length PVC gloves and face-shield or goggles. If product on skin, immediately wash area with soap and water after use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves and face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre. (*Phone Australia 13 11 26; New Zealand 0800 764 766*).

Additional GHS Hazard & Precautionary Statements

•Harmful if swallowed; •Harmful to aquatic life with long lasting effects; •**DO NOT** get on clothing; •**DO NOT** eat, drink or smoke when using this product; •**IF SWALLOWED:** Call a POISON CENTRE or doctor if you feel unwell; •**IF SWALLOWED:** Rinse mouth. Do **NOT** induce vomiting; •**IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing; •If eye irritation persists: Get medical advice; •Not combustible. Use extinguishing media suited to burning materials.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from your supplier or from Imtrade Australia Pty Ltd website at www.imtrade.com.au

CONDITIONS OF SALE

Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Imtrade Australia Pty Ltd has any authority to add to or alter these conditions.

*Other Registered Trademarks

**In a Transport Emergency
Dial 000
Police or Fire Brigade**

APVMA Approval No. 54658/122047