

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. Store in a locked room or a place away from children, animals, food, feedstuffs, seed and fertilizers.

All other non-refillable containers

Triple rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers and product.

Envirodrum - Micro Matic Valve

Store the original sealed Envirodrum in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. **DO NOT** tamper with the Micro Matic valve or the security seal. **DO NOT** contaminate the Envirodrum with water or any other foreign matter. After each use of the product please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase.

Refillable Containers

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Very dangerous. Poisonous if absorbed by skin contact, inhaled or swallowed. Will irritate the eyes, nose, throat and skin. Avoid contact with eyes and skin. **DO NOT** inhale spray mist. When preparing spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves, face shield or goggles, and half-face respirator or disposable respirator. If clothing becomes contaminated with product or wet with spray remove clothing immediately. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, respirator and if rubber wash with detergent and warm water, face shield and goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (*Phone Australia 13 11 26; New Zealand 0800 764 766*). If in eyes, hold open, flood with water for at least 15 minutes and see a doctor.

Refer to attached Leaflet for additional GHS Hazard & Precautionary Statements.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from your supplier or Imtrade Australia Pty Ltd website at www.imtrade.com.au

CONDITIONS OF SALE

Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Imtrade Australia Pty Ltd has any authority to add to or alter these conditions.

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

IMTRADE

DIQUAT 200

NON-RESIDUAL HERBICIDE

ACTIVE CONSTITUENTS: 200 g/L DIQUAT present as
DIQUAT DIBROMIDE MONOHYDRATE

GROUP 22 HERBICIDE

For pre-harvest crop desiccation and the control of a wide range of broadleaf weeds in certain crops as per Directions for Use.

For application through aircraft and ground equipment.

IMPORTANT: Read the attached Leaflet before use.

UN 1760	CORROSIVE LIQUID, N.O.S. (contains DIQUAT)
In a Transport Emergency Dial 000 Police or Fire Brigade	
PG III	HAZCHEM 2X

APVMA Approval No: 58411/141286



GROUP
22

FORM
AC

CONTENTS
5L-1000L



Batch No.
DOM:



17 Ocean St, Kwinana Beach WA 6167
08 9419 0333 | www.imtrade.com.au

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



IMTRADE DIQUAT 200 NON-RESIDUAL HERBICIDE

**ACTIVE CONSTITUENTS: 200 g/L DIQUAT present as
DIQUAT DIBROMIDE MONOHYDRATE**

GROUP 22 HERBICIDE

For pre-harvest crop desiccation and the control of a wide range of broadleaf weeds in certain crops as per Directions for Use.
For application through aircraft and ground equipment.

**IMPORTANT: READ THIS LEAFLET THOROUGHLY
BEFORE OPENING OR USING THIS PRODUCT**

APVMA APPROVAL No: 58411/141286

Imtrade Australia Pty Ltd
ABN 13 090 151 134
17 Ocean Street, Kwinana, Western Australia 6167
Tel: (08) 9419 0333 Fax: (08) 9419 7516
Web: www.imtrade.com.au

RESTRAINTS:**DO NOT spray when weeds are under drought stress or when covered with dust or soil.****DO NOT apply with misting machines or CDA applications****DIRECTIONS FOR USE****PRE-HARVEST CROP DESICCATION**

Crop	State	Rate*	Critical Comments
Cotton (short stapled varieties only)	Qld, NSW, and WA only	2 to 3L/ha^	Apply when 85% of the bolls are open and remaining bolls are mature. Imtrade Diquat 200 Non-Residual Herbicide can damage green bolls.
Dry Beans Dry Peas Lentils Chickpeas Faba beans	All States	2 to 3L/ha^	Spray as soon as the crop has reached full maturity. Helps overcome slow and uneven ripening and weed problems at harvest.
Linseed	All States	2 to 3L/ha^	Spray when the majority of seed heads are mature – 90 to 95% of seed heads have changed from yellow to brown and the seeds rattle inside the bolls. Desiccation reduces the period from maturity to harvest, particularly under wet or humid conditions.
Lupins	All States	2 to 3L/ha^	Spray as soon as the crop has reached full maturity. Helps overcome slow and uneven ripening and weed problems at harvest.
Mung Beans	All States	2 to 3L/ha^	Apply when 80 to 90% of pods are black or brown. Desiccation of weeds and foliage aids timely and efficient harvesting, reduces harvester wear and tear but can increase harvest losses. Harvest 2 to 5 days after spraying.
Perennial Legume Seed Crops	All States	1.5 to 3L/ha^	<p>Lucerne- Spray when 60 to 70% of the pods are Brown/bluish and the seeds are Yellow/brown and easily released from the pods.</p> <p>Red Clover- Spray when majority of seed heads are brown and the seed is purple.</p> <p>White Clover- Spray when majority of seeds are hard and yellow.</p> <p>The use of Imtrade Diquat 200 Non-Residual Herbicide enables direct harvesting instead of cutting and windrowing and may result in higher seed quality. Harvest 3 to 4 days after spraying.</p>

PRE-HARVEST CROP DESICCATION (Continued)

Crop	State	Rate*	Critical Comments
Pigeon Peas	All States	2 to 3L/ha^	Spray as soon as the crop has reached full maturity.
Poppies	Tas only	3 to 4L/ha	Spray after the poppies have reached the stripy capsule stage. Helps overcome slow and uneven ripening and weed problems at harvest.
Potato (Haulm desiccation) Ground stored – pre-harvest weed control	All States	3 to 4L/ha^ 1.5L/ha plus 1.2L Paraquat 250 g/L	Apply as soon as crop is ready to harvest. DO NOT apply during drought periods, particularly when the tops will wilt during the day. In such conditions wait at least 3 days after the soil has been well moistened by rain or irrigation. Leaf kill is rapid following spraying and usually complete within 4 days. Stem kill may take 10 to 14 days. Lift when desiccation is complete but where possible wait for 14 days after spraying to allow skin to harden off. Use high water volumes to obtain coverage of dense haulm. Regrowth may occur if seed crops are desiccated early. To remove weed growth and facilitate digging, spray about 7 days prior to harvest. Where digging has been postponed and tubers stored in the ground often for a lengthy period, weed growth can be heavy and impede mechanical diggers unless removed.
Canola	All States	1.5 to 3L/ha^	Spray when 70% of the pods are yellow and the seeds are brown/bluish and pliable. Canola ripens unevenly and is prone to pod shatter and seed loss. Direct harvest 4 to 7 days after spraying.
Rice	All States	2 to 3L/ha^	Spray when the grain is mature – not more than 2 to 3% of the grain is still at the milky stage and the grain moisture content must be less than 25%.
Sorghum	All States	2 to 3L/ha^	Spray as soon as the seed is mature and the moisture content about 25%. Imtrade Diquat 200 Non-Residual Herbicide will advance harvest and reduce seed losses due to differential ripening, seed shedding and birds.
Soya Beans	All States	2 to 3L/ha^	Spray when 80% of the pods are yellow/brown and the seeds are ripe – yellow and pliable. Desiccation of weeds and foliage aids timely and efficient harvesting, minimizes cost and increases yields. Harvest 4 to 7 days after spraying.

PRE-HARVEST CROP DESICCATION (Continued)

Crop	State	Rate*	Critical Comments
Sugar Cane	Qld and NSW only	2 to 3L/ha [^]	Spray all accessible faces a few days prior to burning to a depth of about 30 metres. The sprayed cane and weed growth quickly dries out and ensures a good burn and removal or trash prior to harvest.
		High volume hand spraying 200 mL [^] / 200L water	Spray to visible wetness.
Sunflower	All States	2 to 3L/ha [^]	Spray when the seed is mature, seed moisture 35% and below, kernel full and firm, the disc spongy when broken, florets loose and bracts browning off. Harvesting can commence as soon as vegetative parts of crop are desiccated, usually 7 to 14 days after spraying.
Sweet Potatoes	All States		Apply 2 weeks prior to harvest.

GENERAL WEED CONTROL

Crop	Weeds Controlled	State	Rate*	Critical Comments
Aquatic areas	Duck Weeds, Red Azolla, Water Hyacinth, Salvinia	All States	5 or 10L/ha	Apply as an overall spray wetting foliage thoroughly. Clear water is necessary for best results as suspended soil particles interfere with herbicidal action. Use the higher rate for heavy infestations or for deep or dirty water. A repeat application 7 to 14 days later may be necessary for control of dense infestations. Oxygen depletion of decaying weeds may occur, therefore not more than ¼ of the area should be treated at once to ensure adequate oxygen supply for fish.
	Marsilea, Water Lilies and Water Lettuce		400 mL plus 150 mL Agral [®] per 100L water	Small areas – spray to wet weeds thoroughly. About 1 mL of product should be sufficient to treat about 1m ² .
	Cattail and Pond Weeds		5L/megalitre water	Apply by injection below the surface or as a surface spray.
Asparagus	Broadleaf weeds	All States	1.4L/ha plus 800 mL Agral [®] in 400L water	Apply to control seedling weeds before spears have emerged.

GENERAL WEED CONTROL (Continued)

Crop	Weeds Controlled	State	Rate*	Critical Comments	
Hops	Annual broadleaf and grass weeds	Vic and Tas only	700 mL to 1.4L/ha^ may be mixed with 1.2 to 1.6L/ha Imtrade Paraquat 250 Herbicide and/or 1.1kg Simazine granules	Apply as a directed inter-row spray prior to crop emerging from winter dormancy, using a minimum of 250L/ha spray volume to ensure good and even coverage weeds.	
Infested areas	Cotton Thistle (<i>Onopordum acanthium</i>)	Tas only	300 mL/ha plus 150 mL Agral® in 100L water	Spot spray at the rosette stage before the centre shoot is 15cm tall. The spray should be applied to give complete wetting of the leaf surface. DO NOT use a lower rate or treat at a later growth stage.	
	Saffron Thistle	All States	2.8L/ha plus 1L Agral® in 200L water/ha	Apply as an overall treatment to prevent seeding.	
			100 mL plus 70 mL Agral® per 15L knapsack	Alternatively spot spray on the same basis.	
Lucerne	Capeweed and <i>Erodium</i> spp.	All States	350 mL/ha^ in 200L water	Early Autumn application	Heavy grazing is necessary to reduce Lucerne to 2cm in height before spraying.
			700 mL/ha^ in 200L water	Late winter application	
Oil seed poppies	Weed control	Tas only	300 mL to 1.5L/ha	Use in accordance with recommendations made by Department of Primary Industries of the poppy contracting company. DO NOT add Agral® or any other wetting agent to the spray solution.	

GENERAL WEED CONTROL (Continued)

Crop	Weeds Controlled	State	Rate*	Critical Comments
Orchard and Vineyards	Capeweed	All States	1.5L plus 1.4L Agral® in 700L water per ha plus 1.6L/ha Imtrade Paraquat 250 Herbicide	Apply as a directed spray under trees or vines. Under most conditions Imtrade Paraquat 250 Herbicide at 1.6 to 3.2L/ha or Imtrade Spraykill 250 Herbicide at 2.4 to 3.2L/ha will give effective control of grasses and broadleaved weeds in orchards, but where heavy infestations of Capeweed occur Imtrade Diquat 200 Non-Residual Herbicide should be added to Imtrade Paraquat 250 Herbicide at the rate of 1.5L/ha. For inter-row or around butts use high volume applications. Paraquat 250 g/L, Imtrade Spraykill 250 Herbicide and Imtrade Diquat 200 Non-Residual Herbicide have no effect on brown bark but care should be taken when spraying around trees to avoid spray contacting green bark or plant material.
Pasture Renovation and establishment	Capeweed and <i>Erodium</i> spp. (Storksbill)	All States	1.5L plus 1.4L Agral® in 700L water per ha plus 1.6L/ha Imtrade Paraquat 250 Herbicide	Apply by boom spray as an overall spray on 'run-down' pasture after heavy grazing. Pasture should not be greater than 4cm long when sprayed. Grazing should be carried out during previous spring, summer and early autumn. Where Capeweed is in the very young seedling stage (2 or 3 true leaves only) rates may be reduced to 350 mL/ha. Where Capeweed infestation is high, oversowing with new pasture seed by direct drilling is advisable. Direct drill 3 to 7 days after spraying using a pasture mixture suitable to the district.
	Barley Grass, Brome Grass, Silver Grass and Sweet Vernal Grass		750 mL to 1.5L/ha [^] plus 1 to 2L/ha Imtrade Paraquat 250 Herbicide in a minimum of 100L water/ha	

GENERAL WEED CONTROL (Continued)

Crop	Weeds Controlled	State	Rate*	Critical Comments	
Row crops, vegetables and market gardens	Broadleaf weeds	All States	1.4L/ha [^]	(Seedling weeds)	Imtrade Spraykill 250 Herbicide and Imtrade Paraquat 250 Herbicide are more generally used for grass and broadleaf weed control in these situations. However where broadleaf weeds dominate, particularly Capeweed, Imtrade Diquat 200 Non-Residual Herbicide should be to tank mixed with Imtrade Paraquat 250 Herbicide or instead of Imtrade Paraquat 250 Herbicide where grass weeds are absent. Apply as a blanket spray prior to crop emergence. Once crops have emerged, or seedling has been transplanted, apply as a shielded spray between crop rows. DO NOT allow spray to contact any part of the crop.
			2.8 [^] to 4L/ha [^] per 200 to 300L water	(Mature weeds)	
Wheat and Oats	Capeweed	Qld, NSW, Vic, Tas, SA only	550 mL/ha in 200L of water	Small seedlings. DO NOT add wetting agent. Spray when the crop is between the 4 (Wheat) or 3 (Oats) leaf and early tillering stage.	
			700 mL/ha in 200L of water	Older seedlings. DO NOT add wetting agent. Spray when the crop is between the 4 (Wheat) or 3 (Oats) leaf and early tillering stage.	
Winter cereals	Pre-harvest weed control	All States	1 to 3L/ha [^]	Spray as soon as the crop is fully mature and ready for harvesting. Under wet spring conditions crops can periodically become infested with weeds, which seriously interfere with harvest operations. Imtrade Diquat 200 Non-Residual Herbicide will control these weeds allowing more efficient harvest.	
Wheat		NSW only	2L/ha [^]	Light to moderate stands	Ensure that spray penetrates deep down into the crop canopy.
			3L/ha [^]	Moderate to heavy stands	

NOTE: Use higher rate for dense or weedy crops

[^] WETTING AGENT: Add Agral® at the rate of 200 mL/100L or Imtrade Penetrate Wetter at 160 mL/100L of prepared spray unless otherwise specified.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

**GRAZING -
DO NOT GRAZE OR CUT SPRAYED VEGETATION FOR STOCK FOOD FOR 1 DAY AFTER APPLICATION**

**HARVEST -
Cotton, Dry beans, Dry peas, Mung Beans, Asparagus, Hops, Orchards and Vineyards, Row Crops, Vegetables and Market Gardens, Oats, Wheat and Winter Cereals: NOT REQUIRED WHEN USED AS DIRECTED**

**Lentils, Chickpeas and Faba Beans: DO NOT HARVEST FOR 2 DAYS AFTER APPLICATION
Pigeon Peas, Canola, Sunflower, Soya Beans, Sugar Cane: DO NOT HARVEST FOR 4 DAYS AFTER APPLICATION**

Rice: DO NOT HARVEST FOR 5 DAYS AFTER APPLICATION

Potatoes: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION

Sweet Potatoes: DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION

Poppies: DO NOT HARVEST FOR 16 DAYS AFTER APPLICATION

DO NOT USE TREATED WATER FOR HUMAN CONSUMPTION, LIVESTOCK WATERING OR IRRIGATION PURPOSES FOR 10 DAYS AFTER APPLICATION.

GENERAL INSTRUCTIONS

Uses
Imtrade Diquat 200 Non-Residual Herbicide is an aqueous solution of diquat, a non-volatile herbicide with unique properties. It very quickly kills green growth with which it comes into contact and is particularly effective against broadleaved weeds. It is inactivated on contact with the soil and crop roots and seeds below the soil remain unharmed. It can be safely applied around bushes and trees which have no green bark. It is non-volatile, easily mixed with water and active at low concentrations.

Mixing
Add the required quantity of Imtrade Diquat 200 Non-Residual Herbicide to water in the spray tank and agitate to give even mixing. Agitate again if left standing. Use clean water only, as suspended soil particles in dirty water will interfere with herbicidal action.

Wetting Agent
Imtrade Diquat 200 Non-Residual Herbicide contains no wetting agent, and a non-ionic wetting agent must be added to the spray mixture unless otherwise specified. Add Agral® at the rate of 200 mL/100L or Imtrade Penetrate Wetter at 160 mL/100L of prepared spray unless otherwise specified.

Application
For best result an even and complete coverage and good penetration of the spray into the target foliage is necessary. Best results will be obtained when application is made in dull weather or at the end of the day. Imtrade Diquat 200 Non-Residual Herbicide is rapidly absorbed and is not affected by rain falling shortly after application.

Application Rates
Use the higher rates specified in the directions for use for dense or weedy crops. For application to seedling weeds Imtrade Diquat 200 Non-Residual Herbicide is generally recommended at 1.4L/ha and Imtrade Paraquat 250 Herbicide at 1.2L/ha. Use Imtrade Diquat 200 Non-Residual Herbicide at 2.8 to 4L/ha and Imtrade Paraquat 250 Herbicide at 1.6 to 3.2L/ha when weeds are at the older stages of growth. Imtrade Paraquat 250 Herbicide is preferred where grasses are dominant and Imtrade Diquat 200 Non-Residual Herbicide where there are mainly broadleaf weeds.

Boom Spraying

A boomsprayer fitted with flat fan nozzle is preferred to ensure even coverage and to minimize drift. The boom should be set at sufficient height above the crop to provide a complete double overlap of the flat spray pattern. Spray drop arms on booms are useful for dense crops such as potatoes. A minimum spray volume of 100L/ha is recommended. Aim for a spray quality in the fine to medium range, Generally a flat fan nozzle operated at 200 to 300 kPa is preferred.

High Volume Spot Spraying

Hand held equipment use 250 mL of product per 100L of water and spray to visible wetness (about 700 to 1000L/ha). Use 50 mL product plus 30 mL Agral® per 15L knapsack.

Aerial Application

Flying height, pressure, nozzle size and positioning on the aircraft should be such as to minimize spray drift. Apply 30 to 60L of spray per hectare. Avoid spraying in high winds or under temperature inversion conditions. Wash any spillage during filling of the aircraft and make sure there are not leaks in the spraying system. Inspect the aircraft regularly for signs of corrosion and ensure the paint work is in good condition.

Caution – Use by Aircraft

Although this product is no different in drift behavior from other chemicals, it has rapid spotting effect on green foliage and, as with all herbicides; special care must be taken to avoid drift on to adjacent crops. Aircraft operators must not apply during periods of thermal (temperature) instability, and should avoid wind conditions and flying heights conducive to drift.

WEED CONTROL IN ROW CROPS, VEGETABLES AND MARKET GARDENS

Pre-planting and pre-crop emergence

To control weeds in seed beds before sowing, or post-sowing pre-crop emergence, apply as a blanket spray with this product using boom spray equipment or knapsack sprayers.

Post-emergence inter-row weed control

Use shielded nozzles for rapid control of weeds in inter-row spaces of row crops, after crop seedlings have emerged, or when transplanted crops are established. Direct the spray so that it does not touch the crop.

Pre-harvest crop desiccation

Green crop foliage and weeds can seriously interfere with harvesting operations of a number of crops. Imtrade Diquat 200 Non-Residual Herbicide can be used to facilitate harvesting by desiccating weeds, accelerating the drying of crops and reducing the moisture content of seeds. Drying costs are reduced, harvesting delays and associated risks avoided.

WARNING

Markers – If possible fixed markers should be used. Human markers are not recommended unless flaggers are protected by engineering controls such as vehicles with cabs.

COMPATIBILITY

This product mixes readily with ChemAg Paraquat 250 Herbicide, the soil residual herbicides Atrazine granules, Diuron 900 g/kg granules and Simazine Granules where prolonged weed control is required as well as a quick knockdown.

RESISTANT WEEDS WARNING

GROUP 22 HERBICIDE

Imtrade Diquat 200 Non-Residual Herbicide is a member of the bipyridyl group of herbicides. Imtrade Diquat 200 Non-Residual Herbicide has the inhibitor of photo-synthesis at photosystem I mode of action. For weed resistance management Imtrade Diquat 200 Non-Residual Herbicide is a Group 22 Herbicide. Some naturally occurring weed biotypes resistant to Imtrade Diquat 200 Non-Residual Herbicide and other inhibitors or Group 22 Herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Imtrade Diquat 200 Non-Residual Herbicide or other inhibitors or Group 22 Herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use. Imtrade Australia Pty Ltd accepts no liability for any losses that may result from the failure of Imtrade Diquat 200 Non-Residual Herbicide to control resistant weeds.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions or from spraying equipment which may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

PROTECTION OF LIVESTOCK

Domestic pets and poultry – keep away from treated areas, Low hazard to bees. No special precautions are required.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or watercourses with the chemical or used containers.

STORAGE AND DISPOSAL

The method of disposal of the container depends on the container type. Read the Storage and Disposal instructions on the label that is attached to the container.

Store in the closed, original container in a dry, cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. Store in a locked room or a place away from children, animals, food, feedstuffs, seed and fertilizers.

All other non-refillable containers

Triple rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers and product.

Envirodrum - Micro Matic Valve

Store the original sealed Envirodrum in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. **DO NOT** tamper with the Micro Matic valve or the security seal. **DO NOT** contaminate the Envirodrum with water or any other foreign matter. After each use of the product please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase.

Refillable Containers

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Very dangerous. Poisonous if absorbed by skin contact, inhaled or swallowed. Will irritate the eyes, nose, throat and skin. Avoid contact with eyes and skin. **DO NOT** inhale spray mist. When preparing spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves, face shield or goggles, and half-face respirator or disposable respirator. If clothing becomes contaminated with product or wet with spray remove clothing immediately. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, respirator and if rubber wash with detergent and warm water, face shield and goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (*Phone Australia 13 11 26; New Zealand 0800 764 766*). If in eyes, hold open, flood with water for at least 15 minutes and see a doctor.

Additional GHS Hazard & Precautionary Statements

- Causes severe skin burns and eye damage; •May cause an allergic skin reaction; •May cause respiratory irritation; •Causes damage to organs through prolonged or repeated exposure; •Very toxic to aquatic life with long-lasting effects; •**DO NOT** breathe fumes, mists or vapours; •**DO NOT** get on clothing; •Wash contacted areas thoroughly after handling; •Use only outdoors or in a well-ventilated area; •Contaminated clothing should not be allowed out of the workplace; •Avoid release to the environment; •**IF SWALLOWED:** Rinse mouth. Do **NOT** induce vomiting; •**IF ON SKIN (or hair):** Remove immediately all contaminated clothing. Rinse skin with water; •**IF INHALED:** Remove victim to fresh air and keep at rest in a position comfortable for breathing; •If skin irritation or rash occurs: Get medical advice; •If eye irritation persists: Get medical advice; •Collect spillage; •Store locked up.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from your supplier or Imtrade Australia Pty Ltd website at www.imtrade.com.au

CONDITIONS OF SALE

Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Imtrade Australia Pty Ltd has any authority to add to or alter these conditions.

UN 1760

CORROSIVE LIQUID, N.O.S
(contains DIQUAT)

**IN A TRANSPORT EMERGENCY,
DIAL 000
POLICE OR FIRE BRIGADE**

PG III

HAZCHEM 2X



APVMA Approval No: 58411/141286

