**STORAGE AND DISPOSAL**
Store in the closed, original container in a dry, cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight.

***Gatekeeper® Drums - (20L)***
**DO NOT** tamper with the valve or the security seal while product remains in drum. After each use of the product please ensure that the valve coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained into the spray tank. When drum is empty, attach a water line to the hose coupling, fill 1/3 with water, invert the drum 3 times and pump the rinsings into the spray tank via the valve coupler. Repeat 3 times. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers and product. **DO NOT** re-use empty container for any other purpose.

**Envirodrum - Micro Matic Valve**
Store the original sealed Envirodrum in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. **DO NOT** tamper with the Micro Matic valve or the security seal. **DO NOT** contaminate the Envirodrum with water or any other foreign matter. After each use of the product please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase.

**Other Containers**
Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

**SAFETY DIRECTIONS**
Very dangerous, particularly the concentrate. **DO NOT** swallow. The product, particularly the concentrate, can kill if swallowed, absorbed through the eyes, or absorbed by skin contact. The liquid can cause burns particularly to the eyes. Will irritate the nose, throat and skin.When handling, **DO NOT** touch or rub eyes, nose or mouth with hand. Avoid contact with eyes and skin, open wounds, and clothing. Protect eyes while using.

If clothing becomes contaminated with product or with wet spray remove clothing immediately. **DO NOT** inhale spray mist. **DO NOT** allow children to play with containers or any equipment that is used.

When connecting, disconnecting and cleaning equipment wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and a washable hat, impervious footwear, elbow-length chemical resistant gloves and a full-face respirator with canister specified for paraquat/diquat OR half face-piece respirator with canister specified for paraquat/diquat and face shield or goggles.

When applying by hand wand wear cotton overalls, over normal clothing, buttoned to the neck and wrist and a washable hat, impervious footwear, elbow-length chemical resistant gloves and a full face piece respirator with a canister specified for paraquat/diquat.

After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, face shield or goggles, respirator (and if rubber wash with detergent and warm water) clothing and footwear.

**FIRST AID**
If poisoning occurs, get to a doctor or hospital quickly. If sprayed on skin, wash thoroughly. If sprayed in mouth, rinse mouth with water. If in eyes, hold open, flood with water for at least 15 minutes and see a doctor.

Refer to attached Leaflet for additional GHS Hazard & Precautionary Statements.

**SAFETY DATA SHEET**
Additional information is listed in the Safety Data Sheet, which can be obtained from your supplier or Imtrade Australia Pty Ltd website at www.imtrade.com.au

# POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY INSTRUCTIONS BEFORE OPENING OR USING

IMTRADE

# DIQUAT 200

## NON-RESIDUAL HERBICIDE

ACTIVE CONSTITUENTS: 200 g/L DIQUAT
present as DIQUAT DIBROMIDE MONOHYDRATE

**GROUP 22 HERBICIDE**

For pre-harvest crop desiccation in Poppies and control of broadleaf weeds in certain crops as per Directions for Use.

**IMPORTANT: Read the attached Leaflet before use.**

**GROUP 22** | **FORM AC** | **CONTENTS 5L-1000L**

**CONDITIONS OF SALE**
Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Imtrade Australia Pty Ltd has any authority to add to or alter these conditions.

GATEKEEPER® is a Registered Trademark of Imtrade Australia Pty Ltd

**UN 1760**

CORROSIVE LIQUID, N.O.S. (contains DIQUAT)

**In a Transport Emergency Dial 000**
**Police or Fire Brigade**

PG III | HAZCHEM 2X

**APVMA Approval No.**
**58411/154991**

drumMUSTER eligible container

Batch No.
DOM:

IMTRADE CropScience
17 Ocean St, Kwinana Beach WA 6167
08 9419 0333 | www.imtrade.com.au

# POISON

**KEEP OUT OF REACH OF CHILDREN**
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

IMTRADE

# DIQUAT 200

## NON-RESIDUAL HERBICIDE

**ACTIVE CONSTITUENT: 200 g/L DIQUAT**
**present as DIQUAT DIBROMIDE MONOHYDRATE**

**GROUP 22 HERBICIDE**

For pre-harvest crop desiccation in Poppies and
control of broadleaf weeds in certain crops
as per Directions for Use

**IMPORTANT: READ THIS LEAFLET THOROUGHLY**
**BEFORE OPENING OR USING THIS PRODUCT**

**APVMA Approval No: 58411/154991**

Imtrade Australia Pty Ltd
ABN 13 090 151 134
17 Ocean Street, Kwinana, Western Australia 6167
Tel: (08) 9419 0333 Fax: (08) 9419 7516
Web: www.imtrade.com.au

**RESTRAINTS:**
**DO NOT** remove contents except for immediate use.
**DO NOT** apply by spraying equipment carried on the back of the user.
**DO NOT** use open mixing/loading equipment. Closed mixing and loading **MUST** be used.
**DO NOT** continue to use if eye irritation or bleeding from the nose occurs.
**DO NOT** spray when weeds are under drought stress or when covered with dust or soil.
**DO NOT** apply if heavy rains or storms are forecast within 3 days.
**DO NOT** irrigate to the point of field runoff for at least 3 days after application.

**For small scale applications up to 6 ha per day:**
**DO NOT** apply using open cab equipment unless using a PF10 respirator.

**For applications greater than 6 ha per day:**
**DO NOT** apply using open cab equipment. Enclosed cab applications **MUST** be used.

**For hand spray applications:**
**DO NOT** use hand wand sprays by spraying out of the window of a vehicle.

**SPRAY DRIFT RESTRAINTS:**
Specific definitions for terms used in this section of the label can be found at apvma.gov.au/spraydrift

**DO NOT** allow **bystanders** to come into contact with the spray cloud.
**DO NOT** apply in a manner that may cause an unacceptable impact to **native vegetation**, **agricultural crops**, **landscaped gardens** and **aquaculture production**, or cause contamination of plant or livestock commodities, outside the **application site** from **spray drift**. The advisory buffer zones in the relevant buffer zone table/s below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.
**DO NOT** apply unless **wind speed** is between 3 and 20 kilometres per hour at the **application site** during the time of application.
**DO NOT** apply if there are **surface temperature inversion** conditions present at the **application site** during the time of application. These conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

**DO NOT** apply by a boom sprayer unless the following requirements are met:

- Spray droplets not smaller than a **MEDIUM** spray droplet size category.
- Minimum distances between the application site and downwind sensitive areas (see 'Mandatory buffer zones' section of the following table titled 'Buffer zones for boom sprayers') are observed.

**Buffer zones for boom sprayers**

| Application Rate | Boom height above the target canopy | Mandatory downwind buffer zones (metres) | | | | |
|---|---|---|---|---|---|---|
| | | Bystander areas | Natural aquatic areas | Pollinator areas | Vegetation areas | Livestock areas |
| Up to 10L/ha* | 0.5m or lower | 110 | 275 | 0 | 35 | 0 |
| Up to 5L/ha* | 0.5m or lower | 50 | 110 | | 20 | |
| | 1.0m or lower | 160 | 350 | | 60 | |
| Up to 4L/ha | 0.5m or lower | 40 | 85 | | 20 | |
| | 1.0m or lower | 130 | 250 | | 55 | |
| Up to 3L/ha | 0.5m or lower | 35 | 65 | | 10 | |
| | 1.0m or lower | 100 | 180 | | 45 | |
| Up to 1.5L/ha | 0.5m or lower | 20 | 40 | | 5 | |
| | 1.0m or lower | 60 | 110 | | 25 | |
| Up to 700mL/ha | 0.5m or lower | 5 | 20 | | 0 | |
| | 1.0m or lower | 35 | 60 | | 15 | |
| Up to 400mL/ha | 0.5m or lower | 0 | 10 | | 0 | |
| | 1.0m or lower | 20 | 10 | | 10 | |

*Aquatic uses only

**DO NOT** apply by a vertical sprayer.

**DO NOT** apply by aircraft unless the following requirements are met:

- Spray droplets not smaller than a **MEDIUM** spray droplet size category.
- For maximum release heights above the target canopy of 3m or 25% of wingspan or 25% of rotor diameter whichever is the greatest, minimum distances between the application site and downwind sensitive areas (see 'Mandatory buffer zones' section of the following table titled 'Buffer zones for aircraft') are observed.
- Application rate does not exceed 1.4L/ha.

**Buffer zones for aircraft**

| Type of aircraft | Mandatory downwind buffer zones (metres) | | | | |
|---|---|---|---|---|---|
| | Bystander areas | Natural aquatic areas | Pollinator areas | Vegetation areas | Livestock areas |
| **Fixed-wing up to 1.4L/ha** | 275 | 525 | 0 | 100 | 0 |
| **Helicopter up to 1.4L/ha** | 180 | 300 | | 80 | |
| **Fixed-wing up to 750mL/ha** | 170 | 275 | | 65 | |
| **Helicopter up to 750mL/ha** | 120 | 190 | | 50 | |

## DIRECTIONS FOR USE

### Pre-Harvest Crop Desiccation

| Crop | State | Rate | Critical Comments |
|---|---|---|---|
| **Poppies** | Tas only | 3 to 4L/ha^ | Spray after the poppies have reached the stripy capsule stage. Helps overcome slow and uneven ripening and weed problems at harvest. |

## General Weed Control

| Crop | Weeds Controlled | State | Rate | Critical Comments |
|---|---|---|---|---|
| **Aquatic areas:**<br>**Farm dams and artificial water bodies** | Duck Weeds, Red Azolla, Water Hyacinth, Salvinia, Marsilea, Water Lilies, Water Lettuce | All States | **DO NOT** apply to water bodies from which the product may contaminate natural aquatic areas.<br>**DO NOT** use treated water for human consumption, livestock watering or irrigation purposes for 10 days after application. | |
| | | | 5 or 10L/ha | Apply as an overall spray wetting foliage thoroughly. Clear water is necessary for best results as suspended soil particles interfere with herbicidal action. Use the higher rate for heavy infestations or for deep or dirty water. A repeat application 7 to 14 days later may be necessary for control of dense infestations. Oxygen depletion of decaying weeds may occur, therefore not more than ¼ of the area should be treated at once to ensure adequate oxygen supply for fish. |
| | | | 400mL plus 150mL Agral® per 100L water | Small areas – spray to wet weeds thoroughly. About 1mL of product should be sufficient to treat about $1m^2$. |
| | Cattail and Pond Weeds | | 5L/ megalitre water | Apply by injection below the surface or as a surface spray. |
| **Asparagus** | Broadleaf weeds | All States | 1.4L/ha plus 800mL Agral® in 400L water | Apply to control seedling weeds before spears have emerged. |
| **Hops** | Annual broadleaf and grass weeds | Vic and Tas only | 700mL to 1.4L/ha^ may be mixed with 1.1kg Simazine granules | Apply as a directed inter-row spray prior to crop emerging from winter dormancy, using a minimum of 250L/ha spray volume to ensure good and even coverage weeds. |

| Crop | Weeds Controlled | State | Rate | Critical Comments | |
|---|---|---|---|---|---|
| **Infested areas** | Cotton Thistle (*Onopordum acanthium*) | Tas only | 300L/ha plus 150mL Agral® in 100L water | Spot spray at the rosette stage before the centre shoot is 15cm tall. The spray should be applied to give complete wetting of the leaf surface. **DO NOT** use a lower rate or treat at a later growth stage. | |
| | Saffron Thistle | All States | 2.8L/ha plus 1L Agral® in 200L water/ha | Apply as an overall treatment to prevent seeding. | |
| **Lucerne** | Capeweed and *Erodium* spp. | All States | 350mL/ha^ in 200L water | Early Autumn application | Heavy grazing is necessary to reduce Lucerne to 2cm in height before spraying. |
| | | | 700mL/ha^ in 200L water | Late winter application | |
| **Oil Seed Poppies** | Weed control | Tas only | 300mL to 1.5L/ha | Use in accordance with recommendations made by Department of Primary Industries of the poppy contracting company. **DO NOT add Agral® or any other wetting agent to the spray solution.** | |
| **Berries and other small fruit (except Grapes)** | Broadleaf weeds | All States | 1.4L/ha^ | Apply as a shielded spray between crop rows. **DO NOT** allow spray to contact any part of the crop. | |
| **Broccoli, Head Cabbages, Cauliflower and Chinese Cabbage (type Pe-tsai)** | | | | Apply as a blanket spray prior to crop emergence. Once crops have emerged, or seedling has been transplanted, apply as a shielded spray between crop rows. **DO NOT** allow spray to contact any part of the crop. | |
| **Bulb Onions** | | | | | |
| **Fruiting Vegetables: other than Cucurbits** | | | | | |
| **Cucumbers** | | | | | |
| **Leafy Vegetables** | | | | | |
| **Legume Vegetables** | | | | | |
| **Root and Tuber Vegetables** | | | | | |

| Crop | Weeds Controlled | State | Rate | Critical Comments |
|---|---|---|---|---|
| **Wheat and Oats** | Capeweed (Small seedlings) | Qld, NSW, Vic, Tas, SA only | 550mL/ha in 200L of water | **DO NOT add wetting agent.** Spray when the crop is between the 4 (Wheat) or 3 (Oats) leaf and early tillering stage. |
| **Pasture Renovation and establishment** | Capeweed and *Erodium* spp. (Storksbill) | All States | 750mL to 940mL/ha plus Agral® in a minimum of 100L water | Apply by boom spray as an overall spray on 'run-down' pasture after heavy grazing. Pasture should not be greater than 4cm long when sprayed. Grazing should be carried out during previous spring, summer and early autumn.<br>Where Capeweed is in the very young seedling stage (2 or 3 true leaves only) rates may be reduced to 350mL/ha. Where Capeweed infestation is high, oversowing with new pasture seed by direct drilling is advisable. Direct drill 3 to 7 days after spraying using a pasture mixture suitable to the district. |
| **^ WETTING AGENT:** Add Agral® at the rate of 200mL/100L or Imtrade Penetrate Wetter at 160mL/100L of prepared spray unless otherwise specified. | | | | |

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIODS**
**GRAZING:**
**DO NOT GRAZE OR CUT SPRAYED VEGETATION FOR STOCK FOOD FOR 1 DAY AFTER APPLICATION**

**HARVEST:**
***Berries and other small fruits, Hops, Broccoli, Cauliflower, Head Cabbage, Chinese Cabbage, Bulb Onions, Fruiting Vegetables (other than Cucurbits), Cucumbers, Leafy Vegetables, Legume Vegetables, Root and Tuber Vegetables, Wheat and Oats:*** **NOT REQUIRED WHEN USED AS DIRECTED.**
***Poppies:*** **DO NOT HARVEST FOR 2 DAYS AFTER APPLICATION**

**EXPORT OF TREATED PRODUCE**
Growers should note that MRLs or import tolerances do not exist in all markets for edible produce treated with IMTRADE DIQUAT 200 NON-RESIDUAL HERBICIDE. If you are growing edible produce for export, please check with Imtrade Australia Pty Ltd for the latest information on MRLs and import tolerances BEFORE using IMTRADE DIQUAT 200 NON-RESIDUAL HERBICIDE.

## GENERAL INSTRUCTIONS

IMTRADE DIQUAT 200 NON-RESIDUAL HERBICIDE is an aqueous solution of diquat, a non-volatile herbicide with unique properties. It very quickly kills green growth with which it comes into contact and is particularly effective against broadleafed weeds. It is inactivated on contact with the soil and crop roots and seeds below the soil remain unharmed. It can be safely applied around bushes and trees which have no green bark. It is non-volatile, easily mixed with water and active at low concentrations.

## MIXING

Add the required quantity of IMTRADE DIQUAT 200 NON-RESIDUAL HERBICIDE to water in the spray tank and agitate to give even mixing. Agitate again if left standing. Use clean water only, as suspended soil particles in dirty water will interfere with herbicidal action.

## WETTING AGENT

IMTRADE DIQUAT 200 NON-RESIDUAL HERBICIDE contains no wetting agent, and a non-ionic wetting agent must be added to the spray mixture unless otherwise specified. Add Agral® at the rate of 200mL/100L or Imtrade Penetrate Wetter at 160mL/100L of prepared spray unless otherwise specified.

## APPLICATION

For best result an even and complete coverage and good penetration of the spray into the target foliage is necessary. Best results will be obtained when application is made in dull weather or at the end of the day. IMTRADE DIQUAT 200 NON-RESIDUAL HERBICIDE is rapidly absorbed and is not affected by rain falling shortly after application.

### Application Rates

Imtrade Paraquat 250 Herbicide is preferred where grasses are dominant and IMTRADE DIQUAT 200 NON-RESIDUAL HERBICIDE where there are mainly broadleaf weeds.

## BOOM SPRAYING

A boom sprayer fitted with flat fan nozzle is preferred to ensure even coverage and to minimize drift. The boom should be set at sufficient height above the crop to provide a complete double overlap of the flat spray pattern. Spray drop arms on booms are useful for dense crops. A minimum spray volume of 100L/ha is recommended.

### High Volume Spot Spraying

**Hand held equipment** - use 250mL of product per 100L of water and spray to visible wetness (about 700 to 1000L/ha).

## AERIAL APPLICATION

Flying height, pressure, nozzle size and positioning on the aircraft should be such as to minimize spray drift. Apply 30 to 60L of spray per hectare. Avoid spraying in high winds or under temperature inversion conditions. Wash any spillage during filling of the aircraft and make sure there are no leaks in the spraying system. Inspect the aircraft regularly for signs of corrosion and ensure the paintwork is in good condition.

## CAUTION – USE BY AIRCRAFT

Although this product is no different in drift behavior from other chemicals, it has rapid spotting effect on green foliage and, as with all herbicides, special care must be taken to avoid drift into adjacent crops. Aircraft operators must not apply during periods of thermal (temperature) instability and should avoid wind conditions and flying heights conducive to drift.

**WEED CONTROL IN** Berries and other small fruit (except Grapes), Broccoli, Head Cabbages, Cauliflower and Chinese Cabbage (type Pe-tsai), Bulb Onions, Fruiting Vegetables (other than Cucurbits), Cucumbers, Leafy Vegetables, Legume Vegetables, Root and Tuber Vegetables.

### Pre-planting and pre-crop emergence

To control weeds in seed beds before sowing, or post-sowing pre-crop emergence, apply as a blanket spray with this product using boom spray equipment or handheld sprayers.

### Post-emergence inter-row weed control

Use shielded nozzles for rapid control of weeds in inter-row spaces of row crops, after crop seedlings have emerged, or when transplanted crops are established. Direct spray so that it does not touch the crop.

## COMPATIBILITY

For information on the compatibility of IMTRADE DIQUAT 200 NON-RESIDUAL HERBICIDE with other products, contact your local Imtrade Australia Pty Ltd representative.

## RESISTANT WEEDS WARNING

**GROUP 22 HERBICIDE**

IMTRADE DIQUAT 200 NON-RESIDUAL HERBICIDE is a member of the bipyridyl group of herbicides. IMTRADE DIQUAT 200 NON-RESIDUAL HERBICIDE has the inhibitor of photo-synthesis at photosystem I mode of action. For weed resistance management IMTRADE DIQUAT 200 NON-RESIDUAL HERBICIDE is a Group **22** Herbicide. Some naturally occurring weed biotypes resistant to IMTRADE DIQUAT 200 NON-RESIDUAL HERBICIDE and other inhibitors or Group **22** Herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by IMTRADE DIQUAT 200 NON-RESIDUAL HERBICIDE or other inhibitors or Group **22** Herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use. Imtrade Australia Pty Ltd accepts no liability for any losses that may result from the failure of IMTRADE DIQUAT 200 NON-RESIDUAL HERBICIDE to control resistant weeds.

## PRECAUTIONS

### Re-Entry period

**DO NOT** allow entry to treated areas until the spray has dried unless using an enclosed cab or wearing cotton overalls and gloves. Workers performing scouting and hand-set irrigation activities must comply with the Re-entry periods specified below:

| | Crop | |
|---|---|---|
| **Rate** | **Scouting (Days)** | **Irrigation (Days)** |
| Up to 700mL/ha | 0 | 0 |
| Up to 1.4L/ha | 0 | 0 |
| Up to 2.8L/ha | 1 | 7 |
| Up to 4L/ha | 4 | 9 |

## PROTECTION OF LIVESTOCK

**Domestic pets and poultry** – keep away from treated areas.

## PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

**Very toxic to aquatic life.**

**DO NOT** contaminate wetlands or watercourses with this product or used containers.

### Toxic to birds

However, the use of this product as directed is not expected to have adverse effects on birds and native mammals.

### Toxic to beneficial arthropods.

Not compatible with Integrated Pest Management (IPM) programs utilizing beneficial arthropods. Minimise spray drift to reduce harmful effects on beneficial arthropods in non-crop areas.

## STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight.

### *Gatekeeper® Drums - (20L)*

**DO NOT** tamper with the valve or the security seal while product remains in drum. After each use of the product please ensure that the valve coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained into the spray tank. When drum is empty, attach a water line to the hose coupling, fill 1/3 with water, invert the drum 3 times and pump the rinsings into the spray tank via the valve coupler. Repeat 3 times. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers and product. **DO NOT** re-use empty container for any other purpose.

### Envirodrum - Micro Matic Valve

Store the original sealed Envirodrum in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. **DO NOT** tamper with the Micro Matic valve or the security seal. **DO NOT** contaminate the Envirodrum with water or any other foreign matter. After each use of the product please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase.

### Other Containers

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

## SAFETY DIRECTIONS

Very dangerous, particularly the concentrate. **DO NOT** swallow. The product, particularly the concentrate, can kill if swallowed, absorbed through the eyes, or absorbed by skin contact. The liquid can cause burns particularly to the eyes. Will irritate the nose, throat and skin. When handling, **DO NOT** touch or rub eyes, nose or mouth with hand. Avoid contact with eyes and skin, open wounds, and clothing. Protect eyes while using.

If clothing becomes contaminated with product or with wet spray remove clothing immediately. **DO NOT** inhale spray mist. **DO NOT** allow children to play with containers or any equipment that is used.

When connecting, disconnecting and cleaning equipment wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and a washable hat, impervious footwear, elbow-length chemical resistant gloves and a full-face respirator with canister specified for paraquat/diquat OR half face-piece respirator with canister specified for paraquat/diquat and face shield or goggles.

When applying by hand wand wear cotton overalls, over normal clothing, buttoned to the neck and wrist and a washable hat, impervious footwear, elbow-length chemical resistant gloves and a full face piece respirator with a canister specified for paraquat/diquat.

After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, face shield or goggles, respirator (and if rubber wash with detergent and warm water) clothing and footwear.

## FIRST AID

If poisoning occurs, get to a doctor or hospital quickly. If sprayed on skin, wash thoroughly. If sprayed in mouth, rinse mouth with water. If in eyes, hold open, flood with water for at least 15 minutes and see a doctor.

### Additional GHS Hazard & Precautionary Statements

•Causes severe skin burns and eye damage; •May cause an allergic skin reaction; •May cause respiratory irritation; •Causes damage to organs through prolonged or repeated exposure; •Very toxic to aquatic life with long-lasting effects; •**DO NOT** breathe fumes, mists or vapours; •**DO NOT** get on clothing; •Wash contacted areas thoroughly after handling; •Use only outdoors or in a well-ventilated area; •Contaminated clothing should not be allowed out of the workplace; •Avoid release to the environment; •**IF SWALLOWED:** Rinse mouth. Do **NOT** induce vomiting; •**IF ON SKIN (or hair):** Remove immediately all contaminated clothing. Rinse skin with water; •**IF INHALED:** Remove victim to fresh air and keep at rest in a position comfortable for breathing; •If skin irritation or rash occurs: Get medical advice; •If eye irritation persists: Get medical advice; •Collect spillage; •Store locked up.

## SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from your supplier or Imtrade Australia Pty Ltd website at www.imtrade.com.au

## CONDITIONS OF SALE

Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Imtrade Australia Pty Ltd has any authority to add to or alter these conditions.

GATEKEEPER® is a Registered Trademark of Imtrade Australia Pty Ltd

| | |
|---|---|
| **UN 1760** | **CORROSIVE LIQUID, N.O.S (contains DIQUAT)** |
| **IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE** | |
| **PG III** | **HAZCHEM 2X** |




**APVMA Approval No: 58411/154991**