

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

STORAGE AND DISPOSAL (5L, 10L & 20L)

Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or an approved waste management facility. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers and product.

Envirodrum 110L Mini Bulk Returnable Container

Store the closed, original sealed Envirodrum in a well-ventilated area, as cool as possible and locked away from children, animals, food, foodstuffs, seed and fertilisers. **DO NOT** store for prolonged periods in direct sunlight. **DO NOT** tamper with the Micro Matic valve or the security seal. **DO NOT** contaminate the Envirodrum with water or any other foreign matter. After each use of the product, please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase.

Refillable containers (1000L only)

Store in the closed, original container in a well-ventilated area, as cool as possible and locked away from children, animals, food, foodstuffs, seed and fertilisers. **DO NOT** store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and preparing spray, wear elbow-length PVC gloves. Wash hands after use. After each day's use, wash gloves. When tank mixing with Imtrade Cyclone 330 EC Herbicide or Imtrade Paraquat 250 Herbicide, consult also the safety directions for those products.

IMTRADE EMBER 240 SL HERBICIDE

ACTIVE CONSTITUENT: 240 g/L IMAZAPIC (present as the ammonium salt)

GROUP 2 HERBICIDE

For the pre-emergence control of certain annual grass and broadleaf weeds in fallow situations, sugarcane and peanuts and early post-emergence control of certain annual grass and broadleaf weed in peanuts and sugarcane as specified in the Directions for Use section of this label.

IMPORTANT: Read the attached Leaflet before use.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre (Phone Australia 13 11 26).

Refer to attached Leaflet for additional GHS
Hazard & Precautionary Statements

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from your supplier or Imtrade Australia Pty Ltd website at www.imtrade.com.au

CONDITIONS OF SALE

Imtrade Australia Pty Ltd warrants that this product conforms to the chemical description on the label. The use of Imtrade Ember 240 SL Herbicide being beyond the control of the manufacturer, Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Imtrade Australia Pty Ltd has any authority to add to or alter these conditions.

In a Transport Emergency Dial 000
Police or Fire Brigade

APVMA Approval No: 64943/106765

GROUP
2

FORM
SL

CONTENTS
1L-1000L



Batch No.
DOM:

IMTRADE
CropScience

17 Ocean St, Kwinana Beach WA 6167
08 9419 0333 | www.imtrade.com.au

READ SAFETY DIRECTIONS BEFORE OPENING OR USING



IMTRADE EMBER 240 SL HERBICIDE

**ACTIVE CONSTITUENT: 240 g/L IMAZAPIC
(present as the ammonium salt)**

GROUP	2	HERBICIDE
--------------	----------	------------------

For the pre-emergence control of certain annual grass and broadleaf weeds in fallow situations, sugarcane and peanuts and early post-emergence control of certain annual grass and broadleaf weed in peanuts and sugarcane as specified in the Directions for Use section of this label.

**IMPORTANT: READ THIS LEAFLET THOROUGHLY
BEFORE OPENING OR USING THIS PRODUCT**

APVMA Approval No. 64943/106765

Imtrade Australia Pty Ltd
ABN 13 090 151 134
17 Ocean Street, Kwinana Western Australia 6167
Tel: 1800 171 799 Fax: 1800 17 1788
Web: www.imtrade.com.au

RESTRAINTS

DO NOT apply to soils saturated with water.

DO NOT apply by aircraft.

DIRECTIONS FOR USE

Situation	Weeds Controlled	Rate /ha	Critical Comments
Fallow prior to planting Winter Crops as directed in the Follow Crops table (Qld, Nth NSW only)	Barnyard Grass (<i>Echinochloa colona</i>), Liverseed Grass (<i>Urochloa panicoides</i>), Stink Grass (<i>Eragrostis ciliaris</i>), Blowaway Grass (<i>Panicum decompositum</i>), Button Grass (<i>Dactyloctenium radulans</i>), Pigweed (<i>Portulaca oleracea</i>), Yellow Vine (<i>Tribulus terrestris</i>), Mintweed (<i>Salvia reflexa</i>), Boggabri weed (<i>Amaranthus mitchelli</i>), Dwarf Amaranth (<i>Amaranthus macrocarpus</i>), Peachvine/Cowvine (<i>Ipomea lonchophylla</i>)	150 to 200 mL	Treatments may be applied from the end of the previous crop harvest but not less than 3 months prior to planting of winter crop (refer to Follow Crops table). Best results are obtained where treatments are applied to dry, weed-free soil prior to weed germination. Treatments can be applied to standing crop residues. Sufficient rainfall is required after application and prior to weed emergence to wet soil to a depth of 5 cm. Use the higher rates for increased residual control of weeds or anticipated high weed pressure. Emergent weeds must be controlled by prior cultivation, or by application of a knockdown herbicide. Where infestation of emergent weeds is light, tank mix treatment with glyphosate or paraquat. <i>continued next page ...</i>

Situation	Weeds Controlled	Rate /ha	Critical Comments
Fallow prior to planting Winter Crops as directed in the Follow Crops table (Qld, Nth NSW only) <i>(Cont'd)</i>	<i>Refer to previous page</i>	150 to 200 mL	<i>continued from previous page...</i> Cultivation following application of treatments may reduce pre-emergence weed control provided by Imtrade Ember 240 SL Herbicide. Prolonged wet soil conditions following treatment will reduce the effective life of pre-emergence treatments. Weed escapes may require follow-up application of knockdown herbicides. DO NOT apply more than 200 mL/ha per season on cropland. DO NOT apply in more than two successive seasons on cropland. DO NOT overspray headlands.
Sugarcane (Qld, NSW and WA only)	Awnless Barnyard Grass (<i>Echinochloa colona</i>), Barnyard Grass (<i>E. crusgalli</i>), Blackberry Nightshade (<i>Solanum nigrum</i>), Blue Billygoat Weed (<i>Ageratum houstonianum</i>), Common Sida (<i>Sida rhombifolia</i>), <i>continued next page...</i>	300 to 400 mL	DO NOT apply more than once a year to the same crop. Pre-emergence application: PLANT CANE: Apply after planting and before emergence of the cane. On sandy soil, ensure thorough coverage of sets to a minimum depth of 5cm. RATOON: Apply after harvest and before new cane growth appears. Use the higher rates of Imtrade Ember 240 SL Herbicide for increased residual control of weeds or anticipated high weed pressure. <i>continued next page...</i>

Situation	Weeds Controlled	Rate /ha	Critical Comments
Sugarcane (Qld, NSW and WA only) (Cont'd)	<i>Refer to previous page</i> Giant/Black Pigweed <i>(Trianthema portulacastrum)</i> , Glossy Nightshade <i>(Solanum americanum)</i> , Green Amaranth <i>(Amaranthus viridis)</i> , Green Summer Grass <i>(Brachiaria subquadrifida)</i> , <i>lipomoea</i> spp.), Guinea Grass <i>(Panicum maximum)</i> Liverseed Grass <i>(Urochloa panicoides)</i> , Milkweed (Mexican Fire Plant) <i>(Euphorbia heterophylla)</i> , Pigweed <i>(Portulaca oleracea)</i> , Summer Grass <i>(Digitaria ciliaris)</i>	300 to 400 mL	<i>continued from previous page...</i> Heavy trash layers may prevent Imtrade Ember 240 SL Herbicide from reaching the soil. Apply to weed-free soil prior to weed germination. Sufficient rainfall, or overhead irrigation, is required after application and prior to weed emergence to wet soil to a minimum depth of 5cm. Emerged weeds must be controlled by prior cultivation, or application of an appropriate herbicide. Weed control can be improved under dry conditions or where incorporating moisture has been insufficient, by cultivating the treated soil above setts. Care needs to be taken to minimise exposure of untreated soil or setts. Pre-emergence weed control may be limited on Krasnozem or red brown Ferrosol soil types, where moisture is not maintained in the top 5cm of soil. Refer to Soil Type in the General Instructions. Heavy rain and/or irrigation within 2 days following application may concentrate Imtrade Ember 240 SL Herbicide in the furrow and cause temporary yellowing and stunting of cane leaves. Prolonged wet soil and/or cool conditions can increase crop damage, especially in sandy soil. Normal growth will return when conditions become warm and drier. Weed escapes may require follow-up application of an appropriate herbicide. Contact Imtrade Australia Pty Ltd for advice on variety susceptibility before using Imtrade Ember 240 SL Herbicide (refer to the CROP SAFETY section of the label).

Situation	Weeds Controlled	Rate /ha	Critical Comments
Sugarcane (Qld, NSW and WA only) (Cont'd)	As per previous page plus Crowsfoot Grass (<i>Eleusine indica</i>)	400 mL plus 3L Imtrade Cyclone 330 EC Herbicide or 2.25L Cyclone 440EC Herbicide	Incorporate into the soil within 7 days of application, either by overhead irrigation, rainfall or mechanical incorporation, to a minimum depth of 5cm. Read all directions for use in Sugarcane as well as the Restraints and Follow Crop restrictions on the Imtrade Cyclone 330 EC Herbicide or Cyclone 440EC Herbicide label.
	As per previous page and the above plus control of those weeds expected with these rates of 250 g/L Paraquat products alone.	300 to 400 mL plus 1.2 to 1.6L of 250 g/L Paraquat products (e.g. Imtrade Paraquat 250 Herbicide)	<p>Post-emergence application (in emerged cane): ALWAYS tank mix Imtrade Ember 240 SL Herbicide with registered and appropriate rates of paraquat for the emerged weeds. Use the higher rates of Imtrade Ember 240 SL Herbicide for increased residual control of weeds or anticipated high weed pressure. Use the higher rates of paraquat for dense, more mature weeds. Apply as a carefully directed inter-row spray to weeds and at the base of the plant, minimising contact with sugarcane foliage. Contact with sugarcane foliage may cause temporary crop damage observed as inter-veinal yellowing and reduction in crop biomass or crop height.</p> <p><i>continued on next page ...</i></p>

Situation	Weeds Controlled	Rate /ha	Critical Comments
<p>Sugarcane (Qld, NSW and WA only) (Cont'd)</p>	<p>As above plus control of those weeds expected with these rates of 250 g/L Paraquat products alone.</p>	<p>300 to 400 mL plus 1.2 to 1.6L of 250 g/L Paraquat products (e.g. Imtrade Paraquat 250 Herbicide)</p>	<p><i>continued from previous page ...</i></p> <p>Refer to Imtrade Paraquat 250 Herbicide (or equivalent) label re spray boom types and precautions. Reduce the incidence of "fines" by using the correct nozzles and operating pressures. Replace worn or old nozzles and calibrate equipment on a regular basis.</p> <p>DO NOT cultivate within 1 hour of application.</p> <p>Heavy rain and/or irrigation within 2 days following application may concentrate Imtrade Ember 240 SL Herbicide in the furrow and cause temporary yellowing and stunting of cane leaves.</p> <p>DO NOT spray plants covered with heavy dew.</p> <p>Prolonged wet soil and/or cool conditions can increase crop damage, especially in sandy soil. Normal growth will return when conditions become warm and drier.</p> <p>Pre-emergence weed control may be limited on Krasnozem or red brown Ferrosol soil types, where moisture is not maintained in the top 5cm of soil. Refer to Soil Type in the General Instructions.</p> <p>Contact Imtrade Australia Pty Ltd for advice on variety susceptibility before using Imtrade Ember 240 SL Herbicide (refer to the CROP SAFETY section of the label).</p>

Situation	Weeds Controlled	Rate /ha	Critical Comments
Peanuts (Qld, NSW and NT only)	Awnless Barnyard Grass <i>(Echinochloa colona)</i> , Barnyard Grass <i>(E. crusgalli)</i> , Blackberry Nightshade <i>(Solanum nigrum)</i> , Blue Billygoat Weed <i>(Ageratum houstonianum)</i> , Common Sida <i>(Sida rhombifolia)</i> , Giant/Black Pigweed <i>(Trianthema portulacastrum)</i> , Glossy Nightshade <i>(Solanum americanum)</i> , Green Amaranth <i>(Amaranthus viridis)</i> , Green Summer Grass <i>(Brachiaria subquadriflora)</i> , <i>lipomoea</i> spp.), Guinea Grass <i>(Panicum maximum)</i> , Liverseed Grass <i>(Urochloa panicoides)</i> , Milkweed (Mexican Fire Plant) <i>(Euphorbia heterophylla)</i> , Pigweed <i>(Portulaca oleracea)</i> , Summer Grass <i>(Digitaria ciliaris)</i>	300 to 400 mL	Pre-emergence application: Optimum weed control will occur when moisture is maintained in the top 5cm of soil. Apply to weed-free soil prior to weed germination. Sufficient rainfall, or overhead irrigation, is required after application and prior to weed emergence to wet soil to a minimum depth of 5cm. Use the higher rates for increased residual control of weeds or anticipated high weed pressure. Pre-emergence weed control may be limited on Krasnozern or red brown Ferrosol soil types, where moisture is not maintained in the top 5cm of soil. Refer to Soil Type in the General Instructions. DO NOT apply to any one crop more than once per season.
	As above plus Nutgrass* <i>(Cyperus rotundus)</i>	400 mL plus Hasten® or Kwicken® at 1L/100L (or a similar esterified oil adjuvant)	Post-emergence application: Apply prior to crop flowering. DO NOT apply to any one crop more than once per season. *Nutgrass: suppression only when applied to nutgrass at the 3 to 4 leaf stage.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

HARVEST: NOT REQUIRED WHEN USED AS DIRECTED.

GRAZING/STOCKFEED:

WHEAT, BARLEY, CHICKPEA, PEANUTS: DO NOT GRAZE TREATED VEGETATION, STUBBLE OR WEEDS, OR CUT FOR STOCKFOOD, FOR 4 WEEKS AFTER APPLICATION.

SUGARCANE: DO NOT GRAZE CROP OR SPRAYED VEGETATION, OR CUT FOR STOCKFOOD, FOR 6 WEEKS AFTER APPLICATION.

GENERAL INSTRUCTIONS

All Applications:

Reduce the incidence of “fines” by using the correct nozzles and operating pressures. Replace worn or old nozzles and calibrate equipment on a regular basis.

Fallow Application:

Imtrade Ember 240 SL Herbicide can be applied in fallow prior to the planting of various crops as outlined in the Fallow Crops table. Imtrade Ember 240 SL Herbicide is a pre-emergence herbicide, which is recommended to provide short term (6 to 12 weeks) pre-emergence control of summer germinating grasses and broadleaved weeds in cultivated or uncultivated fallow. A green weed cover of greater than 50% will significantly reduce the length of pre-emergence weed control.

Sugarcane Application:

Pre-emergence

Where possible apply to soil two days prior to incorporation. Incorporate with adequate irrigation or rainfall to a minimum depth of 5cm. Incorporation by flood irrigation may result in loss of weed control. Imtrade Ember 240 SL Herbicide should be applied to plant or ratoon cane before the cane has emerged. When applying Imtrade Ember 240 SL Herbicide in moist conditions, allow weeds to germinate prior to application and apply mixed with an appropriate knockdown herbicide. Whenever emerged weeds are present, Imtrade Ember 240 SL Herbicide should be mixed with an appropriate knockdown herbicide.

Post-emergence

Imtrade Ember 240 SL Herbicide **MUST** always be mixed with registered rates of paraquat (i.e. Imtrade Paraquat 250 Herbicide) and applied as a carefully directed inter-row spray to weeds and at the base of the sugarcane stalks, minimising spray contact with sugarcane foliage. **DO NOT** add crop oil to this mixture. Other surfactant or adjuvant is not required.

Peanut Application:

Pre-emergence

Apply to soil and incorporate with adequate irrigation or rainfall to a minimum depth of 5cm.

Post-emergence

Imtrade Ember 240 SL Herbicide **MUST** always be mixed with an adjuvant as directed in the DIRECTIONS FOR USE section of the label.

Soil Type

Pre-emergence weed control with Imtrade Ember 240 SL Herbicide can be limited when applied onto soils with pH <5.0 and/or which contain high concentrations of iron (Fe) and/or aluminium (Al). These situations are common in soil types such as Krasnozems and red brown Ferrosols.

MIXING

Imtrade Ember 240 SL Herbicide is an aqueous solution. Part fill the spray tank with water, then with agitator running, add the required amount of product, then fill the tank with water. When tank mixing Imtrade Ember 240 SL Herbicide with other recommended compatible products, first add the other product to the tank and mix thoroughly before adding Imtrade Ember 240 SL Herbicide. For post-emergence applications in peanuts, add adjuvant after the addition of Imtrade Ember 240 SL Herbicide to the tank.

No adjuvants or surfactants are required when Imtrade Ember 240 SL Herbicide is used alone for pre-emergence weed control or with paraquat. **DO NOT** add crop oil when applied post-emergence with paraquat.

APPLICATION

DO NOT apply by aircraft.

Fallow application:

Apply with boom equipment in not less than 50L/ha water using flat fan nozzles. Avoid overlap and **DO NOT** overspray headlands.

Sugarcane:

Ensure thorough coverage of soil and weeds with well-calibrated spray equipment. Apply in a minimum of 200L water/ha using a nozzle configuration to generate a medium to coarse spray pattern for pre-emergence applications or medium spray pattern for post-emergence applications. Replace worn or old nozzles and calibrate equipment on a regular basis.

Peanuts:

Apply with boom equipment in not less than 50L/ha water using flat fan nozzles. Avoid overlap and **DO NOT** overspray headlands. For post-emergence applications, Imtrade Ember 240 SL Herbicide should be applied a minimum of 2 hours before rainfall.

EQUIPMENT CLEAN-UP

Thoroughly flush all spray equipment with water following the use of Imtrade Ember 240 SL Herbicide and before use with other products.

COMPATIBILITY

Imtrade Ember 240 SL Herbicide is compatible with Pendimethalin, Atrazine, Glyphosate, Paraquat, Paraquat & Diquat, Metsulfuron, Triclopyr, Touchdown® and 2,4-D.

It is **NOT** compatible with Dicamba.

RESISTANT WEEDS WARNING

GROUP	2	HERBICIDE
--------------	----------	------------------

Imtrade Ember 240 SL Herbicide is a member of the imadazolinones group of herbicides. Imtrade Ember 240 SL Herbicide has the inhibitors-of-acetolactate-synthase (ALS) mode of action. For weed resistance management, Imtrade Ember 240 SL Herbicide is a Group 2 Herbicide. Some naturally occurring weed biotypes, resistant to the product and other inhibitors-of-acetolactate-synthase (ALS) herbicides, may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population, if these herbicides are used repeatedly. These resistant weeds will not be controlled by Imtrade Ember 240 SL Herbicide or other inhibitors-of-acetolactate-synthase (ALS) herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Imtrade Australia Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant weeds.

FOLLOW CROPS

Under conditions, which do not favour breakdown of Imtrade Ember 240 SL Herbicide, carry-over soil residues can affect susceptible follow crops. As environmental and agronomic factors make it impossible to eliminate all risks associated with Imtrade Ember 240 SL Herbicide, rotational crop injury is always possible. The following minimum re-cropping intervals (months after application) should be observed

Following use in Summer Fallow:

MONTHS AFTER APPLICATION				
3	4	10	24	36
Faba Beans, Peanuts, Mungbeans, maize varieties with CLEARFIELD Technology ONLY : - Pacific Hycorn 62IT - Pacific Hycorn 53IT - Pioneer 3395IR; wheat varieties with CLEARFIELD Technology only:	Wheat* (except for wheat varieties with CLEARFIELD Technology ONLY) Barley*, Lucerne*, Chickpeas	Sorghum [▮] , Maize [▮] (except for maize varieties with CLEARFIELD Technology ONLY), Dryland Cotton [▼]	Irrigate Cotton, Sunflower	Any other crops

* The following additional requirements apply if it is intended to plant Wheat (except for varieties with CLEARFIELD Technology), Barley or Lucerne during the next winter season.

DO NOT apply Imtrade Ember 240 SL Herbicide in areas where rainfall from spraying to sowing of cereals is expected to be below 200mm.

= The following additional requirements apply if it is intended to plant Sorghum or Maize (except for varieties with CLEARFIELD Technology) during the next summer season.

DO NOT apply Imtrade Ember 240 SL Herbicide in areas where rainfall from spraying to sowing of Sorghum or Maize is expected to be below 500mm.

▼ The following additional requirements apply if it is intended to plant Dryland Cotton during the next summer season.

DO NOT apply Imtrade Ember 240 SL Herbicide where rainfall from spraying to sowing of Dryland Cotton is expected to be below 550mm.

Following use in Sugarcane and Peanuts:

MONTHS AFTER APPLICATION						
0	4	5	10	18	24	36
Mungbeans, Peanuts, Soybeans, maize varieties with CLEARFIELD Technology ONLY : - Pacific Hycorn 62IT - Pacific Hycorn 53IT - Pioneer 3395IR;	Chickpeas, Lucerne*, Barley*, Wheat* (except for wheat varieties with CLEARFIELD Technology ONLY)	Lupins, Pasture Legumes	Maize ⁼ (except for maize varieties with CLEARFIELD Technology ONLY), Dryland Cotton [▼]	Sorghum ⁼	Irrigated Cotton, Sunflower	Any other crops

* The following additional requirements apply if it is intended to plant Wheat (except for varieties with CLEARFIELD Technology), Barley or Lucerne during the next winter season.

DO NOT apply Imtrade Ember 240 SL Herbicide later than the end of December.

DO NOT apply Imtrade Ember 240 SL Herbicide in areas where rainfall from spraying to sowing of cereals is expected to be below 200mm.

= The following additional requirements apply if it is intended to plant Sorghum or Maize (except for varieties with CLEARFIELD Technology) during the next summer season.

DO NOT apply Imtrade Ember 240 SL Herbicide in areas where rainfall from spraying to sowing of Sorghum or Maize is expected to be below 500mm.

▼ The following additional requirements apply if it is intended to plant Dryland Cotton during the next summer season.

DO NOT apply Imtrade Ember 240 SL Herbicide where rainfall from spraying to sowing of Dryland Cotton is expected to be below 550mm.

RE-ENTRY PERIOD FOR IMTRADE EMBER 240 SL HERBICIDE PLUS PENDIMETHALIN

When tank-mixing Imtrade Ember 240 SL Herbicide with Pendimethalin, **DO NOT** allow entry into treated areas for 12 hours. When prior entry is necessary, wear cotton overalls button to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

CROP SAFETY

Imtrade Ember 240 SL Herbicide may, in some circumstances, lead to transient crop yellowing and temporary slowing of growth in Sugarcane but plants soon recover and yield is unaffected. This effect may be more pronounced when Imtrade Ember 240 SL Herbicide is used under conditions where soil moisture is high within 2 days following application and on light/sandy soil. Contact Imtrade Australia Pty Ltd for advice on variety susceptibility before using Imtrade Ember 240 SL Herbicide.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate dams, streams, rivers, waterways or drains with this product or used container. **DO NOT** spray within 50m of wetlands or waterways.

STORAGE AND DISPOSAL (1L & 2L only)

Store product in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. Rinse containers before disposal. Add rinsings to the spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers and product.

STORAGE AND DISPOSAL (5L, 10L & 20L)

Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or an approved waste management facility. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers and product.

Envirodrum 110L Mini Bulk Returnable Container: Store the closed, original sealed Envirodrum in a well-ventilated area, as cool as possible and locked away from children, animals, food, foodstuffs, seed and fertilisers. **DO NOT** store for prolonged periods in direct sunlight. **DO NOT** tamper with the Micro Matic valve or the security seal. **DO NOT** contaminate the Envirodrum with water or any other foreign matter. After each use of the product, please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase.

Refillable containers (1000L only)

Store in the closed, original container in a well-ventilated area, as cool as possible and locked away from children, animals, food, foodstuffs, seed and fertilisers. **DO NOT** store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and preparing spray, wear elbow-length PVC gloves. Wash hands after use. After each day's use, wash gloves. When tank mixing with Imtrade Cyclone 330 EC Herbicide or Imtrade Paraquat 250 Herbicide, consult also the safety directions for those products.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre (Phone Australia 13 11 26).

Additional GHS Hazard & Precautionary Statements

●Keep out of reach of children; ●**DO NOT** get on clothing; ●Rinse skin or shower with water; ●**IF SWALLOWED:** Rinse mouth. Do **NOT** induce vomiting; ●Store in a dry place.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from your supplier or Imtrade Australia Pty Ltd website at www.imtrade.com.au

CONDITIONS OF SALE

Imtrade Australia Pty Ltd warrants that this product conforms to the chemical description on the label. The use of Imtrade Ember 240 SL Herbicide being beyond the control of the manufacturer, Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Imtrade Australia Pty Ltd has any authority to add to or alter these conditions.

® Other Registered Trademarks

**In a Transport Emergency
Dial 000
Police or Fire Brigade**

APVMA Approval No. 64943/106765