STORAGE AND DISPOSAL *(20L & 200L only)*
Store in the closed, original container in a well-ventilated area as cool as possible, out of direct sunlight. This container can be recycled if it is clean, dry, free of visible residues and has the drum muster logo visible. Triple or pressure rinse container for disposal. Add rinsings to spray tank. Dispose of the rinsate by adding it to the spray tank. **DO NOT** dispose of undiluted chemicals on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any drum muster collection or similar container management program site. The cap should not be replaced but may be taken separately. If not recycling, break, crush or puncture container and deliver empty packaging for appropriate disposal to an approved waste management facility. Empty containers and product should not be burnt.

Envirodrum Mini Bulk Returnable Container *(110L only)*
DO NOT tamper with the Micro Matic valve or the security seal. **DO NOT** contaminate the Envirodrum with water or any other foreign matter. After each use of the product, please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase.

Refillable containers *(1000L only)*
Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS
Harmful if swallowed. Will damage the eyes. Will irritate the skin. Avoid contact with eyes and skin. When opening the container and preparing the spray, wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length PVC gloves and face shield or goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking, smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, face shield or goggles and contaminated clothing.

FIRST AID
If poisoning occurs, contact a doctor or Poisons Information (Phone Australia 13 11 26). If swallowed, **DO NOT** induce vomiting. Give a glass of water. If in eyes, wash out immediately with water.

Additional GHS Hazard & Precautionary Statements

•May cause respiratory irritation; •Harmful if inhaled; •May damage fertility or the unborn child; •Toxic to aquatic life with long lasting effects; •Avoid breathing fumes, mists, vapours or spray; •**DO NOT** eat, drink or smoke when using this product; •Use only outdoors or in a well-ventilated area; •Contaminated work clothing should not be allowed out of the workplace; •**IF ON SKIN:** Wash with plenty of soap and water; •**IF INHALED:** Remove victim to fresh air and keep at rest in a position comfortable for breathing; •If skin irritation or rash occurs: Get medical advice; •If eye irritation persists: Get medical advice; •Collect spillage; •In case of a fire, use carbon dioxide, dry chemical, foam, water fog; •Store in a dry place.

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

IMTRADE LIONEX SELECTIVE HERBICIDE

ACTIVE CONSTITUENTS: 250g/L MCPA present as the ethyl hexyl ester
25g/L DIFLUFENICAN
SOLVENTS: 325g/L LIQUID HYDROCARBONS
150g/L N-MEHTYL-2-PYRROLIDONE

GROUP F I HERBICIDE

For the control of certain broadleaf weeds in winter cereals and clover as specified in the Directions for Use table

IMPORTANT: READ THIS LABEL AND THE ATTACHED LEAFLET THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT

NET CONTENTS: 5L-1000L

Imtrade Australia Pty Ltd
ABN 13 090 151 134
17 Ocean Street, Kwinana WA 6167
Tel: 1800 171 799 Fax: 1800 171 788
Web: www.imtrade.com.au

SAFETY DATA SHEET
Additional information is listed in the Safety Data Sheet, which can be obtained from your supplier or Imtrade Australia Pty Ltd website at www.imtrade.com.au

CONDITIONS OF SALE
Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Imtrade Australia Pty aLtd has any authority to add to or alter these conditions.

UN 3082	**ENVIRONMENTALLY HAZARDOUS SUBSTANCES, LIQUID, N.O.S.**
In a Transport Emergency Dial 000 Police or Fire Brigade	
PG III	**HAZCHEM •3Z**

APVMA Approval No. 65028/106304

Batch No:
Date of Manufacture:

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY INSTRUCTIONS BEFORE OPENING OR USING

IMTRADE LIONEX SELECTIVE HERBICIDE

ACTIVE CONSTITUENTS: 250g/L MCPA present as the ethyl hexyl ester
25g/L DIFLUFENICAN
SOLVENTS: 325g/L LIQUID HYDROCARBONS
150g/L N-MEHTYL-2-PYRROLIDONE

GROUP F I HERBICIDE

For the control of certain broadleaf weeds in winter cereals and clover as specified in the DIRECTIONS FOR USE table.

IMPORTANT: READ THIS LEAFLET THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT

APVMA Approval No. 65028/106304

Imtrade Australia Pty Ltd
ABN 13 090 151 134
17 Ocean Street, Kwinana Western Australia 6167
Tel: 1800 171 799 Fax: 1800 171 788
Web: www.imtrade.com.au

DIRECTIONS FOR USE
RESTRAINTS:
DO NOT apply if crops or weeds are stressed due to dry or excessively moist conditions
DO NOT apply to crops under stress due to disease or insect damage.
DO NOT apply to frost-affected crops or if frosts are imminent.
DO NOT apply if heavy rain is expected within 4 hours.

CROP	WEEDS CONTROLLED	STAGE OF WEED GROWTH	STATE	RATE/HA	CRITICAL COMMENTS
CEREALS Wheat, Barley Oats, Triticale, Cereal Rye (including Cereals undersown with Clover) **PASTURE** Newly sown and established clover-based pasture, clover for hay and seed production	Wild Radish	Up to the 2-leaf stage and not More than 60mm in diameter	WA only	250 mL	**CROP STAGE** ***Cereals*** **Up to 750 mL** (3 leaf to fully tillered stage – Z13 to 30) **Over 750 mL** (5 leaf to fully tillered stage – Z15 to 30) Optimum results are achieved when sprayed at 3-5 leaf crop stage (generally 4-8 weeks post sowing) **WA only**: **DO NOT** Apply to Barley or Kulin Wheat before the 5 leaf stage (Z15). **Warning:** Imtrade Lionex Selective Herbicide may cause transient crop yellowing of cereals. Some varieties of oats have not been tested. (Refer to **"Crop Tolerance"** section of **General Instructions**). *continued next page ...*
		Up to the 4-leaf stage and not more than 120mm in diameter	All States	500 mL	
		Up to the 6-leaf stage and not more than 150mm in diameter		750 mL	
		Up to the 8-leaf stage and not more than 180mm in diameter		1.0L	

CROP	WEEDS CONTROLLED	STAGE OF WEED GROWTH	STATE	RATE/HA	CRITICAL COMMENTS
CEREALS Wheat, Barley Oats, Triticale, Cereal Rye (including Cereals undersown with Clover) **PASTURE** Newly sown and established clover-based pasture, clover for hay and seed production	Charlock, Hedge Mustard, Indian Hedge Mustard, Shepherd's Purse, Turnip Weed, Wild Turnip	Up to the 2-leaf stage and not more than 60mm in diameter	All States	500 mL	*continued from previous page ...* ***Clover*** Application is recommended prior to the eighth trifoliate leaf stage, however, applications prior to the third leaf stage may result in crop damage especially under stressed conditions and in sandy soils. **DO NOT** apply to Annual Medics or Lucerne. **Warning**: Imtrade Lionex Selective Herbicide may cause transient crop yellowing of clover, and may affect growth and seed set of some varieties of clover. (Refer to **Crop Tolerance –**section of the **General Instructions**).
		Up to the 4-leaf stage and not more than 120mm in diameter		750 mL	
		Up to 6-leaf stage and not more than 150mm in diameter		1.0L	
	London Rocket	Up to the 5-leaf stage and not more than 120mm in diameter	Qld only	750 mL	
	Ward's Weed		SA only		
	Capeweed	Up to the 2-leaf stage and not more than 60mm in diameter	All States	500 mL	
		Up to the 4-leaf stage and not more than 120mm in diameter		1.0L	*continued next page ...*

CROP	WEEDS CONTROLLED	STAGE OF WEED GROWTH	STATE	RATE/HA	CRITICAL COMMENTS
CEREALS Wheat, Barley Oats, Triticale, Cereal Rye (including Cereals undersown with Clover) **PASTURE** Newly sown and established clover-based pasture, clover for hay and seed production	Crassula	Up to the 2-leaf stage	All States	500 mL	*continued from previous page ...* **WEED STAGE** Apply when weeds are actively growing. In most situations the rate specified for each weed size will give satisfactory control. Under certain conditions such as: • high crop and weed density, • late season germinations, • abnormal weed growth (include early flowering), higher rates of product (up to the maximum rate of application specified for that weed) may be required.
		Up to the 4-leaf stage		750 mL	
	Prickly Lettuce	Up to the 2-leaf stage		500 mL	
		Up to the 4-leaf stage		750 mL	
		Up to the 6-leaf stage		1.0L	
	Dense-Flower Fumitory	Up to the 2-leaf stage		750 mL	
	Corn Gromwell, Saffron Thistle, Toad Rush			1.0L	
	Deadnettle	Up to the 2-leaf stage	NSW, Vic, SA only	1.0L	
	Sorrel		Vic only	1.0L	
	Canola (Rape)	Up to the 4-leaf stage	All States	500 mL	*continued next page ...*

CROP	WEEDS CONTROLLED	STAGE OF WEED GROWTH	STATE	RATE/HA	CRITICAL COMMENTS
CEREALS Wheat, Barley Oats, Triticale, Cereal Rye (including Cereals undersown with Clover) **PASTURE** Newly sown and established clover-based pasture, clover for hay and seed production	Purple Goosefoot	Up to 6-leaf stage	Qld only	500 mL	*continued from previous page ...* Imtrade Lionex Selective Herbicide will not effectively control: • regrowth of suppressed weeds, • transplanted weeds, • regrowth from rhizomes or roots, • weeds growing under stress from previous herbicide applications. **GRAZING** Efficacy on larger weeds will be improved by grazing with normal levels of stock after the 7 day withholding period. Refer to **"Protection of Livestock"** for grazing precautions. *continued next page ...*
	Turnip Weed, Wild Turnip	Cotyledon to 2-leaf stage	NSW only (West of Newel Hwy) SA only (Eyre Peninsula north of the line between Venus Bay and Cowell)	350 mL	

CROP	WEEDS CONTROLLED	STAGE OF WEED GROWTH	STATE	RATE/HA	CRITICAL COMMENTS
CEREALS Wheat, Barley, Oats, Triticale, Cereal Rye	Fumitory	2-6 leaf stage	All States	500 mL + 200 mL Terbutryn (500 g/L)	*continued from previous page ...* **APPLICATION** Activity of this product will be reduced if weeds are stressed. Optimum results will be obtained if good soil moisture exists at and after application. Where crop or weed density is high, water volume should be increased. **WILD RADISH** Imtrade Lionex Selective Herbicide will provide residual control of Wild Radish for up to 4 weeks after application. Effective residual activity of this product may be reduced where: • rates lower than 1.0L/ha are used, • dry conditions prevail, • poor coverage of the soil surface is achieved, • crop is planted in non-wetting sand, • soils contain a high content of organic matter. Optimum results will be obtained if soil moisture exists at and after application.

CROP	WEEDS CONTROLLED	STAGE OF WEED GROWTH	STATE	RATE/HA	CRITICAL COMMENTS
CEREALS Wheat, Barley, Oats, Triticale, Cereal Rye (including Cereals undersown with Clover) **PASTURE** Newly sown and established clover-based pasture, clover for hay and seed production	**SUPPRESSION OF THE FOLLOWING WEEDS**				**Refer to Critical Comments on previous page**
	Saffron Thistle	Up to the 6-leaf stage	All States	1.0L	
	Chickweed, Fireweed, Hexham Scent (King Island Melilot), Iceplant, Mouse-Eared Chickweed, Night-Scented Stock, Paterson's Curse, Peppercress, Skeleton Weed, Long Storksbill, Volunteer Lupins	Up to the 4-leaf stage			
	Wireweed (Hogweed)			750 mL	

CROP	WEEDS CONTROLLED	STAGE OF WEED GROWTH	STATE	RATE/HA	CRITICAL COMMENTS
	SUPPRESSION OF THE FOLLOWING WEEDS				**Refer to Critical Comments on previous page**
CEREALS Wheat, Barley, Oats, Triticale, Cereal Rye (including Cereals undersown with Clover) **PASTURE** Newly sown and established clover-based pasture, clover for hay and seed production	Common Sowthistle (Milk Thistle), Cowvine, Dock, Doublegee (Spiny Emex), Fat Hen, Horehound, Hyssop Loosestrife, Marshmallow Rough Poppy, Scarlet Pimpernel, Stemless Thistle, Tree Hogweed, Variegated Thistle, Vetch (Tares)	Up to the 2-leaf stage	All States	1.0L	

CROP	WEEDS CONTROLLED	STAGE OF WEED GROWTH	STATE	RATE/HA	CRITICAL COMMENTS
CEREALS Wheat, Barley, Oats, Triticale, Cereal Rye	Wild Radish	Up to the 4-leaf stage and not more than 120mm in diameter	All States	350 mL plus 200 mL MCPA LVE (500g/L)	**Refer also to all Critical Comments relating to weed stage, grazing, application and Wild Radish above.** [T]Reduced efficacy (suppression only) may be achieved on Wild Radish larger than 8-leaf or greater then 180mm in diameter. **DO NOT** use this tank-mix if cereals are undersown with Lucerne or Annual Medics. **Crop Stage** Imtrade Lionex Selective Herbicide 350 mL + MCPA LVE 200 mL: Apply from 3-leaf to fully tillered (Zadok's Z13 to Z30). Imtrade Lionex Selective Herbicide 500 mL + MCPA LVE 200 mL: Apply from 3-leaf to fully tillered (Zadok's Z13 to Z30). Imtrade Lionex Selective Herbicide 500 mL + MCPA LVE 400 mL: Apply from 5-leaf stage to fully tillered (Zadok's Z15 to Z30). Optimum results are achieved when sprayed at 3-5 leaf crop stage (generally 4-8 weeks post sowing). **WA only: DO NOT** apply to Barley or Kulin Wheat before the 5 leaf stage (Z15). **Warning:** Imtrade Lionex Selective Herbicide may cause transient crop yellowing of cereals. Some varieties of oats have not been tested. (Refer to "**Crop Tolerance**" section of **General Instructions**). Observe instructions also on MCPA LVE product label.
		Up to the 6-leaf stage and not more than 150mm in diameter		500 mL plus 200 mL MCPA LVE (500g/L)	
		Up to the 8-leaf stage and not more than 180mm in diameter[T]		500 mL plus 400 mL MCPA LVE (500g/L)	

***Pasture**:* The tolerance of clover varieties to Imtrade Lionex Selective Herbicide can vary with rate of application, soil type, crop health, stage of growth and degree of moisture and temperature stress.

Warning

Imtrade Lionex Selective Herbicide may result in transient crop yellowing and suppression of growth with a resultant initial reduction in dry matter, particularly at rates in excess of 500 mL/ha and in areas of double spray. For this reason we recommended application prior to the 8 trifoliate leaf stage. However, at the lower rates (500 mL/ha and less) and under normal growing conditions, subsequent growth and seed yield should not be affected.

Under normal growing conditions the following varieties have shown acceptable levels of foliage tolerance to Imtrade Lionex Selective Herbicide applied at 500 mL/ha.

ArrowLeaf: Zulu **Balansa:** Paradana
Berseem: Sacromonte **Persian**: Kyambro, Lupers Maral **White**: Haifa
Subterranean Clover: Daliak, Dalkeith, Denmark, Esperance, Geraldton, Goulburn, Karridale, Larissa, Leura, Mt Barker, Nungarin, Rosedale, Seaton Park, Trikkala and Woogenellup.

The effects of Imtrade Lionex Selective Herbicide on clover seed yield have been tested on the following varieties. Under normal growing conditions they show acceptable levels of tolerance to Imtrade Lionex Selective Herbicide applied at 500 mL/ha.

Subterranean Clover: Esperance, Goulburn, Larrisa, Seaton Park and Trikkala.

Warning

Rose and Strawberry Clover have shown increased sensitivity to Imtrade Lionex Selective Herbicide. Imtrade Lionex Selective Herbicide may affect the seed yield of Subterranean Clover variety Woogenellup. Some pasture grasses, including Phalaris and Cocksfoot, may show some initial reduction in vegetative growth after application of Imtrade Lionex Selective Herbicide. Care should be exercised if sensitive clover varieties or grasses are included in the pasture sward. Varieties not listed should be tested before using Imtrade Lionex Selective Herbicide over large areas.

Subsequent Crops

To reduce effect on subsequent susceptible crops (e.g. Canola), ensure thorough cultivation of soil prior to the sowing of these crops.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS
CROP HARVEST: NOT REQUIRED WHEN USED AS DIRECTED
All crops: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION

GENERAL INSTRUCTIONS

RESISTANT WEEDS WARNING

GROUP F I HERBICIDE

Imtrade Lionex Selective Herbicide is a member of the phenoxy and nicotinanilide groups of herbicides and acts by inhibiting carotenoid biosynthesis and disrupting plant cell growth. For weed resistance management Imtrade Lionex Selective Herbicide is both a Group **F** and a Group **I** Herbicide. Some naturally occurring weed biotypes resistant to Imtrade Lionex Selective Herbicide and other Group **F** and **I** herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Imtrade Lionex Selective Herbicide or other Group F or Group I herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Imtrade Australia Pty Ltd accepts no liability for any losses that may result from the failure of Lionex to control resistant weeds.

TOLERENCE

Some pre-emergence herbicides, such as atrazine, can cause stress to certain crops resulting in an increase in crop damage when using this product. Subterranean Clover is particularly sensitive.
***Cereals**:* After application some transient crop yellowing may occur. This usually appears as yellow or white banding on leaves. Provided the crop is not under stress from pre-emergent herbicide, root disease, insect damage, frost, dry or excessively moist conditions, the development of the crop and subsequent growth will be unaffected.

Warning (Oats)
The tolerance of oat varieties Esk and Nile (the two main varieties grown in Tasmania) to Imtrade Lionex Selective Herbicide has not been tested. Test a small area of crop before using Imtrade Lionex Selective Herbicide over large areas.

MIXING

To ensure even mixing, half fill the spray tank with clean water and add the required amount of product. Agitate thoroughly then add the remainder of the water. Agitate again before spraying commences. Reseal part used product container immediately after use. Spray mixtures containing Imtrade Lionex Selective Herbicide should not be left to stand overnight. Prolonged periods of exposure to cold temperatures could result in setting out of the product in the mixture.

Warning

The rubber components present in some spraying units may be affected by exposure to the solvents in Imtrade Lionex Selective Herbicide and some other agricultural products. To reduce this risk it is recommended that the spray unit be thoroughly washed with a boom cleaner and fresh water after use. Contact the spray unit manufacturer to determine the suitability of the rubber components for use with agricultural products.

APPLICATION

Boom Sprayer: A minimum of 50L of water per hectare should be used, however, for optimum results water rates of 70-100L/ha are recommended. Increase the water volume if weed infestation is heavy or crop cover is dense. Complete coverage of weeds is essential.

Aircraft (NSW, Vic, SA only): Apply in a minimum 30L water per hectare. Effective weed control will only be achieved where good coverage of leaf surface is achieved.

COMPATIBILITY

The following products are physically compatible with Imtrade Lionex Selective Herbicide as a two-way mixture in the spray tank but should only be used for the crops specified:

Crop	Imtrade Lionex Selective Herbicide	Compatible Product
Wheat, Triticale and Cereal Rye only	Up to 750mL/ha	Diclofop-methyl (also Barley) Puma® Progress (Wild Oats only), Tristar® Advance (1.5L rate only), ChemAg Maddog Selective Herbicide (Wild Oats only)
Cereals (Including undersown)	All rates	Chlorpyrifos (500g/L), Dimethoate, Thiodan®
Cereals (not undersown)	Up to 500 mL/ha	Metsulfuron, Chlorsulfuron, MCPA LV E, Imtrade Mandate 750 WG Selective Herbicide
	All rates	Bromoxynil 200g/L, 2,4-D Amine 500 Herbicide, Imtrade Rally 750 SG Herbicide, Tordon® 50-D, Cadence (up to 115g only), Eclipse®
Wheat, Barley, Triticale, and Cereal Rye only (not undersown)		ChemAg Attain 400 WG Herbicide
Wheat only (not undersown)		Imtrade Subject 240 EC Selective Herbicide
Clover	Up to 750mL/ha	Targa®, Fusilade®
Subterranean Clover		Simazine (500g/L), Simazine (500g/L) + Paraquat (200g/L) mixture
	Up to 1.0L/ha	2,4-D Amine (500g/L)

When mixing with other hebicides, crop yellowing may be enhanced. When mixing with Hoegrass®, ChemAg Maddog Selective Herbicide or Tristar® Advance some reduction in the efficacy and speed of action of these products may occur. If the crop is stressed, the application of the herbicide tank-mixtures may cause yield reduction. When mixing with Cadence® a temporary wilting may be evident in some crops after application. Growers should seek advice before spraying recently released cereal varieties.

Use the recommended rates for both herbicides in the tank-mixture as well as the surfactant recommendation of the grass herbicide. If another herbicide is applied as a tank mix, observe the plantback restrictions on that label. **DO NOT** add surfactant when mixing Imtrade Lionex Selective Herbicide and Ally® (Metsulfuron-methyl).

Simazine: Refer to the Simazine label for correct application rates, especially with regard to soil types.

This product may be mixed in the spray tank with one of the following insecticides according to the directions for use on this product: Alpha-cypermethrin 100g/L, Deltamethrin 27.5g/L, Lambda-cyhalothrin 250g/L and Talstar®.

Warning

DO NOT use crop oils with Imtrade Lionex Selective Herbicide or Imtrade Lionex Selective Herbicide tank mixtures with other products in cereals. As formulations of other manufacturer's products are beyond the control of Imtrade Australia Pty Ltd, all mixtures should be tested prior to mixing commercial quantities.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment that may cause drift onto nearby susceptible plants/crops, cropping lands or pastures. Avoid spray drift and vapour movement onto susceptible crops such as cotton, tobacco, tomatoes, vines, lupins, fruit trees and ornamentals.

PROTECTION OF LIVESTOCK

Grazing Precaution

Sprayed weeds may become more palatable to stock and a higher intake of some weeds may result in stock poisoning and death from causes such as nitrate poisoning. Care should be taken especially where Capeweed, Patterson's Curse and Variegated Thistles predominate in the pasture. Avoid grazing with young or breeding stock. **DO NOT** graze horses or pigs on Paterson's Curse. If in doubt, contact your nearest Department of Agriculture.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish.

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a well-ventilated area as cool as possible, out of direct sunlight. This container can be recycled if it is clean, dry, free of visible residues and has the drum muster logo visible. Triple or pressure rinse container for disposal. Add rinsings to spray tank. Dispose of the rinsate by adding it to the spray tank. **DO NOT** dispose of undiluted chemicals on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any drum muster collection or similar container management program site. The cap should not be replaced but may be taken separately. If not recycling, break, crush or puncture container and deliver empty packaging for appropriate disposal to an approved waste management facility. Empty containers and product should not be burnt.

Envirodrum 110L Mini Bulk Returnable Container

DO NOT tamper with the Micro Matic valve or the security seal. **DO NOT** contaminate the Envirodrum with water or any other foreign matter. After each use of the product, please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase.

Refillable containers (1000L only)

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if swallowed. Will damage the eyes. Will irritate the skin. Avoid contact with eyes and skin. When opening the container and preparing the spray, wear cotton overalls buttoned to the neck and wrist, washable hat, elbow- length PVC gloves and face shield or goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking, smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information (Phone Australia 13 11 26). If swallowed, **DO NOT** induce vomiting. Give a glass of water. If in eyes, wash out immediately with water.

Additional GHS Hazard & Precautionary Statements

•May cause respiratory irritation; •Harmful if inhaled; •May damage fertility or the unborn child; •Toxic to aquatic life with long lasting effects; •Avoid breathing fumes, mists, vapours or spray; •**DO NOT** eat, drink or smoke when using this product; •Use only outdoors or in a well-ventilated area; •Contaminated work clothing should not be allowed out of the workplace; •**IF ON SKIN:** Wash with plenty of soap and water; •**IF INHALED:** Remove victim to fresh air and keep at rest in a position comfortable for breathing; •If skin irritation or rash occurs: Get medical advice; •If eye irritation persists: Get medical advice; •Collect spillage; •In case of a fire, use carbon dioxide, dry chemical, foam, water fog; •Store in a dry place.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from your supplier or Imtrade Australia Pty Ltd website at www.imtrade.com.au

CONDITIONS OF SALE

Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Imtrade Australia Pty aLtd has any authority to add to or alter these conditions.

UN 3082	**ENVIRONMENTALLY HAZARDOUS SUBSTANCES, LIQUID, N.O.S.**
IN A TRANSPORT EMERGENCY **DIAL 000** **POLICE OR FIRE BRIGADE**	
PG III	**HAZCHEM •3Z**

APVMA Approval No. 65028/106304