

#### STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Shake the bag contents into spray tank until the bag is empty. **DO NOT** dispose of undiluted chemicals on-site. Puncture or shred and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty bags and product should not be burnt.

#### SAFETY DIRECTIONS

May irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. **DO NOT** inhale dust. When opening the container and preparing spray, wear cotton overalls over normal clothing, buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves, goggles and disposable dust mask covering mouth and nose. When using the prepared spray, wear cotton overalls over normal clothing, buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

#### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 131126). Avoid giving alcohol.

#### Additional GHS Hazard & Precautionary Statements

•Suspected of damaging fertility or the unborn child; •Very toxic to aquatic life; •**DO NOT** get on clothing; •Contaminated clothing should not be allowed out of the workplace; •**IF SWALLOWED:** Rinse mouth. **DO NOT** induce vomiting; •**IF ON SKIN:** Wash with plenty of soap and water; •If eye irritation persists: Get medical advice; •If exposed or concerned: Get medical advice; •If skin irritation or rash occurs: Get medical advice; •In case of fire, note the following: Water fog or fine spray is the preferred medium for large fires. Try to contain spills, minimise spillage entering drains or water courses; •Store locked up.

#### SAFETY DATA SHEET

Additional information is listed in the Data Sheet, which is available from the supplier or Imtrade Australia Pty Ltd website at [www.imtrade.com.au](http://www.imtrade.com.au)

#### CONDITIONS OF SALE

Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill of judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Imtrade has any authority to add to or alter these conditions.

## CAUTION

**KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**



# IMTRADE MANCOZEB 750 DF FUNGICIDE

ACTIVE CONSTITUENT: 750 g/kg MANCOZEB

GROUP **M3** FUNGICIDE

**For the control of certain fungal diseases of fruit, field crops, tobacco, turf, vegetables and ornamentals as per the Directions for Use Table**

**IMPORTANT: READ THIS LABEL AND THE ATTACHED LEAFLET THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT**

**NET CONTENTS:  
25kg**

Imtrade Australia Pty Ltd  
ABN 13 090 151 134  
17 Ocean Street, Kwinana WA 6167  
Tel: 1800 171 799 Fax: 1800 171 788  
Web: [www.imtrade.com.au](http://www.imtrade.com.au)

# UN 3077

ENVIRONMENTALLY  
HAZARDOUS SUBSTANCE,  
SOLID, N.O.S. MANCOZEB

In a Transport Emergency  
Dial 000  
Police or Fire Brigade



APVMA Approval No:  
63513/25/0909

Batch No:  
Date of Manufacture:

# CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



# IMTRADE MANCOZEB 750 DF FUNGICIDE

ACTIVE CONSTITUENT: 750 g/Kg MANCOZEB

**GROUP M3 FUNGICIDE**

For the control of certain fungal diseases of fruit,  
field crops, tobacco, turf, vegetables and ornamentals  
as per the Directions for Use Table

**IMPORTANT: READ LEAFLET THOROUGHLY  
BEFORE OPENING OR USING THIS PRODUCT**

**APVMA Approval No: 63513/0909**

Imtrade Australia Pty Ltd  
ACN 13 090 151 134  
17 Ocean Street, Kwinana, Western Australia 6167  
Tel: 1800 171 799 Fax: 1800 171 788  
Web: [www.imtrade.com.au](http://www.imtrade.com.au)

**DIRECTIONS FOR USE**

For use in All States where appropriate for crop and/or disease.

**Restrains:** DO NOT incorporate this product with an Integrated Mite Control programme.**TREE AND VINE CROPS**

RATE In the following table, all rates are given for dilute spraying. For concentrate spraying, refer to the Application section.				Critical Comments. For all uses in this table: Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods.
CROP	DISEASE/PEST	RATE	WHP Days Harvest (H) Grazing (G)	CRITICAL COMMENTS
Citrus	Black Spot	200g/100L plus spray oil (use according to its label instructions)	Not required	<b>On Heavy Soil:</b> Spray at 8 and 15 weeks following a Bordeaux spray (5-5-100) at petal fall. <b>On Light Soil:</b> Add an adjuvant (spreader/sticker) according to label recommendations per 100L. Spray at 6 and 13 weeks following a Bordeaux spray (3.5-3.5-100) at petal fall. <b>Qld, NT:</b> Spray at 6 and 12 weeks after a copper spray at ½ to ¾ petal fall. Apply at first sign of mite activity on fruit, usually between December and May. Choose a spray oil recommended for use on citrus.
	Brown Citrus Mite, Citrus Rust Mite			
	Citrus Bud Mite			Apply when blossom malformation is observed, usually February. Choose a spray oil recommended for use on citrus.
Custard Apples	Pseudocercospora Fruit Spot	200g/100L	1 day (H)	<b>DO NOT</b> apply during flowering. Apply at 3-4 week intervals from first fruit set until harvest. Apply at 2 week intervals in wet, cloudy weather.
Grapevines	Black Spot, Downy Mildew	200g/100L	30 days (H)	For Blackspot control apply every 10-14 days starting at bud burst. If Downy Mildew is expected, continue spray programme at intervals of 10-14 days until the threat of disease has passed. Use the shorter spray interval during weather conditions favourable to disease development.
	Phomopsis Cane and Leaf Spot	150 to 200g/100L		Apply at budburst and then repeat 7 to 10 days later.

TREE AND VINE CROPS continued

CROP	DISEASE/PEST	RATE	WHP Days Harvest (H) Grazing (G)	CRITICAL COMMENTS
Mangoes	Anthraxnose ( <i>Collectotrichum gloeosporioides</i> )	200g/100L	1 day (H)	Spray at weekly intervals during flowering then monthly until near harvest.
Passionfruit	Brown Spot ( <i>Alternaria passiflorae</i> ), Septoria Leaf Spot	150g/100L	1 day (H)	Apply at 10-14 day intervals from October to May and every 21-28 days for the remainder of the year.
	Anthraxnose (NSW only)	200g/100L + 600 mL/100L miscible summer oil		Spray at weekly intervals during flowering and then fortnightly until near harvest.
Pawpaws (Papaya)	Black Spot	200g/100L	1 day (H)	Apply with a wetting agent at 10 to 14 day intervals during periods conducive to disease.
Pome Fruit Apples & Pears	Apple Scab (Black Spot), Bitter Rot, Target Spot, Fly Speck, Sooty Blotch	150g to 200g/100L	14 days (H)	Apply at 7-14 day intervals following a cover copper spray at green tip. Use higher rate or shorter interval between sprays where conditions favour disease. Imtrade Mancozeb 750 DF Fungicide can be mixed with Systhane 400* WP Fungicide for the control of Apple Scab (Black Spot). Read and follow all the recommendations on the label of the tank mix partner. Imtrade Mancozeb 750 DF Fungicide may be harmful to <i>Typhlodromus pyri</i> and its use in orchards where integrated control is practiced should be avoided.
Stone Fruit Almonds Apricots Cherries Nectarines Peaches Plums <sup>^</sup>	Brown Rot, Rust, Shot Hole, Freckle	150g to 200g/100L	14 days (H)	Apply at early bloom (1-10%), mid to full bloom (50-100%), petal fall and shuck fall, then at intervals of 14 days.  Continue with a protective spray programme at 2 week intervals.  <sup>^</sup> May be phytotoxic to Wilson, Wickson and Bellarose varieties of plums. It is advisable to test any new plum varieties on a small scale before applying to the whole crop.

## NON-TREE AND VINE CROPS

CROP	DISEASE	RATE (per ha)	RATE (per 100L)	WHP Days Harvest (H) Grazing (G)	CRITICAL COMMENTS
FRUIT Bananas	Leaf Spot	2.2 to 4.5 kg/ha plus spray oil (used in accordance to its label directions)	200g/100L plus spray oil (used in accordance to its label directions)	NIL (H)	Apply when weather conditions favour disease outbreak. Repeat treatments in accordance with locally recommended spray schedules. Use sufficient water to ensure good coverage. <b>NSW: Ground application</b> – Apply at 3 week intervals from December or January until May. <b>Aerial application</b> – Apply at 7 to 10 day intervals from December or January until May. <b>South Qld:</b> Apply at 3 week intervals from December or January until May. <b>North Qld:</b> Apply every 2 weeks during the wet season. Extend to 3 weeks after this and then every 4 to 5 weeks during the drier months returning to 3 weeks prior to the next wet season.
	Leaf Speckle, Cordana Leaf Spot, Fruit Speckle, Black Pit	2.2kg /ha	200g/100L		

## VEGETABLES Peas and Beans grown for fresh consumption

Broadbeans/ Faba Beans	Rust, Cercospora Leaf Spot	1.7-2.2 kg/ha	–	7 days (H) 7 days (G)	Apply when disease symptoms first appear and then repeat at 7-10 day intervals. Use the higher rate and shorter spray interval during weather conditions favourable to disease development. Spray when leaves, flowers or pods show first signs of infection and weather conditions are likely to remain humid or wet and favour disease development.
	Chocolate Spot, Ascochyta Leaf Blight (suppression only)	1.7-2.5 kg/ha			

**NON-TREE AND VINE CROPS** continued

CROP	DISEASE	RATE (per ha)	RATE (per 100L)	WHP Days Harvest (H) Grazing (G)	CRITICAL COMMENTS
Peas ( <i>Pisum sativum</i> )	Rust	1.7-2.2 kg/ha	–	7 days (H) 7 days (G)	Apply when disease symptoms first appear and then repeat at 7-10 day intervals. Use the higher rate and shorter spray interval during weather conditions favourable to disease development. Spray when leaves, flowers or pods show first signs of infection and weather conditions are likely to remain humid or wet and favour disease development.
	Ascochyta Leaf Blight (suppression only)	1.7-2.5 kg/ha			If disease pressure is severe and weather conditions favour the spread of disease, repeat the spray treatment at 2-3 week intervals in order to protect new growth and developing pods. Use the higher rate for dense crops and if disease is severe. <b>Volume of applications:</b> for ground apply 100L/ha; for aircraft apply at least 30L/ha.
Green Beans ( <i>Phaseolus</i> spp.)	Angular Leaf Spot, Anthracnose Rust	1.7-2.2 kg/ha			Apply when disease symptoms first appear and then repeat at 7-10 day intervals. Use the higher rate and shorter spray interval during weather conditions favourable to disease development. Spray when leaves, flowers or pods show first signs of infection and weather conditions are likely to remain humid or wet and favour disease development.

**NON-TREE AND VINE CROPS** continued

CROP	DISEASE	RATE (per ha)	RATE (per 100L)	WHP Days Harvest (H) Grazing (G)	CRITICAL COMMENTS
<b>VEGETABLES</b>					
Beetroot Silver Beet Spinach	Cercospora Leaf Spot, Downy Mildew	1.7-2.2 kg/ha	150-200g/1 00L	14 days (H)	Spray at first sign of disease, then at 7-10 day intervals. Use the higher rate and shorter spray interval during weather conditions that are favourable to disease development.
Capsicum	Target Spot	2-3 kg/ha			
Carrots	Cercospora Leaf Spot, Alternaria Leaf Spot	1.7-2.2 kg/ha	7 days (H)		
Celery	Septoria Leaf Spot				
<b>Cole Crops</b> Broccoli Brussels Sprout Cabbage Cauliflower	Alternaria Leaf Spot, Anthracnose, Downy Mildew, Ring Spot				
<b>Curcubits</b> Cantalope Cucumber Melons Pumpkin Squash Zucchini	Downy Mildew, Anthracnose, Gummy Stem Blight, Septoria Spot				

**NON-TREE AND VINE CROPS** continued

CROP	DISEASE	RATE (per ha)	RATE (per 100L)	WHP Days Harvest (H) Grazing (G)	CRITICAL COMMENTS
Fennel (Tas only)	Leaf Blight ( <i>Cercosporidium</i> )	2.2 kg/ha	–	14 days (H)	Apply as a ground spray if disease symptoms are present in November. Allow for one repeat treatment 10-14 days later. Aerial application may be required in prolonged wet periods.
Lettuce	Downy Mildew, Anthracnose, Septoria Leaf Spot	1.7-2.2 kg/ha	150-200g/1 00L		Spray at first sign of disease, then at 7-10 day intervals. Use the higher rate and shorter spray interval during weather conditions that are favourable to disease development.
Onions Garlic	Downy Mildew, Purple Blotch	2.2-3.5 kg/ha	150-200g/1 00L	7 days (H)	Apply when disease symptoms first appear and then repeat at 7-10 day intervals. Use higher rate and shorter intervals during weather conditions that are favourable to disease development.
Potatoes	Early and Late Blight	1.7 to 2.2 kg/ha	150-200g/1 00L	Not required	
Rhubarb				Downy Mildew, Rust	
Tomatoes	Early Blight (Target Spot), Late Blight, Irish Blight, Leaf Mould, Anthracnose, Phoma Rot, Grey Leaf Spot	2-3 kg/ha	150-200g/1 00L	7 days (H)	



**NON-TREE AND VINE CROPS** continued

CROP	DISEASE	RATE (per ha)	RATE (per 100L)	WHP Days Harvest (H) Grazing (G)	CRITICAL COMMENTS
<b>ORNAMENTALS</b>					
Carnations	Rust, Alternaria Leaf Spot	150-200g/100L	-	-	Apply when disease symptoms first appear and then repeat at 7-10 day intervals. Use higher rate and shorter intervals during weather conditions that are favourable to disease development.  <b>Caution:</b> Imtrade Mancozeb 750 DF Fungicide may be phytotoxic to some ornamental species, in particular some varieties of ferns. It is recommended to test a few plants before widespread use.
Chrysanthemum, Gladioli	Botrytis, Septoria Leaf Spot				
Roses	Black Spot				
<b>FIELD CROPS</b>					
Cotton Pima var only	Alternaria Leaf Spot	2.5 kg/ha	-	14 days (H)	Imtrade Mancozeb 750 DF Fungicide is a protectant fungicide. To maximize control ensure thorough coverage of all plant surfaces. Begin applications as soon as disease symptoms appear. Repeat before each infection period (e.g. dew or rain) but no later than 7 to 10 days after the last application. <b>DO NOT</b> apply more than 4 sprays per season.
Peanuts	Cercospora Leaf Spot, Peanut Rust	1.7-2.2 kg/ha	-	14 days (H) 14 days (G)	Apply at intervals of 10-14 days, starting when the disease symptoms first appear. If rust is anticipated, apply at 7 to 10 day intervals. Use the higher rate and the shorter spray interval during weather conditions favourable to disease development.
Poppies	Downy Mildew	2.5 kg/ha	-	7 weeks (H)	Apply as a protectant spray when weather conditions favour disease development. Repeat spray at 14 day intervals if conditions persist.

**NON-TREE AND VINE CROPS** continued

CROP	DISEASE	RATE (per ha)	RATE (per 100L)	WHP Days Harvest (H) Grazing (G)	CRITICAL COMMENTS
<b>PULSES grown for dry beans:</b>					
Broadbeans/ Faba Beans	Ascochyta Blight, Chocolate Spot, Rust, Cercospora	1-2.2 kg/ha	-	4 weeks (H) 14 days (G)	<p>Spray when leaves, flowers or pods on plants first show signs of infection and weather conditions are likely to remain humid or wet and favour disease development.</p> <p>If disease pressure is severe and conditions favour the spread of disease, repeat sprays at 2-3 weeks to protect new growth and developing pods. Use the higher rate for dense crops and if disease is severe. Repeat more frequently if severe disease persists. For improved coverage add a registered surfactant at 10-20 mL/100L of spray.</p>
Chickpeas	Ascochyta Blight, Botrytis Grey Mould				
Field Peas	Ascochyta Blight, Black Spot, Botrytis Grey Mould, Rust				
Lentils and Vetch	Ascochyta Blight, Botrytis Grey Mould, Rust				
Lupins	Botrytis Grey Mould, Anthracnose				
Beans (Phaseolus spp.)	Ascochyta Blight, Angular Leaf Spot, Anthracnose, Rust				

**NON-TREE AND VINE CROPS** continued

CROP	DISEASE	RATE (per ha)	RATE (per 100L)	WHP Days Harvest (H) Grazing (G)	CRITICAL COMMENTS
<b>OTHER CROPS</b>					
Tobacco (Field)	Blue Mould, Brown Spot, Frog Eye Leaf Spot	2.2 kg/ha	200g/100L	Not required	<b>Field treatment:</b> Apply at 7 day intervals commencing 10 days after transplanting and ceasing 7 to 10 days before the first harvest. Increase spray volume accordingly as plant size increases.  <b>Seed bed treatments:</b> Apply when seedlings are 1.5cm across and then repeat twice weekly. Wet the seedlings evenly without run-off.
Turf	Helminthosporium, Fusarium, Rhizoctonia	200-250g/100m <sup>2</sup>		–	Apply as a protective spray schedule at 7 day intervals throughout the susceptible season. Use sufficient water to ensure adequate coverage. <b>DO NOT</b> feed grass clippings from treated areas to poultry or animals.

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIOD**

**HARVEST**

**BANANAS, CITRUS, POTATOES & TOBACCO: NOT REQUIRED WHEN USED AS DIRECTED**

**MANGOES, CUSTARD APPLES,**

**PAWPAWS (PAPAYA) & PASSIONFRUIT:**

**DO NOT APPLY LATER THAN 1 DAY BEFORE HARVEST**

**BEANS, CARROTS, CELERY, COLE**

**CROPS, CUCURBITS, GARLIC, ONIONS,**

**PEAS, TOMATOES:**

**DO NOT APPLY LATER THAN 7 DAYS BEFORE HARVEST**

**WITHHOLDING PERIOD**

**HARVEST** *continued*

ALMONDS, BEETROOT, COTTON,  
CAPSICUMS, FENNEL, LETTUCE,  
PEANUTS, POME FRUIT, RHUBARB,  
SILVERBEET, SPINACH,  
STONE FRUIT:

**DO NOT APPLY LATER THAN 14 DAYS BEFORE HARVEST**

GRAPEVINES:

**DO NOT APPLY LATER THAN 30 DAYS BEFORE HARVEST**

POPPIES:

**DO NOT APPLY LATER THAN 7 WEEKS BEFORE HARVEST**

GREEN LEGUMES, PULSE CROPS:

GREEN CROPS(fresh consumption):

**DO NOT APPLY LATER THAN 7 DAYS BEFORE HARVEST**

PULSE CROPS (dry beans):

**DO NOT APPLY LATER THAN 4 WEEKS BEFORE HARVEST**

**WITHHOLDING PERIOD:**

**GRAZING**

COTTON:

**DO NOT ALLOW LIVESTOCK TO GRAZE COTTON CROP, STUBBLE OR GIN TRASH  
THAT HAS BEEN TREATED WITH IMTRADE MANCOZEB 750 DF FUNGICIDE**

GREEN LEGUMES, PEANUTS,  
PULSE CROPS (dry beans):

**DO NOT GRAZE OR CUT FOR STOCKFOOD FOR 14 DAYS**

AFTER THE LAST APPLICATION

PEAS and BEANS

(fresh consumption):

**DO NOT GRAZE OR CUT FOR STOCKFOOD FOR 7 DAYS AFTER THE LAST APPLICATION**

PAWPAWS (PAPAYA):

**DO NOT USE OR SUPPLY TREATED PAPAYA INCLUDING LEAVES FOR STOCKFOOD**

## GENERAL INSTRUCTIONS

Imtrade Mancozeb 750 DF Fungicide is a protectant Fungicide for the control of certain diseases in many crops. For best results, apply as a regular spray programme. Thorough coverage of the treated crop is essential.

## APPLICATION

### Dilute spraying:

- Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to point of run-off. Avoid excessive run-off.
- The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice.
- Add the required amount of product specified in the Directions for Use for each 100L of water. Spray to the point of run-off.
- The required dilute spray volume will change as sprayer set up and operation may also need to be changed, as the crop grows.

### Concentrate spraying:

- Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume.
- Determine the appropriate dilute spray (see Dilute Spraying above) for crop canopy. This is needed to calculate the concentrate mixing rate.
- The mixing rate for concentrate spraying can then be calculated in the following way:

#### EXAMPLE ONLY:

- (i) Dilute spray volume as determined above: for example 1500L/ha
  - (ii) Your chosen concentrate spray volume: for example 500L/ha
  - (iii) The concentration factor in this example is:  $3 \times$  (ie  $1500L \div 500L = 3$ )
  - (iv) If the dilute label rate is 10mL/100L, then the concentrate rate becomes  $3 \times 10$ , that is 30mL/100L of concentrate spray.
- The chosen spray volume, amount of product per 100L of water, and the sprayer set up and operation may need to be changed as the crop grows.
  - For further information on concentrate spraying, users are advised to consult the relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

For fruit trees and some row crops apply as a high volume dilute spray.

The spray volume will vary according to the type of crop to be treated and should be increased in heavy growth and on large plants to ensure thorough coverage. The spray volume should also be increased as the size of treated crops increases throughout the growing season. As a guide for mature crops:

#### **Crops**

Bananas, Tobacco, Vines and other row crops  
Pome and Stone Fruit  
Citrus

#### **Spray volume**

200 to 1100L/ha  
1100 to 4000L/ha  
4000 to 8000L/ha

#### **Aerial Application:**

May be applied by agricultural aircraft.

Use at least 30 to 50L/ha of spray mixture per hectare (refer to Directions for Use table).

#### **Spray Timing:**

Treatments should begin prior to disease infection and continue until the threat of disease passes. Repeated applications may be required to protect new growth. Reduce the spray interval when weather conditions favour disease development.

#### **FUNGICIDE RESISTANCE WARNING**

##### **GROUP M3 FUNGICIDE**

Multi-site activity group of fungicides. For fungicide resistance management Imtrade Mancozeb 750 DF Fungicide is a Group **M3** Fungicide. Some naturally occurring individual fungi resistant to Imtrade Mancozeb 750 DF Fungicide and other Group **M3** Fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. Imtrade Mancozeb 750 DF Fungicide and other Group **M3** Fungicides will not control these resistant fungi, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, Imtrade Australia Pty Ltd accepts no liability for any losses that may result from the failure of Imtrade Mancozeb 750 DF Fungicide to control resistant fungi.

#### **MIXING**

Slowly add the required amount of this product to the spray tank as it is being filled or thoroughly premix in a nurse tank for concentrate and aerial spraying. Add other products after Imtrade Mancozeb 750 DF is in suspension. Continue agitation while spraying to prevent product settling out and to ensure a uniform spray mixture. Rinse out spray tank, pumps, nozzles at the end of the day. When preparing spray solutions for use in a hand sprayer, premix as a slurry in a small container, and then add to sprayer containing one-third to half the desired final water volume.

#### **COMPATIBILITY**

This product may be harmful to *Typodromus pyri* and its use in orchards where integrated pest control is practiced should be avoided.

Imtrade Mancozeb 750 DF Fungicide is compatible with most commonly used insecticides and miticides such as Azinphos, Carbaryl, Dicofol, Endosulfan, Pirimicarb, Propargite, Sulphur Power, Wetting Agent or Spreader-Sticker. Also compatible with D-C-Tron.

The user is advised to check the label recommendation for the companion product before preparing tank mixes.

**Surfactants** - The addition of surfactants will improve initial spray deposits, fungicide redistribution and weatherability. Refer to manufacturers label instructions for further details.

### **PROTECTION OF LIVESTOCK**

**DO NOT** graze treated turf/grass; or feed turf/grass clippings from any treated area to poultry or livestock.

### **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

**Toxic to fish.**

Drift and run-off from treated areas may be hazardous to aquatic organisms in neighbouring areas.

**DO NOT** contaminate streams, rivers or waterways with the chemical or used containers.

**DO NOT** re-use container.

### **STORAGE AND DISPOSAL**

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Shake the bag contents into spray tank until the bag is empty. **DO NOT** dispose of undiluted chemicals on-site. Puncture or shred and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty bags and product should not be burnt.

### **SAFETY DIRECTIONS**

May irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. **DO NOT** inhale dust. When opening the container and preparing spray, wear cotton overalls over normal clothing, buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves, goggles and disposable dust mask covering mouth and nose. When using the prepared spray, wear cotton overalls over normal clothing, buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

### **FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 131126). Avoid giving alcohol.

#### **Additional GHS Hazard & Precautionary Statements**

- Suspected of damaging fertility or the unborn child;
- Very toxic to aquatic life;
- DO NOT** get on clothing;
- Contaminated clothing should not be allowed out of the workplace;
- IF SWALLOWED:** Rinse mouth. **DO NOT** induce vomiting;
- IF ON SKIN:** Wash with plenty of soap and water;
- If eye irritation persists: Get medical advice;
- If exposed or concerned: Get medical advice;
- If skin irritation or rash occurs: Get medical advice;
- In case of fire, note the following: Water fog or fine spray is the preferred medium for large fires. Try to contain spills, minimise spillage entering drains or water courses;
- Store locked up.

## SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which is available from the supplier or Imtrade Australia Pty Ltd website at [www.imtrade.com.au](http://www.imtrade.com.au)


## CONDITIONS OF SALE

Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill of judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Imtrade has any authority to add to or alter these conditions.

# UN 3077

ENVIRONMENTALLY  
HAZARDOUS SUBSTANCE,  
SOLID, N.O.S. MANCOZEB

IN A TRANSPORT EMERGENCY  
**DIAL 000**  
POLICE OR FIRE BRIGADE



APVMA Approval Number: 63513/0909