

#### STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. Keep from contact with fertilizers, insecticides, fungicides and seeds. Triple or preferably pressure rinse container before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Empty containers and product should not be burnt.

#### SAFETY DIRECTIONS

Harmful if inhaled or swallowed. Will irritate the eyes. Avoid contact with eyes and skin. **DO NOT** inhale dust or spray mist. Repeated minor exposure may have a cumulative poisoning effect. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow length PVC gloves and face shield or goggles. Wash hands after use.

#### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. *Phone Australia 13 11 26.*

Refer to attached Leaflet for additional GHS  
Hazard & Precautionary Statements

#### SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from your supplier or Imtrade Australia Pty Ltd website at [www.imtrade.com.au](http://www.imtrade.com.au)

#### CONDITIONS OF SALE

Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Imtrade Australia Pty Ltd has any authority to add to or alter these conditions.

## POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

IMTRADE

# METRIBUZIN 750 WG

HERBICIDE

ACTIVE CONSTITUENT: 750 g/kg METRIBUZIN

GROUP **5** HERBICIDE

For selective weed control in cereals, pastures and other crops, including vegetables  
as indicated in the Directions for Use table.

**IMPORTANT:** Read the attached Leaflet before use.

GROUP  
**5**

FORM  
**WG**

CONTENTS  
**1kg-50kg**



Batch No.  
DOM:

**IMTRADE**  
CropScience

17 Ocean St, Kwinana Beach WA 6167  
08 9419 0333 | [www.imtrade.com.au](http://www.imtrade.com.au)

UN 3077	ENVIRONMENTALLY HAZARDOUS SUBSTANCE, SOLID, N.O.S. (contains METRIBUZIN)
	In a Transport Emergency Dial 000 Police or Fire Brigade
PG III	HAZCHEM 2Z

APVMA Approval No: 63658/60417



# **POISON**

**KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**



# **IMTRADE METRIBUZIN 750 WG HERBICIDE**

**ACTIVE CONSTITUENT: 750 g/kg METRIBUZIN**

<b>GROUP</b>	<b>5</b>	<b>HERBICIDE</b>
--------------	----------	------------------

For selective weed control in cereals, pastures and other crops, including vegetables as indicated in the Directions for Use table

**IMPORTANT: READ THIS LEAFLET THOROUGHLY  
BEFORE OPENING OR USING THIS PRODUCT**

**APVMA Approval No: 63658/60417**

Imtrade Australia Pty Ltd  
ABN 13 090 151 134  
17 Ocean Street, Kwinana WA 6167  
Tel: 1800 171 799 Fax: 1800 171 788  
Web: [www.imtrade.com.au](http://www.imtrade.com.au)

## RESTRAINTS

**DO NOT** apply to crops that are stressed by any cause (such as severe weather conditions, water-logging, drought, excessive soil acidity or alkalinity, poor nutrient status, disease, nematode or insect damage) as crop injury may result.

## DIRECTIONS FOR USE

Crop/ Situation	Weeds Controlled	State	Rate/ha	Critical Comments
Barley	Capeweed, Catsear, Charlock, Chickweed, Common Cotula, Corn Gromwell (Sheepweed, White Ironweed), Corn Spurry, Creeping Speedwell, Crown Beard (Dogweed), Deadnettle (Henbit <i>Lamium</i> spp.), Dock seedlings, Fat Hen, Field Madder, Fumitory, Hare's Ear (Treacle Mustard), Heliotrope, Hogweed (Wireweed), Horehound, Indian Hedge Mustard, Lesser Swinecress, Mouse-ear Chickweed, Mountain Sorrel, (Purple Calandrinia), Rough Poppy, Scarlet Pimpernel, Scrub Nettle (Stinging Nettle), Shepherds Purse, Sorrel seedlings, Sowthistle, Spiny Emex (Doublegee, Threecornered Jack(s)), Storksbill, Toad Rush, Wild Radish, Wild Turnip, Winter Grass, Yellow Burrweed.	Vic & Tas only	180g (Light sandy soils, i.e., sandy loams to loamy sands)  280g (Medium soils, loams, silt plus clay 40-60%)  380g (Heavy soils, clay loams, silt plus clay over 60%)	Apply to moist soils while weeds are young and actively growing and barley plants are at the 5-leaf stage.  <b>DO NOT</b> use on Barley grown on alkaline light sandy soils with pH over 7.  <b>DO NOT</b> use on Barley grown on light sandy soils (Tas).
Barley	Buxbaum's Speedwell, Powell's Amaranth, Stagger weed, Sun Spurge, Great Brome	Tas only		

Crop/ Situation	Weeds Controlled	State	Rate/ha	Critical Comments
Barley (all varieties excluding Morrell)	<p>Capeweed, Common Cotula, Doublegee, Fumitory, Indian Hedge Mustard, Toadrush, Wild Radish, Wild Turnip, Wireweed.</p> <p>Brome grass (<i>Bromus diandrus</i> &amp; <i>B.rigidus</i>)</p> <p>Annual Ryegrass, Barley grass</p>	WA only (sandy- sandy loam soils)	135-150g plus Trifluralin 480 at 1L/ha and incorporated by sowing (IBS)	<p><b>DO NOT</b> apply to Morrell Barley. Use the lower rate in lighter soils, and the higher rate in heavier soils. Apply to moist soil for best efficacy. Seed below herbicide band at 5cm. Reduced control may occur under high weed burdens (&gt;100/m<sup>2</sup>). <b>Ensure Imtrade Metribuzin 750 WG Herbicide is fully dispersed in the boom spray tank prior to the addition of Trifluralin 480. Refer to the application section of this and the Trifluralin label for complete instructions on product use.</b></p>
Chick Peas, Faba Beans, Lentils, Vetch	Refer to the Barley section	SA, Vic & WA only	<p>180 g (Light sandy soils, i.e., sandy loams to loamy sands)</p> <p>280g (Medium soils, loams, silt plus clay 40-60%)</p> <p>380g (Heavy soils, clay loams, silt plus clay over 60%)</p>	<p><b>Crop:</b> ONLY apply post-sowing, pre-emergence at least 5 cm deep. <b>DO NOT apply post-emergence as crop injury will occur.</b></p> <p><b>Weeds:</b> Weeds should be from pre-emergence to 3-leaf stage except Wireweed (Hogweed), which should not be beyond the cotyledon stage. Best results for grass and Wireweed control are obtained when rain or irrigation follows within a few days of spraying.</p>

Crop/ Situation	Weeds Controlled	State	Rate/ha	Critical Comments
Lupins	Capeweed, Doublegee, Wild Radish	WA only	100-150g plus 100 mL Difen Options Selective Herbicide	For use on varieties Gunurru, Meritt, Myallie and Kayla only. <b>DO NOT</b> use on Danja, Kiev mutant or Albus lupins. Suppression of larger Doublegee only. Wild Radish weed size up to 250mm only. Apply to the actively growing lupins from the 3-4 leaf stage onwards. Use higher rates for heavy weed burden and larger weed size. Apply before weeds are covered by crop canopy. <b>DO NOT</b> apply wetting agents or spay oils with this mixture. Some transient leaf burn may occur after application, but this will not affect yield. Best results are obtained if good soil moisture exists at, and is received soon after application. Results may be less effective south of the Great Eastern Highway. Refer to critical comments section on the Difen Options Selective Herbicide label.

Crop/ Situation	Weeds Controlled	State	Rate/ha	Critical Comments
Peas (Field and Green) ( <b>not</b> Snow Peas)	<p>Refer to Barley section plus: Buchan Weed, Ball Mustard, Fog Grass (SA only), Dwarf Nettle (Tas only)</p> <p><b>Suppression only:</b> Sub clover and Docks (Broadleaved Dock – Tas only), Wild Oats</p>	NSW, Vic, SA & Tas only	180g (Light sandy soils, i.e. sandy loams to loamy sands)	For use on varieties Gunurru, Meritt, Myallie and Kayla only. <b>DO NOT</b> use on Danja, Kiev mutant or Albus lupins. Suppression of larger Doublegee only. Wild Radish weed size up to 250mm only. Apply to the actively growing lupins from the 3-4 leaf stage onwards. Use higher rates for heavy weed burden and larger weed size. Apply before weeds are covered by crop canopy. <b>DO NOT</b> apply wetting agents or spay oils with this mixture. Some transient leaf burn may occur after application, but this will not affect yield. Best results are obtained if good soil moisture exists at, and is received soon after application. Results may be less effective south of the Great Eastern Highway. Refer to critical comments section on the Difen Options Selective Herbicide label.
			280g (Medium soils, loams, silt plus clay 40-60%)	
			380g (Heavy soils, clay loams, silt plus clay over 60%)	
Green Peas (post-emergence) ( <b>not</b> Snow Peas)		Qld only	470g	Apply to medium soils
			700g	Apply to soils high in clay or organic matter content when grass weeds are a problem.
			370 - 470g	Refer to Critical Comments for Peas (Field and Green)

Crop/ Situation	Weeds Controlled	State	Rate/ha	Critical Comments
Potatoes	Refer to Barley section plus: Buchan Weed, Ball Mustard, Fog Grass (SA only), Dwarf Nettle (Tas only)  <b>Suppression only:</b> Sub clover and Docks (Broadleaved Dock – Tas only), Wild Oats	ACT, NSW, Vic, SA, Tas & WA only	470g	Apply after rain or irrigation when crop has no more than 6 leaves or is no more than 10 cm high (many plants may be smaller). <b>CROPS HILLED OR MOLDED UP</b> -Hill or mold up as the crop emerges. Apply 1-4 weeks later as soon as weed seedlings have emerged. If surface crusting is not a problem no further cultivation should be necessary. <b>CROPS NOT HILLED AT OR SOON AFTER PLANTING</b> -Spray when weed seedlings have emerged. No cultivation should be necessary until crop is ready for final hilling. Additional cultivation may be necessary when surface crusting is a problem.
			700g	Where summer grasses are a problem.
		Vic, Bellarine Peninsula only	280 - 370g	Apply to winter crops on light soils only.
		Qld only (Winter crop)	470g	Apply at first emergence of potatoes to control broadleaf weeds.
		South Qld only (Summer crop)	470g - 950g	Where summer grasses such as <i>Echinochloa crus galli</i> , <i>Eleusine</i> , <i>Digitaria</i> and <i>Panicum</i> spp. are a problem.
		Atherton Tablelands (Qld only)	950g	Apply to summer crops at first emergence of potatoes to control broadleaved weeds and more troublesome grasses such as <i>Eleusine</i> , <i>Digitaria</i> and <i>Panicum</i> spp.

Crop/ Situation	Weeds Controlled	State	Rate/ha	Critical Comments
Irrigated Soybeans	Apple of Peru, Cobbler's Pegs, Pigweed, Noogoora Burr, Common Thornapple, Summer Grass, Caltrop	ACT, NSW, NT only	500g	Apply pre-plant incorporated. Where grass weeds other than summer grass are a problem apply in combination with Trifluralin 480. (Refer to label for directions for use and incorporation techniques). Use only on soils classified as silty clay-heavy clay.
Irrigated Soybeans (Variety Davis only)	Refer to weeds listed for soybeans plus Anoda weed, Awnless Barnyard grass. Suppression only of Bellvine	Qld, NT only	750g	Apply pre-plant incorporated. Where grass weeds other than summer grass are a problem apply in combination with Trifluralin 480. (Refer to label for directions for use and incorporation techniques). Use only on soils classified as silty clay-heavy clay.
Tomatoes Inter-row spaces of furrow irrigated tomatoes	Refer to Peas section plus: Barnyard grass, Summer grass, Crowsfoot grass, Blackberry Nightshade, Pigweed, Black Pigweed, Bull's Head, Green Summer Grass		950g or 9.5g/100 m of row	Apply in a 1 metre band centered on the interrow either during or shortly after the formation of the final furrow. Apply to soil moist at the surface.  <b>DO NOT</b> exceed spraying pressures of 200 kPa as excessive drift will occur.



Crop/ Situation	Weeds Controlled	State	Rate/ha	Critical Comments
Wheat (Blade only)	<p>Capeweed, Common Cotula, Doublegee, Fumitory, Indian Hedge Mustard, Toad Rush, Wild Turnip, Wild Radish, Wireweed.</p> <p>Brome Grass (<i>Bromus diandrus</i> and <i>B. rigidus</i>)</p> <p>Annual Ryegrass, Barley Grass (refer to critical comments)</p>	WA (Great northern region only) sandy- sandy loam soils	<p>200g and incorporated by sowing (IBS)</p> <p>200g/ha plus a Trifluralin 480 at 1L/ha and incorporated by sowing (IBS)</p>	<p>Imtrade Metribuzin 750 WG Herbicide has increased crop safety and gives more reliable weed control when incorporated by sowing and trailing harrows.</p> <p>Reduced grass control may occur under high weed burdens (&gt;100/m<sup>2</sup>).</p> <p>To achieve good control of Annual Ryegrass and Barley grass add Trifluralin 480 at 1L/ha.</p> <p><b>Ensure Imtrade Metribuzin 750 WG Herbicide is fully dispersed in the boom spray tank prior to the addition of Trifluralin 480.</b></p> <p><b>Refer to the application section of this and the Trifluralin label for complete instructions on product use.</b></p>
Wheat and Barley	Toad Rush	Vic, ACT, NSW, SA only	100g	<p>Toad rush should be sprayed at the 2-4 leaf stage for optimum control.</p> <p>Spray when soil moisture is plentiful and soil is moist to the surface. Take advantage of dew on the soil surface.</p>
Oats		Vic only		

Crop/ Situation	Weeds Controlled	State	Rate/ha	Critical Comments
White Lupins	Buchan Weed, Capeweed, Charlock, Chickweed, Deadnettle ( <i>Lamium</i> spp.), Fat Hen, Fumitory (Common and Wall Fumitory - Tas), Wireweed (Hogweed), Lesser Swinecress, Purple Calandrinia, Scarlet Pimpernel, Spiny Emex, Corn Spurry (Spurry -Tas), Stinging Nettle (Dwarf Nettle -Tas), Toad Rush, Winter grass, Heliotrope, Wild Oats, Wild Turnip, Wild Radish. <b>Suppression only:</b> Sub clover and Docks (Broadleaved Dock -Tas)	Tas only	400g	<b>Spring application</b> – Apply after sowing but before crop emerges.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL  
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIOD**

**TOMATOES: DO NOT APPLY LATER THAN 21 DAYS BEFORE HARVEST.**

**BARLEY: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION.**

**OTHER CROPS: NOT REQUIRED IF USED AS DIRECTED.**

## GENERAL INSTRUCTIONS:

Imtrade Metribuzin 750 WG Herbicide is non-corrosive to equipment, non-flammable and non-volatile. The degree of weed control and duration of effect will vary with the amount of herbicide applied, soil texture, rainfall and other conditions.

## CROP SAFETY

**DO NOT** apply to crops that are stressed by any cause (such as severe weather conditions, water-logging, drought, excessive soil acidity or alkalinity, poor nutrient status, disease, nematode or insect damage) as crop injury may result.

- Barley: **DO NOT** apply to crops grown on sandy soils or soils low in organic matter.
- Peas **DO NOT** apply to Laura, Pennant or White Brunswick varieties as phytotoxicity may occur.
- (Field & Green): **DO NOT** apply any later than the one node stage to Alma and Dinkum peas.  
**DO NOT** spray pea crops in soils or areas where collar or stem rot is likely to occur.
- Potatoes: **DO NOT** apply to crops grown on sandy soils as injury to the crops may result, appearing as chlorosis or minor necrosis.  
**DO NOT** apply to Kurell potatoes as phytotoxicity may occur.  
**DO NOT** apply to Atlantic and Pontiac varieties – WA only.  
**DO NOT** use on dry-grown, non-irrigated crops.
- Soybeans: **DO NOT** use on varieties Cannapolis, Hill and Semstar as crop injury may occur.  
**DO NOT** use on soils with a calcareous surface layer as crop injury may occur.  
**DO NOT** apply over atrazine residues as crop injury may result.  
**DO NOT** apply in combination with soil applied organophosphate insecticides as crop injury may result.  
**DO NOT** apply to soils with less than 0.5% organic matter as crop injury may result.  
**DO NOT** apply to dryland soybeans.  
**DO NOT** apply to soils of pH greater than 8 that are prone to water-logging as crop injury may occur.
- Lupins: **DO NOT** apply to varieties Danja, Kiev Mutant or Albus Lupins.  
**DO NOT** apply to Lupin crops that have received 2.5 litres or more of Simazine pre-seeding, or lupin crops that are showing symptoms of Simazine or Atrazine damage.

## **COMPATIBILITY**

Imtrade Metribuzin 750 WG Herbicide is compatible with the herbicides Chlorsulfuron, Metsulfuron, Glyphosate, Pendimethalin, Propargite, Difen Options Selective Herbicide and Trifluralin 480.

## **MIXING**

Imtrade Metribuzin 750 WG Herbicide is a water dispersible granule formulation to be mixed with water and applied as a spray. Before spraying, calibrate equipment to determine the quantity of water necessary to uniformly cover the measured area to be treated. Carefully weigh the amount of Imtrade Metribuzin 750 WG Herbicide required and add to partially filled spray tank with agitation operating. Top up to the correct volume with water.

## **THE MATERIAL MUST BE KEPT IN SUSPENSION AT ALL TIMES BY CONTINUOUS AGITATION.**

In tank mixes Imtrade Metribuzin 750 WG Herbicide must be in suspension before adding the companion herbicide, insecticide, spray adjuvant or liquid fertiliser.

### **Pre-planting Incorporation Treatment:**

#### **Imtrade Metribuzin 750 WG Herbicide plus Trifluralin 480**

Add Imtrade Metribuzin 750 WG Herbicide to water in spray tank with agitator running as above. Maintain agitation then add Trifluralin 480 at the recommended rate (Refer to Trifluralin 480 label).

## **APPLICATION - CROPS**

### **General:**

Best results are obtained if treatment is made to moist soil and moisture is supplied by rainfall or sprinkler irrigation (6-12mm) within 2 weeks after application. If heavy rains occur soon after application, injury to the crop may result.

### **Aerial Application- (Qld only):**

**Peas** - apply in a minimum of 20 litres water per hectare.

**Potatoes** - apply in a minimum of 50 litres water per hectare. Avoid overlapping of spray swath. Avoid spraying in still conditions and in wind likely to cause drift.

**Pre-emergence and Post-emergence (To crop):**

Imtrade Metribuzin 750 WG Herbicide should not be applied when weeds are in the young seedling stage (no more than 3 true leaves). Under dry conditions there is no root uptake of Imtrade Metribuzin 750 WG Herbicide and results against grass weeds and Wireweed (Hogweed) will be poor. Except for small areas, use fixed boom power sprayers set at 200-300kPa (for tomatoes do not exceed 200kPa) and fitted with fan nozzles. Apply a minimum of 80L-100L prepared spray/ha. Openings in screens should be equal to or larger than 50 mesh. Agitate by mechanical or hydraulic means in spray tank. When spraying avoid overlapping. Shut off spray booms whilst starting, turning, slowing or stopping otherwise crop injury may result.

**Pre-plant incorporated - Soybeans only:****Imtrade Metribuzin 750 WG Herbicide alone or in combination with Trifluralin 480**

Spray anytime from 4 weeks before sowing until just before sowing. Use fixed boom power sprayers set at 200-300 kPa and fixed with fan nozzles. Use 120-500 litres water per hectare. Openings in screens should be equal to or larger than 80 mesh. Agitate by mechanical or hydraulic means in the spray tank. When spraying avoid overlapping. Shut off spray booms while starting, turning, slowing and stopping otherwise crop injury may result.

**Imtrade Metribuzin 750 WG Herbicide alone**

If possible spray and incorporate within the same operation. If not possible incorporate spray within 24 hours of application using Go-devil or Lilliston cultivators set to work 5 cm deep.

**Imtrade Metribuzin 750 WG Herbicide plus Trifluralin 480**

After spraying mix soil within 4 hours. Spray and incorporate during the one operation if possible. (Refer to Trifluralin 480 label for specific incorporation details).

**Sprayer Cleanup**

Thoroughly clean all traces of Imtrade Metribuzin 750 WG Herbicide and/or Trifluralin 480 from application equipment immediately after use. Flush tank, pump, hoses and boom with several changes of water after removing nozzle tips and screens (clean these parts separately).

## RESISTANT WEEDS WARNING

### GROUP 5 HERBICIDE

Imtrade Metribuzin 750 WG Herbicide is a member of the Triazines group of herbicides. Imtrade Metribuzin 750 WG Herbicide has the inhibitors of photosynthesis at photosystem II mode of action. For weed resistance management Imtrade Metribuzin 750 WG Herbicide is a Group 5 Herbicide. Some naturally occurring weed biotypes resistant to Imtrade Metribuzin 750 WG Herbicide and other Group 5 Herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides used repeatedly. These resistant weeds will not be controlled by Imtrade Metribuzin 750 WG Herbicide or other Group 5 Herbicides.

Annual Ryegrass biotypes resistant to diclofop-methyl and other ryegrass specific herbicides may be resistant to Imtrade Metribuzin 750 WG Herbicide. Imtrade Australia Pty Ltd recommends resistance testing of the population to ensure susceptibility prior to use.

Since the occurrence of resistant weeds is difficult to detect prior to use, Imtrade Australia Pty Ltd accepts no liability for any losses that may result from the failure of Imtrade Metribuzin 750 WG Herbicide to control resistant weeds. Large numbers of healthy surviving weeds can be an indication that resistance is developing. Efforts should be made to prevent seed set of the survivors.

**DO NOT** make more than one application of a Group 5 Herbicide to a crop in the same season, either pre-sowing incorporated by sowing or post crop and weed emergence. If the user suspects that Group 5 Herbicide resistant weeds are present, Imtrade Metribuzin 750 WG Herbicide or other Group 5 Herbicides should not be used.

Strategies to minimise the risk of herbicide resistance are available. Consult your farm chemical supplier, consultant, local Department of Agriculture or Primary Industries.

## **PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS**

Prevent drift of spray to sensitive plants. Keep from contact with fertilisers, insecticides, fungicides and seeds.

**DO NOT** apply (except as recommended for crop use) or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots, as injury or loss may occur.

**DO NOT** plant treated area to sensitive crops such as brassicas, sunflowers, beetroot, cucurbits, lettuce or onions for at least 6 months following application of Imtrade Metribuzin 750 WG Herbicide as crop injury may result. Thoroughly work the soil prior to planting of these crops.

**DO NOT** apply under meteorological conditions or from spraying equipment, which could be expected to cause spray to drift onto nearby susceptible plants, adjacent crops, crop lands or pastures.

Tomatoes - **DO NOT** plant follow-on crops of cucurbits or capsicums when applying Imtrade Metribuzin 750 WG Herbicide to autumn plantings.

## **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

**DO NOT** contaminate streams, rivers or waterways by cleaning of equipment, disposal of wastes or used containers.

## **STORAGE AND DISPOSAL**

Store in the closed, original container in a dry, cool well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. Keep from contact with fertilizers, insecticides, fungicides and seeds. Triple or preferably pressure rinse container before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Empty containers and product should not be burnt.

## **SAFETY DIRECTIONS**

Harmful if inhaled or swallowed. Will irritate the eyes. Avoid contact with eyes and skin. **DO NOT** inhale dust or spray mist. Repeated minor exposure may have a cumulative poisoning effect. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow length PVC gloves and face shield or goggles. Wash hands after use.

## FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (*Phone Australia 13 11 26*).

### Additional GHS Hazard & Precautionary Statements

•Very toxic to aquatic life with long-lasting effects; •**DO NOT** get on clothing; •Wash contacted areas thoroughly after handling; •**DO NOT** eat, drink or smoke when using this product; •Rinse skin or shower with water; •**IF SWALLOWED:** Call a POISON CENTRE or doctor if you feel unwell; •**IF SWALLOWED:** Rinse mouth. Do **NOT** induce vomiting; •If eye irritation persists: Seek medical advice; •Suitable extinguishing media are carbon dioxide, dry chemical, foam, water fog.

## SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from your supplier or Imtrade Australia Pty Ltd website at [www.imtrade.com.au](http://www.imtrade.com.au)

## CONDITIONS OF SALE

Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Imtrade Australia Pty Ltd has any authority to add to or alter these conditions.

<b>UN 3077</b>	<b>ENVIRONMENTALLY HAZARDOUS SUBSTANCES, SOLID, N.O.S.</b>
<b>IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE</b>	
<b>PG III</b>	<b>HAZCHEM 2Z</b>

APVMA Approval No: 63658/60417

