

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

STORAGE AND DISPOSAL (5L, 10L & 20L only)

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Triple or (preferably) pressure rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers or product.

SAFETY DIRECTIONS

Avoid inhaling spray mist. Will irritate the eyes and skin. Avoid contact with eyes and skin. When preparing spray and using the prepared spray, wear rubber gloves. Wash hands after use. After each day's use, wash gloves.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26).

Refer to attached Leaflet for additional GHS
Hazard & Precautionary Statements

IMTRADE ORDNANCE 250 SL HERBICIDE

ACTIVE CONSTITUENTS: 250 g/L IMAZAPYR present as the isopropylamine salt

GROUP **2** HERBICIDE

For the control of various annual and perennial weeds in non-crop situations
as indicated in the Directions for Use table.

Important: Read the attached Leaflet before use.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet which can be obtained from your supplier or Imtrade Australia Pty Ltd website at www.imtrade.com.au

CONDITIONS OF SALE

Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever, or howsoever arising through negligence or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express conditions that the purchaser does not rely on Imtrade Australia's skill or judgement in purchasing or using the same and every person dealing with this product does so at his/her own risk absolutely. No representative of Imtrade Australia Pty Ltd has any authority to alter these conditions.

In a Transport Emergency Dial 000
Police or Fire Brigade

APVMA Approval No: 68366/58231

GROUP
B

FORM
SL

CONTENTS
1L-20L



Batch No.
DOM:

 **IMTRADE**
CropScience
17 Ocean St, Kwinana Beach WA 6167
08 9419 0333 | www.imtrade.com.au

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



IMTRADE ORDNANCE 250 SL HERBICIDE

ACTIVE CONSTITUENT: 250 g/L IMAZAPYR
present as the isopropylamine salt

| | | |
|--------------|----------|------------------|
| GROUP | 2 | HERBICIDE |
|--------------|----------|------------------|

For the control of various annual and perennial weeds in
non-crop situations as indicated in the Directions for Use table

**IMPORTANT: READ THIS LEAFLET THOROUGHLY
BEFORE OPENING OR USING THIS PRODUCT**

APVMA Approval No: 68366/58231

Imtrade Australia Pty Ltd
ABN 13 090 151 134
17 Ocean Street, Kwinana WA 6167
Tel: 1800 171 799 Fax: 1800 171 788
Web: www.imtrade.com.au

DIRECTIONS FOR USE**RESTRAINTS**

DO NOT apply by aircraft as spray drift may cause damage to desirable surrounding species.

| SITUATION | WEEDS CONTROLLED | STATES | RATE | CRITICAL COMMENTS |
|--|---|------------|-------|---|
| Around agricultural buildings and other farm non-crop situations, commercial, industrial and public service areas, rights of way and waste land away from desirable vegetation | Annual Ryegrass (<i>Lolium rigidum</i>), Barnyard Grass (<i>Echinochloa colona</i>), Blady Grass (<i>Imperata cylindrica</i>), Buckbush (<i>Salsola kali</i>), Deadnettle (<i>Lamium amplexicaule</i>), Feathertop Rhodes Grass (<i>Chloris virgate</i>), Fleabane (<i>Conyza canadensis</i>), Hexham Scent (<i>Melilotus indicus</i>), Liverseed Grass (<i>Urochloa panicoides</i>), London Rocket (<i>Sisymbrium mirio</i>), Mayne's Pest (<i>Verbena tenuisecta</i>), Medics (<i>Medicago</i> spp.), Mintweed (<i>Salvia reflexa</i>), Blackberry Nightshade (<i>Solanum nigrum</i>), Paterson's Curse (<i>Echium plantagineum</i>), Plains Grass (<i>Stipa aristiglumis</i>), Prairie Grass (<i>Bromus unioloides</i>), Prickly Lettuce (<i>Lactuca serriola</i>), Purple Top (<i>Verbena bonariensis</i>), Scented Top (<i>Capillipedium</i> spp.), Sedges (<i>Cyperus</i> spp.), Slender Celery (<i>Apium leptophyllum</i>), Sowthistle (<i>Sonchus oleraceus</i>), St Barnaby's Thistle (<i>Centaurea solstitialis</i>), Summer Grasses (<i>Digitaria</i> spp.), Thornapples (<i>Datura</i> spp.), Turnip Weed (<i>Rapistrum rugosum</i>), Variegated Thistle (<i>Silybum marianum</i>), Vasey Grass (<i>Paspalum urvillei</i>), Warrego Grass (<i>Paspalidium jubiflorum</i>), Wild Oats (<i>Avena</i> spp.), Wild Turnip (<i>Brassica tournefortii</i>), Wireweed (<i>Polygonum aviculare</i>) | All States | 4L/ha | Applications can be made pre- or post-emergence. Post-emergence applications also provide pre-emergence control. Medium Volume: 4L of Imtrade Ordinance 250 SL Herbicide should be applied using spray volumes ranging between 50 and 200L/ha. High Volume: Apply sufficient spray to wet the surfaces visibly without producing runoff. Measure how much spray is required to cover 10m x 10m. Add Imtrade Ordinance 250 SL Herbicide at the rate of 40 mL in this measured volume. |

| SITUATION | WEEDS CONTROLLED | STATES | RATE | CRITICAL COMMENTS |
|--|---|------------|-------|--|
| Around agricultural buildings and other farm non-crop situations, commercial, industrial and public service areas, rights of way and waste land away from desirable vegetation | Bokhara Clover (<i>Melilotus albus</i>), Bracken (suppression only) (<i>Pteridium esculentum</i>), Capeweed (<i>Arctotheca calendula</i>), Couch (<i>Cynodon dactylon</i>), Docks (<i>Rumex</i> spp.), Johnson Grass (<i>Sorghum halepense</i>), Panics (suppression only) (<i>Panicum</i> spp.), Paspalum (<i>Paspalum dilatatum</i>), Queensland Bluegrass (<i>Dichanthium sericeum</i>), Stinging Nettle (<i>Urtica incise</i>) | All States | 6L/ha | Applications can be made pre- or post-emergence. Post-emergence applications also provide pre-emergence control. Medium Volume: 6L of Imtrade Ordnance 250 SL Herbicide should be applied using spray volumes ranging between 50 and 200L/ha. High Volume: Apply sufficient spray to wet the surfaces visibly without producing runoff. Measure how much spray is required to cover 10m x 10m. Add Imtrade Ordnance 250 SL Herbicide at the rate of 60 mL in this measured volume. |

| SITUATION | WEEDS CONTROLLED | STATES | RATE | CRITICAL COMMENTS |
|--|--|----------|----------------|---|
| Around agricultural buildings and other farm non-crop situations, commercial, industrial and public service areas, rights of way and waste land away from desirable vegetation | Rubber Vine (<i>Cryptostegia grandiflora</i>) | Qld only | 4 mL/L water | Post-emergence application: Apply sufficient spray to wet the surface visibly to the point of runoff. For effective control, apply when the plant is actively growing and not stressed, yellowing or bearing pods. Flowering is a good indication of the best time for application. Use high volume application as detailed in the General Instructions section. |
| | Thunbergia (<i>Thunbergia</i> spp.) | | 7.5 mL/L water | Post-emergence application: Apply sufficient spray to wet the surface visibly to the point of runoff. For effective control, apply when the plant is actively growing. Use high volume application as detailed in the General Instructions section. |

| SITUATION | WEEDS CONTROLLED | STATES | RATE | CRITICAL COMMENTS |
|--|--|------------|-------|--|
| Irrigation channels – supply channels and tail drains leading to recirculation dams only | Barnyard Grass (<i>Echinochloa colona</i>), Bathurst Burr (<i>Xanthium spinosum</i>), Bellvine (<i>Ipomoea plebian</i>), Bladder Ketmia (<i>Hibiscus trionum</i>), Brown Beetle Grass (<i>Diplachne fusca</i>), California Burr (<i>Xanthium orientale</i>), Caltrop (<i>Tribulus terrestris</i>), Couch Grass (<i>Cynadon dactylon</i>), Cumbungi (<i>Typha spp.</i>), Gooseberry (<i>Physalis viscosa</i>), Johnson Grass (<i>Sorghum halepense</i>), Liverseed Grass (<i>Urochloa panicoides</i>), Mintweed (<i>Salvia reflexa</i>), Noogoora Burr (<i>Xanthium pungens</i>), Nutgrass (<i>Cyperus rotundus</i>), <i>Paspalum spp.</i> , Pumpkin Vine (<i>Operculina brownii</i>), Rosella (<i>Hibiscus heterophyllous</i>), Sesbania Pea (<i>Sesbania brachycarpa</i>), Sowthistle (<i>Sonchus oleraceus</i>), Spider Grass (<i>Brachyachne convergens</i>), Summer Grass (<i>Digitaria ciliaris</i>), Thornapples (<i>Datura spp.</i>), Turnip Weed (<i>Rapistrum rugosum</i>), Volunteer Cereals, Watermelon (<i>Citrullus lanatus</i>), Wireweed (<i>Polygonum aviculare</i>) | All States | 3L/ha | <p>Apply during the non-crop season and when the channel is not in use. Drain channels to be treated prior to use. Apply only to irrigation delivery and recirculation channels which may be flushed to a recirculation dam or settling pond. DO NOT apply where flushing water cannot be disposed of safely on-farm.</p> <p>Waater from the initial flushing of treated channels must be retained on-farm.</p> <p>Applications may be made pre or post-emergence. Post-emergence applications also provide pre-emergence control. For boom sprays, apply in 50 to 200L water/ha.</p> <p>For high volume e.g. handguns, measure how much spray volume is required to cover 10m x 10m and add Imtrade Ordnance 250 SL Herbicide at the rate of 30 mL in this measured volume. Apply to wet surfaces visibly, without producing runoff. Apply only to exposed irrigation channels.</p> <p>DO NOT apply to water.</p> <p>DO NOT apply to irrigation channels in situations where there is a risk of soil or silt movement or water run-off leading to Imtrade Ordnance 250 SL Herbicide entering irrigation water.</p> |

| SITUATION | WEEDS CONTROLLED | STATES | RATE | CRITICAL COMMENTS |
|--|--|------------|-------|--|
| Irrigation channels – supply channels and tail drains leading to recirculation dams only | Barnyard Grass (<i>Echinochloa colona</i>), Bathurst Burr (<i>Xanthium spinosum</i>), Bellvine (<i>Ipomoea plebian</i>), Bladder Ketmia (<i>Hibiscus trionum</i>), Brown Beetle Grass (<i>Diplachne fusca</i>), California Burr (<i>Xanthium orientale</i>), Caltrop (<i>Tribulus terrestris</i>), Couch Grass (<i>Cynodon dactylon</i>), Cumbungi (<i>Typha spp.</i>), Gooseberry (<i>Physalis viscosa</i>), Johnson Grass (<i>Sorghum halepense</i>), Liverseed Grass (<i>Urochloa panicoides</i>), Mintweed (<i>Salvia reflexa</i>), Noogoora Burr (<i>Xanthium pungens</i>), Nutgrass (<i>Cyperus rotundus</i>), <i>Paspalum spp.</i> , Pumpkin Vine (<i>Operculina brownii</i>), Rosella (<i>Hibiscus heterophyllous</i>), Sesbania Pea (<i>Sesbania brachycarpa</i>), Sowthistle (<i>Sonchus oleraceus</i>), Spider Grass (<i>Brachyachne convergens</i>), Summer Grass (<i>Digitaria ciliaris</i>), Thornapples (<i>Datura spp.</i>), Turnip Weed (<i>Rapistrum rugosum</i>), Volunteer Cereals, Watermelon (<i>Citrullus lanatus</i>), Wireweed (<i>Polygonum aviculare</i>) | All States | 3L/ha | <p>DO NOT apply Imtrade Ordnance 250 SL Herbicide to flooded or saturated soil.</p> <p>If the entire (empty) channel surface is to be treated, delay reuse of the channel for as long as possible (at least 6 weeks) for Imtrade Ordnance 250 SL Herbicide to be absorbed into the soil particles. Then flush with water, returning flushing water to a settling pond, before allowing irrigation water to flow through the channel.</p> <p>To avoid the risk of drift damage, application should NOT be made to channel banks at a time when susceptible crops are in the direct vicinity.</p> <p>Applications should NOT be undertaken when wind is blowing towards crops.</p> <p>Applications in early Autumn (March) to channels with minimum weed growth will give best performance. Application should give 6 months effective weed control.</p> <p>Imtrade Ordnance 250 SL Herbicide should NOT be applied more than once per annum to the same channel.</p> <p>REFER TO 'PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS' SECTION OF LEAFLET.</p> |

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

GENERAL INSTRUCTIONS

Imtrade Ordnance 250 SL Herbicide is an aqueous solution of herbicide which will kill a range of standing weeds and emerging seedlings. It stops growth rapidly, although symptoms may take over 30 days to develop. Affected plants are discoloured and often growth is distorted.

The active material in Imtrade Ordnance 250 SL Herbicide is rapidly absorbed through foliage and roots, and translocated throughout plants. Foliage treatments will provide pre-emergence control as the active material is released to the soil following plant decay.

Residual control of weeds, for which the lower rate of Imtrade Ordnance 250 SL Herbicide is recommended, will increase when the higher rate is used.

RESISTANT WEEDS WARNING

GROUP 2 HERBICIDE

Imtrade Ordnance 250 SL Herbicide is a member of the imadazolinones group of herbicides. Imtrade Ordnance 250 SL Herbicide has the inhibitors-of-acetolactate-synthase (ALS) mode of action. For weed resistance management, Imtrade Ordnance 250 SL Herbicide is a Group 2 Herbicide. Some naturally occurring weed biotypes resistant to Imtrade Ordnance 250 SL Herbicide and other inhibitors-of-acetolactate-synthase (ALS) herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Imtrade Ordnance 250 SL Herbicide or other inhibitors-of-acetolactate-synthase (ALS) herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Imtrade Australia Pty Ltd accepts no liability for any loss that may result from the failure of Imtrade Ordnance 250 SL Herbicide to control resistant weeds.

APPLICATION

Faster control of standing weeds will result if growing conditions are favourable at and after application.

Rainfall within two hours of the spray mixture drying on leaves may result in reduced weed control.

SPRAYING INSTRUCTIONS

Medium Volume Spraying: Apply through a spray boom calibrated to ensure even coverage. Use spray nozzles suitable to prevent spray drift onto surrounding desirable vegetation.

High Volume Application: Use a handgun or knapsack to apply sufficient spray to wet the surface visibly without producing runoff.

MIXING

Pour the required amount of Imtrade Ordnance 250 SL Herbicide into a spray container which already holds the total amount of water required. Mix thoroughly. If Imtrade Ordnance 250 SL Herbicide is added during filling, foaming may occur. If excessive foaming is a problem, add a suitable antifoaming agent. Consult your distributor for specific information.

Imtrade Ordnance 250 SL Herbicide may be applied in hard or soft water.

Imtrade Ordnance 250 SL Herbicide is corrosive to mild steel. Use **ONLY** stainless steel, fiberglass, plastic or plastic-line containers for mixing, storage and application.

COMPATIBILITY

For faster knockdown of weeds, certain other herbicides may be mixed with Imtrade Ordnance 250 SL Herbicide.

Bipyridyl herbicides (e.g. Paraquat) are **NOT** compatible with Imtrade Ordnance 250 SL Herbicide. Mixtures of this type will result in poor weed control.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures or other non-target aquatic areas.

DO NOT apply on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved onto contact with their roots.

DO NOT apply where runoff water may flow onto agricultural land or other desirable vegetated areas.

Dispose of washings away from desirable plants and their roots.

PROTECTION OF LIVESTOCK

Low hazard to bees – may be applied to plants at any time.

DO NOT graze or cut for stock feed, vegetation which has been treated with Imtrade Ordnance 250 SL Herbicide.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with this product or used container.

STORAGE AND DISPOSAL (1L only)

Store product in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. Rinse containers before disposal. Add rinsings to the spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers or product.

STORAGE AND DISPOSAL (5L, 10L & 20L only)

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Triple or (preferably) pressure rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers or product.

SAFETY DIRECTIONS

Avoid inhaling spray mist. Will irritate the eyes and skin. Avoid contact with eyes and skin. When preparing spray and using the prepared spray, wear rubber gloves. Wash hands after use. After each day's use, wash gloves.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26).

Additional GHS Hazard & Precautionary Statements

●Harmful to aquatic life with long-lasting effects; ●Wash contact areas thoroughly after handling; ●**DO NOT** eat, drink or smoke when using this product; ●**IF SWALLOWED:** Rinse mouth. Do **NOT** induce vomiting; ●**IF ON SKIN:** Wash with plenty of soap and water; ●Not combustible. Use extinguishing media suited to burning materials.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet which can be obtained from your supplier or Imtrade Australia Pty Ltd website at www.imtrade.com.au

CONDITIONS OF SALE

Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever, or howsoever arising through negligence or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express conditions that the purchaser does not rely on Imtrade Australia's skill or judgement in purchasing or using the same and every person dealing with this product does so at his/her own risk absolutely. No representative of Imtrade Australia Pty Ltd has any authority to alter these conditions.

**IN A TRANSPORT EMERGENCY
DIAL 000
POLICE OR FIRE BRIGADE**

APVMA Approval No: 68366/58231