

STORAGE AND DISPOSAL (1L, 5L, 10L, 20L, 60L & 200L only)

Store in the closed original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemical on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Empty containers and product should **NOT** be burnt.

Envirodrum - Micro Matic Valve (110L)

Store the original sealed Envirodrum in a cool, well ventilated area. **DO NOT** store for prolonged periods in direct sunlight. **DO NOT** tamper with the Micro Matic valve or the security seal. **DO NOT** contaminate the Envirodrum with water or any other foreign matter. After each use of the product please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase.

SAFETY DIRECTIONS

Will damage eyes. Will irritate the skin. Avoid contact with eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. **DO NOT** inhale vapour or spray mist. When opening the container and preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length butyl rubber gloves and face shield or goggles. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (*Phone Australia 13 11 26; New Zealand 0800 764 766*). If swallowed **DO NOT** induce vomiting. Give a glass of water. If in eyes wash out immediately with water.

Refer to attached Leaflet for additional GHS Hazard & Precautionary Statements.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from your supplier or Imtrade Australia Pty Ltd website at www.imtrade.com.au

CONDITIONS OF SALE

Imtrade Australia Pty Ltd warrants that this product conforms to the chemical description on the label. The use of Imtrade Oxen 240 EC Herbicide being beyond the control of the manufacturer, Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Imtrade Australia Pty Ltd has any authority to add to or alter these conditions.

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

IMTRADE

OXEN 240 EC

HERBICIDE

ACTIVE CONSTITUENT: 240 g/L OXYFLUOREN
SOLVENTS: 605 g/L LIQUID HYDROCARBONS
100 g/L N-METHYL PYRROLIDONE

GROUP 14 HERBICIDE

For selective weed control of broadleaf weeds and some grasses as specified in the Directions for Use table.

IMPORTANT: Read the attached Leaflet before use

GROUP
14

FORM
EC

CONTENTS
1L-200L

UN 3082	ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S. (CONTAINS OXYFLUORFEN)
In a Transport Emergency Dial 000 Police or Fire Brigade	
PG III	HAZCHEM *3Z

APVMA Approval No: 56798/106771



Batch No.
DOM:

IMTRADE
CropScience

17 Ocean St, Kwinana Beach WA 6167
08 9419 0333 | www.imtrade.com.au

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



IMTRADE OXEN 240 EC HERBICIDE

ACTIVE CONSTITUENT: 240 g/L OXYFLUORFEN
SOLVENTS: 605 g/L LIQUID HYDROCARBONS
100 g/L N-METHYL PYRROLIDONE

GROUP	14	HERBICIDE
--------------	-----------	------------------

For Selective weed control of
broadleaf weeds and some grasses as
specified in the Directions for Use Table

**IMPORTANT: READ THIS LEAFLET THOROUGHLY
BEFORE OPENING OR USING THIS PRODUCT**

APVMA Approval No: 56798/106771

Imtrade Australia Pty Ltd
ABN 13 090 151 134
17 Ocean Street, Kwinana WA 6167
Tel: 1800 171 799 Fax: 1800 171 788
Web: www.imtrade.com.au

DIRECTIONS FOR USE

RESTRAINT: DO NOT disturb weeds by cultivation or sowing for 1 day following application to annual weeds and 7 days for perennial weeds to ensure herbicide absorption, unless specified in the CRITICAL COMMENTS.

CROP/SITUATION	WEEDS CONTROLLED	TIME OF APPLICATION	RATE	CRITICAL COMMENTS
<p>Trees (Fruit & Nut) and Vines not less than 3 years old as a Dormant Application, including: Grapevine, Olive Trees, Pome Fruit (e.g. Apple, Pear, Nashi, Quince), Stone Fruit (e.g. Apricot, Cherry, Nectarines, Peach, Plum), Tree Nuts (e.g. Almond, Macadamia, Pecan, Walnut)</p>	<p>Refer to Weeds Controlled List</p>	<p>Weed free soil</p>	<p>3 or 4L/ha</p>	<p>DO NOT apply Imtrade Oxen 240 EC Herbicide once bud swell has occurred. Apply to freshly cultivated weed free soil. Use the higher rate when longer residual control is required (up to 4 months). Where grass weeds are expected to be a major problem, or when control of a wider weed spectrum is required, mix the lower rate with 4-5L Oryzalin 500g/L per treated hectare. Read and follow all label directions.</p>
		<p>Emerged weeds present (4-6 leaf stage)</p>		<p>DO NOT apply Imtrade Oxen 240 EC Herbicide once bud swell has occurred. Use the higher rate when longer residual control is required (up to 4 months). When light infestations of weeds are present, apply as a tank mix with Glyphosate 360g/L or Paraquat 200g/L to produce both knockdown and residual control. Read and follow all label directions. Where weed growth is large and dense, weeds must be eliminated prior to application of Imtrade Oxen 240 EC Herbicide, using Glyphosate 360 g/L or mechanical means. Macadamias: Apply in 200 to 500L water/ha. Apply after harvest to prevent spray contacting nuts. Avoid spray contact with the foliage and stem. DO NOT apply to nuts on the ground.</p>

CROP/SITUATION	WEEDS CONTROLLED	TIME OF APPLICATION	RATE	CRITICAL COMMENTS
Fruit and Nut Trees, Vines including: Grapevine, Olive Trees, Pome Fruit (e.g. Apple, Pear, Nashi, Quince), Stone Fruit (e.g. Apricot, Cherry, Nectarines, Peach, Plum), Tree Nuts (e.g. Almond, Macadamia, Pecan, Walnut)	Annual Weeds Refer to Weeds Controlled list and Glyphosate 360 g/L label.	Actively growing emerged weeds	75 mL/ha plus Glyphosate 360g/L at recommended label rate	Addition of Imtrade Oxen 240 EC Herbicide to Glyphosate 360g/L will improve knockdown and increase the speed at which treated weeds develop visible symptoms of phytotoxicity (compared to results achieved with Glyphosate applied alone), and give control Annual Nettles (<i>Urtica</i> spp.) and Marshmallow (<i>Malva parviflora</i>). For rates of Glyphosate 360g/L, refer to the appropriate label. Read and follow all label directions. DO NOT apply the tank mix of Imtrade Oxen 240 EC Herbicide and Glyphosate 360g/L near trees or vines less than 3 years old unless they are effectively shielded from spray and spray drift.
	Most annual grasses and broadleaf weeds. Refer to Weeds Controlled List and Imtrade Spraykill 250 Herbicide or Imtrade Diquat 200 Non-Residual Herbicide labels	Emerged weeds present (4-6 leaf stage)	250 mL plus Imtrade Spraykill 250 Herbicide or Imtrade Diquat 200 Non-Residual Herbicide at recommended label rates OR 25 to 30 mL/100L plus Imtrade Spraykill 250 Herbicide or ChemAg Diquat 200 Non-Residual Herbicide at recommended label rates	The addition of Imtrade Oxen 240 EC Herbicide to Imtrade Spraykill 250 Herbicide or Imtrade Diquat 200 Non-Residual Herbicide will improve control of Small Flowered Mallow, Evening Primrose and other weeds sensitive to Imtrade Oxen 240 EC Herbicide. Apply as a tank mix with Imtrade Spraykill 250 Herbicide or Imtrade Diquat 200 Non-Residual Herbicide, refer to the appropriate label and follow all label directions.

CROP/SITUATION	WEEDS CONTROLLED	TIME OF APPLICATION	RATE	CRITICAL COMMENTS
Duboisia (After transplanting of young seedlings or after harvest of mature plants.	Refer to Weeds Controlled List (Best results are achieved when applied to moist soil free of weeds)	Weed free soil	4 or 8L/ha	Apply after transplanting or young seedlings or after harvest of mature plants. Use the higher dosage where longer residual control (up to 6 months) is required. "Over-the-top" application will be tolerated.
		Weeds at the 4 to 6 true leaf stage		Best results are achieved when applied to moist soil free of weeds. However, some recently germinated small seedling grasses and broadleaf weeds (4 to 6 true leaves) will be controlled with these rates. Established larger weeds must be eliminated by mechanical means prior to the application of Imtrade Oxen 240 EC Herbicide. Add a non-ionic surfactant (1000g/L) at 250 mL/100L spray solution to enhance activity.
Tropical and Sub-Tropical Fruit (inedible peel) including: Avocado, Cherimoya, Custard Apple, Durian, Feijoa, Guava, Jackfruit, Kiwi Fruit, Longan, Lychee, Mango, Mangosteen, Papaya, Passionfruit, Persimmon, Rambutan, Star Apple	Refer to Weeds Controlled List	Weed free soil	4L/ha	Best results are obtained when applied to moist soil free of weeds.
		Emerged weeds present		If weeds are present, Imtrade Oxen 240 EC Herbicide should be applied as a tank mix with recommended rates of Glyphosate 360g/L, Paraquat 200 g/L or Paraquat 250g/L.

CROP/SITUATION	WEEDS CONTROLLED	TIME OF APPLICATION	RATE	CRITICAL COMMENTS
Brassica Crops Cabbages, Cauliflower, Broccoli	Refer to Weeds Controlled List	Weed free soil	1.5 to 2L/ha	Apply Imtrade Oxen 240 EC Herbicide to prepared ground 4 to 7 days prior to transplanting. If soil is dry, irrigation or rainfall is required prior to transplanting for activation of Imtrade Oxen 240 EC Herbicide. Utilise transplanting techniques which cause minimal soil disturbance. Excessive soil disturbance will lessen herbicidal activity. Use the higher rate in situations where weed pressure is known to be heavy.
Onions (seeded)	Blackberry Nightshade	Hook leaf stage	50-100 mL/ha	DO NOT exceed 100 mL/ha at the hook leaf stage as excessive crop damage may occur.
	Docks, Crowsfoot/ Storksbill, Fumitory Weed (Pink and White Weed), Hogweed (Wireweed), Plantain, Sorrel, Volunteer Potato, Potato Weed, Milk Thistle, Deadnettle, Common Cotula, Groundsel.	1½ to 2½ true leaf stage	350-500 mL/ha	Best results are obtained when the hook leaf stage treatment is followed by a further application of Imtrade Oxen 240 EC Herbicide when onions are at the 1½ to 2½ true leaf stage. Apply Imtrade Oxen 240 EC Herbicide in at least 100L/ha water at pressure below 150kPa. DO NOT exceed 150kPa pressure because excessive crop damage could result. Multiple treatments may be applied provided the total does not exceed 2L/ha in one season. For control of other weeds, use in a program with other registered herbicides. NOTE: Can cause flecking, twisting or stunting of onion plants although such symptoms are normally short-lived. DO NOT apply in conditions of very high humidity. DO NOT apply to weeds or crops which are stressed by prevailing weather conditions, disease or mechanical damage. DO NOT use wetting agents or other adjuvants.

CROP/SITUATION	WEEDS CONTROLLED	TIME OF APPLICATION	RATE	CRITICAL COMMENTS
Tobacco	Refer to Weeds Controlled List	Weed free soil	4L/ha	Use to control weeds along spray lines only. DO NOT apply to tobacco crop. Apply to soil after solid-set irrigation system has been laid out in the field. Imtrade Oxen 240 EC Herbicide should be applied to a moist soil. Where very small weeds (2-3 leaf) emerge prior to spraying, the addition of a wetting agent to the spray mixture is necessary for effective control. Should the weeds be more advanced, the addition of 2L of Imtrade Diquat 200 Non-Residual Herbicide is required. Avoid spray drift.
Coffee (Apply after transplanting coffee plants or under established bushes)	Refer to Weeds Controlled List	Weed free soil	2 to 4L/ha	Apply after transplanting coffee plants or under established bushes. Use the higher rate where longer residual activity (up to 4 months) is required. Best results are achieved when applied to moist soil, free of weeds with follow-up irrigation.
		Weeds at the 2 to 4 true leaf stage		When seedlings are present, apply as a tank mix with Paraquat 200g/L to produce both knockdown and residual control. A non-ionic surfactant (1000g/L) should be used in the spray mixture at 250 mL/100L. Apply as a directed spray to avoid contact with coffee plants. Mature established weeds must be eliminated by mechanical or chemical means prior to application

CROP/SITUATION	WEEDS CONTROLLED	TIME OF APPLICATION	RATE	CRITICAL COMMENTS
Forestry <i>Eucalyptus</i> spp. <i>Pinus</i> spp.	Refer to Weeds Controlled List	Weed free soil (either before or preferably within 4 weeks of transplanting)	3 or 4L/ha	Apply to weed free soil either before or preferably soon after transplanting. Disturbance of the herbicidal barrier on the soil surface by the transplanting process may reduce the length of weed control. Use higher rate for longer residual control. Apply as a directed or over-the-top spray. Imtrade Oxen 240 EC Herbicide can be applied in a tank mix with Simazine to extend the spectrum and length of weed control. Application to foliage that is soft and has not hardened off and/or if temperatures exceed 20°C may increase the likelihood of foliar damage to trees, Eucalypts especially. Imtrade Oxen 240 EC Herbicide has been found to be safe to commonly planted species used in forestry. Sensitivity of less common species should be checked prior to large-scale application.

CROP/SITUATION	WEEDS CONTROLLED	TIME OF APPLICATION	RATE	CRITICAL COMMENTS
Forestry <i>Eucalyptus</i> spp. <i>Pinus</i> spp.	Refer to Weeds Controlled List	Weeds at the 4 to 6 true leaf stage	3 or 4L/ha	Apply as an over-the-top spray to eucalypts and pines. Best results are achieved when applied to moist soil free of weeds. However, some newly germinated small seedling grasses and broadleaf weeds (4 to 6 true leaves) will be controlled at these rates. Addition of a non-ionic surfactant (1000g/L) at recommended rates will improve activity. Established weeds must be eliminated prior to application. Imtrade Oxen 240 EC Herbicide may be tank mixed with Simazine 500g/L or other simazine products to extend the spectrum and length of weed control. Application to foliage that is soft or has not been hardened off and/or if temperatures exceed 20°C may increase the likelihood of foliar damage to trees, eucalypts especially. Imtrade Oxen 240 EC Herbicide has been found to be safe to commonly planted species used in forestry. Sensitivity of less common species should be checked prior to large scale application.
Forestry Trees	Broadleaf weeds and grasses	Weeds at the 4 to 6 true leaf stage	4L/ha or 4 mL/10m ²	For the establishment of trees for approved farm practices such as wind breaks, erosion control, woodlots and forestry plantations. When applying as a post-plant spray, ensure spray is directed to the base of seedlings, or that seedlings are protected. DO NOT apply under hot or windy conditions.

CROP/SITUATION	WEEDS CONTROLLED	TIME OF APPLICATION	RATE	CRITICAL COMMENTS
Pyrethrum – as bare rooted transplants or seedlings	Refer to Weeds Controlled list (except Chickweed)	Pre-plant incorporated into weed free soil worked to a fine tilth	4 to 6L/ha	Apply prior to final soil preparation. The preferred implements for final soil preparation would be either a multiple tyne cultivator or rotary harrows. Use the 6L rate for heavy black clay soils only (as found in the Derwent and Coal River Valleys). Imtrade Oxen 240 EC Herbicide will not provide consistent control of Chickweed.
Pyrethrum – greater than 4 leaves	Blackberry Nightshade (<i>Solanum nigrum</i>), Fumitory (<i>Fumaria</i> spp.), Hogweed (<i>Polygonum aviculare</i>), Sorrel (<i>Rumex acetosella</i>), Volunteer Potato (<i>Solanum tuberosum</i>), Cleavers (<i>Galium aparine</i>), Field Bindweed (<i>Convolvulus arvensis</i>), Groundsel (<i>Senecio vulgaris</i>)	Emerged weeds present	100 to 150 mL/ha	Apply when pyrethrum growth stage is greater than 4 true leaf.

CROP/SITUATION	WEEDS CONTROLLED	TIME OF APPLICATION	RATE	CRITICAL COMMENTS
Pyrethrum – established crops greater than 1 year old	As above plus Sowthistle (<i>Sonchus</i> spp.), Spear Thistle (<i>Cirsium vulgare</i>)	Emerged weeds present	200 mL to 4L/ha	Apply when pyrethrum is greater than 100mm rosettes. Apply rates of greater than 1L/ha ONLY between 1st February and 31 st of March. DO NOT apply later than 25 weeks before harvest.
Prior to Sowing Winter Cereals, Cotton	Annual Weeds Refer to Weeds Controlled list and Glyphosate 360 g/L labels	Actively growing emergent weeds	75 mL/ha plus Glyphosate 360g/L at recommended label rates	Addition of Imtrade Oxen 240 EC Herbicide to Glyphosate 360g/L will improve knockdown and increase the speed at which treated weeds develop visible symptoms of phytotoxicity (compared to results achieved with Glyphosate 360g/L applied alone), and give control to Annual Dead Nettles (<i>Urtica</i> spp.) and Marshmallow (<i>Malva parviflora</i>).

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

GRAZING: DO NOT GRAZE TREATED WEEDS.

HARVEST: NOT REQUIRED WHEN USED AS DIRECTED.

WEEDS CONTROLLED BY IMTRADE OXEN 240 EC HERBICIDE INCLUDE THE FOLLOWING:

Amsinckia (<i>Amsinckia</i> spp.)	Liverseed Grass (<i>Urochloa panicoides</i>)	Starburr (<i>Acanthospermum hispidum</i>)
Barley Grass (<i>Horealum leporinum</i>)	Lovegrass (<i>Eragrostis</i> spp.)	Stinkgrass (<i>Eragrostis cilianensis</i>)
Barnyard Grass (<i>Echinochloa</i> spp.)	Pigeon Grass (<i>Setaria</i> spp.)	Summer Grass (<i>Digitaria</i> spp.)
Blackberry Nightshade (<i>Solanum nigrum</i>)	Pigweed (<i>Portulaca oleracea</i>)	Thornapple (<i>Datura stramonium</i>)
Bladder Ketmia (<i>Hibiscus trionum</i>)	Prickly Lettuce (<i>Lactuca</i> spp.)	White Eye (<i>Richardia brasiliensis</i>)
Burrgrass (<i>Cenchrus australis</i>)	Red Natal Grass (<i>Rhynchelytrum repens</i>)	Wild Mustard (<i>Sisymbrium</i> spp.)
Caltrop (<i>Tribulus terrestris</i>)	Redshank (<i>Amaranthus cruentus</i>)	Wild Radish (<i>Raphanus raphanistrum</i>)
Capeweed (<i>Arctotheca calendula</i>)	Ryegrass (<i>Lolium</i> spp.)	Wireweed (<i>Polygonum aviculare</i>)
Chickweed (<i>Stellaria media</i>)	Sesbania Pea (<i>Sesbania cannabina</i>)	
Crowsfoot (<i>Eleusine indica</i>)	Shepherd's Purse (<i>Capsella bursa-pastoris</i>)	
Deadnettle (<i>Lamium amplexicaule</i>)	Smallflower Mallow (<i>Malva parviflora</i>)	
Fat Hen (<i>Chenopodium album</i>)	Soursob (<i>Oxalis pes-caprae</i>)	
Giant Pigweed (<i>Trianthema portulacastrum</i>)	Sowthistle (<i>Sonchus oleraceus</i>)	

SEEDLINGS (LESS THAN 4 TRUE LEAF) WEEDS CONTROLLED BY IMTRADE OXEN 240 EC HERBICIDE INCLUDE THE FOLLOWING:

Amsinckia (<i>Amsinckia</i> spp.)	Pigweed (<i>Portulaca oleracea</i>)
Bellvine (<i>Ipomoea</i> spp.)	Potato Weed (<i>Galinsoga parviflora</i>)
Capeweed (<i>Arctotheca calendula</i>)	Redshank (<i>Amaranthus cruentus</i>)
Common Cotula (<i>Cotula australis</i>)	Shepherd's Purse (<i>Capsella bursa-pastoris</i>)
Crowsfoot Grass (<i>Eleusine indica</i>)	Sowthistle (<i>Sonchus oleraceus</i>)
Deadnettle (<i>Lamium amplexicaule</i>)	Stinging Nettle (<i>Urtica urens</i>)
Groundsel (<i>Senecio vulgaris</i>)	Stinkgrass (<i>Eragrostis cilianensis</i>)
Liverseed Grass (<i>Urochloa panicoides</i>)	Wild Radish (<i>Raphanus raphanistrum</i>)
Marshmallow (<i>Malva parviflora</i>)	

GENERAL INSTRUCTIONS

Imtrade Oxen 240 EC Herbicide is a selective herbicide for the control of certain annual grass and broadleaf weeds in dormant Apples, Grapes, Peaches, Pears, Plums, Apricots, Almonds, Macadamia, Duboisia, Tobacco, Coffee, Pyrethrum, Winter Cereals, Tropical/Sub-tropical crops, Brassicas, Onions, Eucalyptus and Pinus spp. And other forestry trees, applied wither to weed free soil or to seedling weeds up to the 4 to 6 true-leaf stage.

Imtrade Oxen 240 EC Herbicide applied to well prepared, weed free soil should not be disturbed or incorporated after application. Weed control for up to 6 months can be expected with high label rates, but spot treatment of escape weeds or perennial grasses may be necessary with knockdown herbicides. When Imtrade Oxen 240 EC Herbicide is applied to seedling weeds at the 4 to 6 leaf stage, a non-ionic surfactant should be added at recommended rates to improve activity, where suggested in the CRITICAL COMMENTS.

Imtrade Oxen 240 can also be used at low rates as a "spike" to improve the weed spectrum of knockdown herbicides such as Glyphosate and Paraquat or Diquat/Paraquat mixtures.

RESISTANT WEEDS WARNING

GROUP	14	HERBICIDE
--------------	-----------	------------------

Imtrade Oxen 240 EC Herbicide is a member of the Diphenyl ether group of herbicides. The mode of action of Imtrade Oxen 240 EC Herbicide is to inhibit protoporphyrinogen oxidase. For weed resistance management Imtrade Oxen 240 EC Herbicide is a Group **14** Herbicide. Some naturally occurring weed biotypes resistant to Imtrade Oxen 240 EC Herbicide and other Group **14** Herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if Imtrade Oxen 240 EC Herbicide or other Group **14** herbicides are used repeatedly. These resistant weeds will not be controlled by Imtrade Oxen 240 EC Herbicide or other Group **14** Herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Imtrade Australia Pty Ltd accepts no liability for any losses that may result from the failure of Imtrade Oxen 240 EC Herbicide to control resistant weeds.

TIMING

Residual Control

For optimum residual weed control, Imtrade Oxen 240 EC Herbicide should be applied to the soil surface prior to weed emergence after all other agricultural operations, such as mechanical cultivation and re-shaping of irrigation furrows, have been completed. The area should be left undisturbed during the period for which weed control is desired. When applied to seedling weeds, they should be young and actively growing. Weed control for up to 6 months is expected by spot treatment, with knockdown herbicides, for escape weeds and perennial grasses may be necessary.

Post-emergence Weed Control

For optimum post-emergence weed control, Imtrade Oxen 240 EC Herbicide + Glyphosate tank mixes should be applied to small seedling weeds up to 4 to 6 true-leaf stage. Use of a non-ionic surfactant (1000g/L) is recommended to improve activity. Weeds should be actively growing and free from environmental stress (drought, cold, insect attack, nutrient deficiency). Cultivation after treatment and prior to or at planting is beneficial for final fallow weed control.

MIXING

When using Imtrade Oxen 240 EC Herbicide alone, fill the spray tank at least one-third full with clean water, add the recommended amount of Imtrade Oxen 240 EC Herbicide while the pump and agitator are running, then complete filling the spray tank.

A non-ionic surfactant (1000g/L), if required by label directions, should be added near the end of the filling process to minimise foaming. When tank mixing with Oryzalin 500g/L, add to one-third filled tank and then add the Imtrade Oxen 240 EC Herbicide during the filling operation. Maintain agitation during mixing and until the spraying is completed.

When mixing with Glyphosate 360g/L, Imtrade Spraykill 250 Herbicide or Imtrade Diquat 200 Non-Residual Herbicide, add these after Imtrade Oxen 240 EC Herbicide during the filling operation. When using Imtrade oxen 240 EC Herbicide in combination with Glyphosate 360g/L or Oryzalin 500g/L, refer to the appropriate label and read and follow all label directions.

Imtrade Oxen 240 EC Herbicide + Glyphosate tank mixes

Ensure thorough agitation when mixing, filling the spray tank and during application irrespective of glyphosate formulations used. Follow recommended order and directions for tank mixing Imtrade Oxen 240 EC Herbicide and glyphosate. Use all spray mix immediately after preparation.

DO NOT tank mix Imtrade Oxen 240 EC Herbicide and Glyphosate without agitation.

DO NOT allow mix to stand unagitated.

DO NOT store Imtrade Oxen 240 EC Herbicide and Glyphosate tank mixes.

DO NOT mix other agrochemical products with Imtrade Oxen 240 EC Herbicide and Glyphosate tank mixes.

APPLICATION

Spray equipment should be calibrated before use. Imtrade Oxen 240 EC Herbicide should be applied uniformly as a directed treatment to the base of tree and vine crops using flat fan or off-centre nozzles. Complete coverage of seedling weeds is required for maximum knockdown effect.

A water volume of 250 to 500L/ha is recommended for treatments on bare soil.

A spray volume of 100 to 1350L/ha is recommended where seedling weeds (4 to 6 leaf) are present. Ensure both weed foliage and the soil surface are sprayed. Use the higher volumes where weed density is high.

Tank mixtures of 75 mL/ha of Imtrade Oxen 240 EC Herbicide with Glyphosate 360g/L herbicides should be applied in 30 to 200L spray volume/ha. For maximum residual control, Imtrade Oxen 240 EC Herbicide should **NOT** be incorporated or disturbed after application.

CROP SAFETY

Imtrade Oxen 240 EC Herbicide may be applied as directed around dormant peach, plum, apricot, almond, apple and pear trees and grape vines of all ages when applied at rates of less than 1.0L/ha. When applied at 3.0L/ha and above, the trees and grape vines should be at least 3 years of age. **DO NOT** apply Imtrade Oxen 240 EC Herbicide once bud swell has occurred when using rates greater than 1.0L/ha. Duboisia seedlings and mature plants will tolerate "over-the-top" applications of Imtrade Oxen 240 EC Herbicide. When using the 75 mL/ha rate, Imtrade Oxen 240 EC Herbicide may be applied as directed prior to sowing winter cereals.

SAFE SOWING INTERVALS

Imtrade Oxen 240 EC Herbicide at up to 75 mL/ha may be safely applied 1 day prior to planting broadacre crops such as Cereals (Wheat, Barley, Oats, Triticale), Canola, Pulses (Lupins, Faba Beans, Field Peas) and undersown pastures (Lucerne, Clover, Medics, Ryegrass, Phalaris, Cocksfoot) and 7 days minimum prior to planting Cotton or Soybeans, provided minimum tillage planting equipment is used with minimal soil disturbance. Inversion, mixing of surface soil with that in the planting zone or covering seed with treated soil may result in injury to emerging crop seedlings.

Avoid covering the seed with soil treated with Imtrade Oxen 240 EC Herbicide during the planting operation to minimize crop injury.

Imtrade Oxen 240 EC Herbicide has residual soil activity, especially when applied at rates greater than 75 mL/ha and on small-seeded horticultural crops. Plantback intervals in the following table for horticultural crops must be observed if more than 75 mL/ha Imtrade Oxen 240 EC Herbicide has been applied.

Safe Sowing Intervals for Horticultural Crops (Days)

Imtrade Oxen 240 EC Herbicide	Up to 75 mL/ha	1L/ha
Beans	7	60
Brassicas	14	90
Capsicums	14	90
Carrots	14	90
Cucurbits	14	60
Lettuce	14	90
Onions	21	180
Potatoes	7	60
Tomatoes	14	60

COMPATIBILITY

Imtrade Oxen 240 EC Herbicide is compatible with Glyphosate 360g/L, Oryzalin 500g/L, Imtrade Paraquat 250 Herbicide, Imtrade Diquat 200 Non-Residual Herbicide, Touchdown*, Imtrade Spraykill 250 Herbicide and Glufosinate-Ammonium 200g/L.

RE-ENTRY PERIOD

DO NOT allow entry into treated areas until the spray has dried. If prior entry is necessary, wear chemical resistant footwear.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

PROTECTION OF LIVESTOCK

Use with care when applying in areas frequented by stock.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

This product is highly toxic to wildlife and fish. DO NOT contaminate streams, rivers or waterways with the chemical or used containers. Use care when applying in areas frequented by wildlife or adjacent to any body of water.

STORAGE AND DISPOSAL (1L, 5L, 10L, 20L, 60L & 200L only)

Store in the closed original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemical on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Empty containers and product should **not** be burnt.

Envirodrum - Micro Matic Valve (110L)

Store the original sealed Envirodrum in a cool, well ventilated area. **DO NOT** store for prolonged periods in direct sunlight. **DO NOT** tamper with the Micro Matic valve or the security seal. **DO NOT** contaminate the Envirodrum with water or any other foreign matter. After each use of the product please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase.

SAFETY DIRECTIONS

Will damage eyes. Will irritate the skin. Avoid contact with eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. **DO NOT** inhale vapour or spray mist. When opening the container and preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length butyl rubber gloves and face shield or goggles. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (*Phone Australia 13 11 26; New Zealand 0800 764 766*). If swallowed **DO NOT** induce vomiting. Give a glass of water. If in eyes wash out immediately with water.

Additional GHS Hazard & Precautionary Statements

●May be harmful if swallowed and enters airways; ●May cause respiratory irritation; ●May damage the unborn child; ●Very toxic to aquatic life with long-lasting effects; ●Avoid breathing fumes; ●**DO NOT** get on clothing; ●Wash contacted areas thoroughly after handling; ●Use only outdoors or in a well-ventilated area; ●Call a POISON CENTRE or doctor if you feel unwell; ●**IF INHALED:** Remove victim to fresh air and keep at rest in a position comfortable for breathing; ●If exposed or concerned: Get medical advice; ●If skin irritation occurs: Get medical advice; ●If eye irritation persists: Get medical advice; ●Store locked up.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from your supplier or Imtrade Australia Pty Ltd website at www.imtrade.com.au

CONDITIONS OF SALE

Imtrade Australia Pty Ltd warrants that this product conforms to the chemical description on the label. The use of Imtrade Oxen 240 EC Herbicide being beyond the control of the manufacturer, Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Imtrade Australia Pty Ltd has any authority to add to or alter these conditions.

*Other Registered Trademarks

UN 3082	ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S. (CONTAINS OXYFLUORFEN)
IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE	
PG III	HAZCHEM *3Z



APVMA Approval No: 56798/106771