

#### STORAGE AND DISPOSAL (5L, 10L, 20L, 200L only)

Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. **DO NOT** store near food, fertilizers or seed. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Empty containers and product should **NOT** be burnt.

#### Envirodrum - Micro Matic Valve (110L only)

Store the closed, original sealed Envirodrum in a well-ventilated area, as cool as possible and locked away from children, animals, food, foodstuffs, seed and fertilisers. **DO NOT** store for prolonged periods in direct sunlight. **DO NOT** tamper with the Micro Matic valve or the security seal. **DO NOT** contaminate the Envirodrum with water or any other foreign matter. After each use of the product, please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase.

#### Refillable containers (1000L only)

Store in the closed, original container in a well-ventilated area, as cool as possible and locked away from children, animals, food, foodstuffs, seed and fertilisers. **DO NOT** store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

#### SMALL SPILL MANAGEMENT

Wear protective equipment (See Safety Directions). Apply absorbent material such as earth, sand, clay granules or cat litter to the spill. Sweep up material for disposal when absorption is completed and contain in a refuse vessel for disposal. (See Storage and Disposal section). If necessary, wash the spill area with an alkali detergent and water and absorb as above the wash liquid for disposal.

#### SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When preparing the spray, wear cotton overalls buttoned to the neck and wrists, a washable hat, elbow-length PVC gloves and a face shield or goggles. If the product is in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

# POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

## IMTRADE PICKER HERBICIDE

ACTIVE CONSTITUENTS: 300 g/L TRICLOPYR present as butoxyethyl ester  
100 g/L PICLORAM present as hexyloxypropylamine salt

**GROUP 4 HERBICIDE**

For control of a range of environmental and noxious woody and herbaceous weeds as specified in the Directions for Use table.

**Important: Read the attached Leaflet before use.**

GROUP  
4

FORM  
EC

CONTENTS  
1L-1000L



Batch No.  
DOM:

#### FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre (Phone Australia 13 11 26).

Refer to attached Leaflet for additional GHS  
Hazard & Precautionary Statements

#### SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet which can be obtained from your supplier or Imtrade Australia Pty Ltd website at [www.imtrade.com.au](http://www.imtrade.com.au)

#### CONDITIONS OF SALE

Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Imtrade Australia Pty Ltd has any authority to add to or alter these conditions.

In a Transport Emergency Dial 000  
Police or Fire Brigade

APVMA Approval No: 56453/54458

 **IMTRADE**  
CropScience

17 Ocean St, Kwinana Beach WA 6167  
08 9419 0333 | [www.imtrade.com.au](http://www.imtrade.com.au)

**POISON**

**KEEP OUT OF REACH OF CHILDREN**

**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**



# **IMTRADE PICKER HERBICIDE**

**ACTIVE CONSTITUENT: 300 g/L TRICLOPYR**  
**present as butoxyethyl ester**  
**100 g/L PICLORAM**  
**present as hexyloxypropylamine salt**

**GROUP 4 HERBICIDE**

For control of a range of environmental and  
noxious woody and herbaceous weeds  
as specified in the DIRECTIONS FOR USE table

**IMPORTANT: READ THIS LEAFLET THOROUGHLY  
BEFORE OPENING OR USING THIS PRODUCT**

**SHAKE WELL BEFORE USE**

**APVMA Approval No: 56453/54458**

Imtrade Australia Pty Ltd  
ABN 13 090 151 134  
17 Ocean Street, Kwinana, Western Australia 6167  
Tel: (08) 9419 0333 Fax: (08) 9419 7516  
Web: [www.imtrade.com.au](http://www.imtrade.com.au)

## DIRECTIONS FOR USE

### RESTRAINTS:

**DO NOT** apply to weeds which have been stressed (not actively growing) due to prolonged periods of extreme cold, moisture stress (water-logged or drought affected), poor nutrition, presence of disease, damage or previous herbicide treatment, as reduced levels of control may result.

**DO NOT** spray if rain is likely within one hour or if foliage is wet from rain or dew.

**DO NOT** burn off, cut or clear blackberry or other woody weeds for at least six months after spraying.

**DO NOT** apply by aerial application in wind in excess of 15km/hr and air temperatures above 35°C. In areas prone to flooding treatment should commence after any flooding as such areas flooded within 9 months following application may have reduced results.

## WOODY WEED SITUATIONS

**Table A: High Volume Spraying**

See **GENERAL INSTRUCTIONS – APPLICATION** section for application method details.

<b>AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY.</b>				
<b>WEEDS CONTROLLED</b>	<b>WEED GROWTH STAGE</b>	<b>STATE</b>	<b>RATE/ 100L water</b>	<b>CRITICAL COMMENTS</b>
African Boxthorn	Less than 2m tall	All States	500 mL	Apply when bushes have good leaf cover, growth and no leaf fall.
<i>Angophora</i> spp.	1 to 3m tall		350 mL	
Australian Blackthorn	Less than 2m tall		500 mL	Apply from late spring to early Autumn.
<i>Banksia</i> spp.	1 to 3m tall		350 mL	
Biddy Bush, (Chinese Shrub), (Sifton Bush)	Autumn when actively growing	ACT and NSW only	500 mL	Add a 100% concentrate non-ionic surfactant (e.g. ChemAg Wetter 1000) at 125 mL /100L of water for best results.
Blackberry in association with: Docks, Ragwort, Smartweed, Thistles	Late Spring to Autumn	All States except NT	350 or 500 mL	<b>Use the higher rate on plants that have been damaged by grazing stock or insects and on known difficult to kill Blackberry.</b> Where herbicides other than Group 4 Herbicides have been used, allow two seasons re-growth to occur before respraying with Imtrade Picker Herbicide
Blue Heliotrope	Flowering	NSW and Qld only	500 mL	Apply in a minimum spray volume of 1250L/ha
Brooms: Cape, English, Flax Leaf, Montpellier	Spring to mid-Summer prior to pod formation	All States except NT	250 mL	Apply as a thorough foliage spray.
	Autumn to Winter		350 mL	

**Table A: High Volume Spraying (Continued)**

See GENERAL INSTRUCTIONS – APPLICATION section for application method details.

<b>AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY.</b>				
<b>WEEDS CONTROLLED</b>	<b>WEED GROWTH STAGE</b>	<b>STATE</b>	<b>RATE/ 100L water</b>	<b>CRITICAL COMMENTS</b>
Camphor Laurel	Less than 2m tall	All States	350 mL	Apply as a thorough foliage spray.
	Above 2m tall		500 mL	
<i>Casuarina</i> spp.	1 to 3m tall		350 mL	
Chinese Apple	Less than 2m tall	Qld and WA only	350 mL	Add a 100% concentrate non-ionic surfactant (e.g. ChemAg Wetter 1000) at 100 mL /100L of water for best results.
Cockspur Thorn	Spring to Autumn	ACT, NSW and Qld only	350 mL	Apply as a thorough foliage spray.
Common Sensitive Plant	Any time when actively growing	NT, Qld and WA only	200 mL	To avoid leaves closing during application, spray plants while moving forward.  Add a 100% concentrate non-ionic surfactant (e.g. ChemAg Wetter 1000) at 100 mL /100L of water for best results.
Crofton Weed	Spring to Autumn	ACT, NSW and Qld only	350 mL	Apply as a foliar spray.
Eastern Cotton Bush ( <i>Mairana microphylla</i> )	Spring to Autumn	NSW and Qld only	500 mL	Add Uptake® Spraying Oil at 500 mL/100L of water. Some bushes may require a follow-up spray to control regrowth.
<i>Eucalyptus</i> spp.	Seedling and regrowth from small lignotubers 1 to 3m tall	All States	350 or 500 mL	<b>Apply the high rate where difficult to control species of Eucalyptus regrowth is present.</b>  Addition of an adjuvant may improve results – contact Imtrade Australia Pty Ltd for details.
Galenia	Fresh growth in Spring to Summer	NSW only	500 mL	Use 2000L of spray mixture/ha.
Giant Bramble	Spring to Autumn	NT, Qld and WA only	500 mL	Penetration of thick clumps may be difficult and re-spraying may be necessary. Add a 100% concentrate non-ionic surfactant (e.g. ChemAg Wetter 1000) at 100 mL/100L water for best results.

**Table A: High Volume Spraying (Continued)**

See GENERAL INSTRUCTIONS – APPLICATION section for application method details.

<b>AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY.</b>				
<b>WEEDS CONTROLLED</b>	<b>WEED GROWTH STAGE</b>	<b>STATE</b>	<b>RATE/ 100L water</b>	<b>CRITICAL COMMENTS</b>
Gorse	1 to 1.5m tall	All States except NT	250 mL	Spring and Summer treatment only.
	Over 1.5m tall or Autumn treatment		350 mL	Add a 100% concentrate non-ionic surfactant (e.g. ChemAg Wetter 1000) at 100 mL/100L of water for best results.
	Winter treatment	All States except NT	500 mL	Brownout may not be complete until summer. Add a 100% concentrate nonionic surfactant (e.g. ChemAg Wetter 1000) at 100 mL/100L of water for best results.
Groundsel Bush ( <i>Baccharis halimifolia</i> )	1 to 1.5m tall in Spring to Summer		250 mL	Apply as a thorough foliage spray.
	Over 1.5m tall or Autumn treatment	350 mL		
Green Cestrum	Late spring to early Autumn	ACT, NSW and Qld only	500 mL	One application may give satisfactory control. Any subsequent re-growth and seedlings must be re-sprayed at approximately 1 metre high.
Hawthorn	Less than 2m tall	All States		Apply from late spring to early Autumn.
Horehound	Pre-flowering	All States	350 mL	Apply as a thorough foliar spray.
Japanese Sunflower		NSW and Qld only		
Lantana ( <i>Lantana camara</i> ), ( <i>Lantana montevidensis</i> )	Up to 1m tall in Summer to Autumn	All States	350 mL	Add one of the following adjuvants, when using 350 mL rate: Uptake® Spraying Oil @ 0.5% v/v. Pulse® Penetrant @ 0.1% v/v.
	1 to 2m tall in Summer to Autumn		500 or 750 mL	Thoroughly wet foliage, stems and soil around base of the plants. Use higher rate on known harder to kill varieties.
Lion Tail ( <i>Leonatis nepetifolia</i> )	Pre-flowering	Qld only	200 mL	Apply as a thorough foliar spray. Add a 100% concentrate non-ionic surfactant (e.g. ChemAg Wetter 1000) at 100 mL/100L of water for best results.

**Table A: High Volume Spraying (Continued)**

See GENERAL INSTRUCTIONS – APPLICATION section for application method details.

<b>AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY.</b>				
<b>WEEDS CONTROLLED</b>	<b>WEED GROWTH STAGE</b>	<b>STATE</b>	<b>RATE/100L water</b>	<b>CRITICAL COMMENTS</b>
Limebush	Any time of year with good leaf cover and soil moisture	NT and Qld only	350 mL	Penetration of thick clumps may be difficult and re-spraying may be required. Addition of an adjuvant may improve results contact Imtrade Australia Pty Ltd for details.
Manuka	At flowering	Vic only	500 mL	For optimum results, add Pulse® Penetrant at 200 mL/100L of spray. Thoroughly wet foliage, stems and soil around the base of the plants.
Mesquite ( <i>Prosopis</i> spp.)	Seedling, full leaf and flowering before podding	NSW, NT, Qld and WA only	350 mL	<b>DO NOT</b> spray plants bearing pods. Add a 100% concentrate non-ionic surfactant (e.g. ChemAg Wetter 1000) at 100 mL/100L of water for best results
<i>Prosopis velutina</i>		Qld only	670 mL	
Mistflower	Spring to Autumn	ACT, NSW and Qld only	670 mL	Apply as a thorough foliar spray
Mother-of-Millions	Flowering	NSW and Qld only	500 mL	Add a 100% concentrate non-ionic surfactant (e.g. ChemAg Wetter 1000) at 100 mL/100L of water.
Paddy's Lucerne	Active growth	NSW only		Plants that have been continually slashed or grazed over many seasons may be difficult to control and re-growth may occur.
Parkinsonia	Under 2m tall	NT, Qld and WA only	350 mL	Add Uptake® Spraying Oil @ 500 mL/100L water. Avoid spraying under dry conditions when plants are stressed and bearing pods. Thoroughly wet foliage.
Prickly Pear (common Smooth Tree Pear)	Active phyllode growth	All States	500 mL	Apply as a thorough foliage spray. Re-growth may occur, so a follow-up application may be necessary
Rubber Vine (Not infected with rust)	Up to 1.5m tall at flowering	NT, Qld and WA only	350 mL	Spray all leaves and stems just to the point of runoff and thoroughly spray the base of the plant. With larger, more dense stands, re-growth may occur. Subsequent control of any re-growth should be done by basal bark spraying.
	Dense stands greater than 1.5m tall at flowering		500 mL	

**Table A: High Volume Spraying (Continued)**

See GENERAL INSTRUCTIONS – APPLICATION section for application method details.

<b>AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY.</b>				
<b>WEEDS CONTROLLED</b>	<b>WEED GROWTH STAGE</b>	<b>STATE</b>	<b>RATE/100L water</b>	<b>CRITICAL COMMENTS</b>
Siam Weed	Active growth	Qld and WA only	350 mL	Add a 100% concentrate non-ionic surfactant (e.g. ChemAg Wetter 1000) at 100 mL/100L of water for best results.
Sticklepod	Up to flowering	NT, Qld and WA only	200 mL	<b>DO NOT</b> apply to podding plants. Add a 100% concentrate non-ionic surfactant (e.g. ChemAg Wetter 1000) at 100 mL/100L of water for best results.
St John's Wort	From flowering to early seed set	All States	500 mL	Late spring to early Summer.
Sweet Briar	Up to 1.5m tall	All States except NT	350 mL	Add metsulfuron-methyl at 10g/100L water to obtain more reliable results with the lower rate of Imtrade Picker Herbicide.
			500 mL	Full leaf to ripe fruit prior to leaf fall. Thorough wetting including the crown is recommended.
Tobacco Weed	Actively growing plants	NT, Qld and WA only	300 mL	Add a 100% concentrate non-ionic surfactant (e.g. ChemAg Wetter 1000) at 100 mL/100L of water for best results.
Wattle ( <i>Acacia</i> spp.) (except Corkwood Wattle)	1 to 3m tall	All States	350 mL	
Wild Rosemary ( <i>Cassinia laevis</i> )	Active growth, 0.5 to 1.0m tall	Qld only	350 to 500 mL	Use lower rate on seedlings 0.5m tall. Apply as a thorough foliar spray.
Wild Tobacco Tree	Spring to Autumn up to 2m tall	ACT, NSW and Qld only	350 mL	

**Table B: Aerial Application**

See GENERAL INSTRUCTIONS – APPLICATION section for application method details.

<b>AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY.</b>				
<b>WEEDS CONTROLLED</b>	<b>WEED GROWTH STAGE</b>	<b>STATE</b>	<b>RATE/ha</b>	<b>CRITICAL COMMENTS</b>
Blackberry	Summer to Autumn	NSW, Qld, SA, Vic and WA only	10L	Where herbicides other than Group 4 Herbicides have been used, allow two seasons re-growth to occur before respraying with Imtrade Picker Herbicide. <b>WARNING:</b> Eucalyptus species up to 4m may be killed if sprayed during the treatment. Mature trees, which are 15 to 20m tall may be partially defoliated but are likely to recover.
Gorse		Tas only		Helicopter application only.
Cockspur Thorn, Crofton Weed, Lantana, Mistflower	Late Autumn	NT and Qld only (Helicopter only)	1.5L plus 7.5L 2,4-D Amine (500g/L formulation)	Spray with calibrated equipment using the half overlap opposite pass technique applying a minimum spray volume of 150L/ha. <i>Follow-up re-spraying will be required.</i>
Lantana			10L	Helicopter application only.
Rubber Vine (Not infected with rust)	When flowering	NT and Qld only (Helicopter only)	3L to 5L	Use rates will depend upon the density and height of the rubber vine stand. The higher rate should be used on dense stands, however, complete coverage and penetration may be difficult. <i>Follow-up re-spraying will be required.</i> Any re-growth should be sprayed with a suitable Basal Bark herbicide.
St John's Wort	Flowering to early seed set (Nov-Jan)	NSW only	4L	Helicopter application only. <i>Follow-up spraying will be required in the following season.</i>

<b>AGRICULTURAL NON-CROP AREAS ON FLOOD PLAINS.</b>				
<b>WEEDS CONTROLLED</b>	<b>WEED GROWTH STAGE</b>	<b>STATE</b>	<b>RATE/ha</b>	<b>CRITICAL COMMENTS</b>
Parkinsonia	Seedlings 1-2m tall, or 12-24 months old	NT and Qld only (Helicopter only)	3L	Add Uptake® Spraying Oil at 1L/ha.



**Table C: Controlled Droplet Application (C.D.A.)**

See GENERAL INSTRUCTIONS APPLICATION section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY.				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE/ha	CRITICAL COMMENTS
Blackberry in association with: Docks, Ragwort, St John's Wort, Thistles	Summer to Autumn	All States except NT	Apply undiluted	One application may give satisfactory control but subsequent re-growth and seedlings should be re-sprayed after hardening of. Where herbicides other than Group 4 Herbicides have been used, allow two seasons re-growth to occur before re-spraying with Imtrade Picker Herbicide.

**Table D: Low Volume High Concentrate Application Techniques (Gas Powered Gun, Sprinkler Sprayer)**

See GENERAL INSTRUCTIONS APPLICATION section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY.				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE/ha	CRITICAL COMMENTS
Blackberry	Late Spring to Autumn	ACT, NSW, Qld, SA, Tas and WA only	335 mL	Apply to actively growing bushes, which can be sprayed on all sides. For larger bushes, the high volume application technique is recommended.
Camphor Laurel, Cockspur Thorn, Crofton Weed.	Less than 1.5m high	ACT, NSW and Qld only	500 mL	
<i>Eucalyptus</i> species	Seedlings up to 2m tall	All States	335 mL	Apply to actively growing bushes, which can be sprayed on all sides. For larger bushes, the high volume application technique is recommended.
Mistflower	Less than 1.5m high	ACT, NSW and Qld only	500 mL	
Sweet Briar	1.5m tall, full leaf to ripe fruit	NSW only		
St John's Wort	During flowering to early seed set	NSW, Tas and Vic only		<b>Gas Powered Gun Only:</b> One application should provide control. Minor re-growth and seedlings may be retarded the following Summer.
Wild Tobacco Tree	Less than 1.5m high	ACT, NSW and Qld only		Apply to actively growing bushes, which can be sprayed on all sides. For larger bushes, the high volume application technique is recommended.

**Table E: Boom Application**

See GENERAL INSTRUCTIONS APPLICATION section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY.				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE/ha	CRITICAL COMMENTS
Galenia	Fresh growth during Spring to Summer	NSW only	5L	Rough mine sites will require adequate spray equipment such as boomless nozzles for effective coverage.
Sicklepod	Up to flowering	NT and Qld only	3L	<b>DO NOT</b> apply to podding plants. Add a 100% concentrate non-ionic surfactant (e.g. ChemAg Wetter 1000) at 100 mL/100L of water.
St John's Wort	Flowering to early seed set (Nov-Jan)	NSW only	2 to 4L	Use the higher rate on dense infestation and when longer residual control is required. Follow-up respraying will be required in the following season.

**2. FALLOW SITUATIONS****Table A: Boom Application**

See GENERAL INSTRUCTIONS APPLICATION section for application method details.

FALLOW				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE/100L water	CRITICAL COMMENTS
Blackberry Nightshade <b>Suppression only</b>	10 to 25cm tall, prior to flowering	NSW and Qld only	200 to 400mL + 1.2L Glyphosate 450 Herbicide + adjuvant	<b>FOR USE BY GROUND EQUIPMENT ONLY.</b> Plants must be actively growing.
Camel Melon, Prickly Paddy Melon, Cucumber Melon ( <i>Cucumis melo</i> )	From 2-leaf to 50cm diameter			Use the lower rate on the smaller weeds, as specified in the weed growth stage (or up to 5cm diameter for <i>Polymeria pusilla</i> ).
Common Sowthistle	From 8-leaf to flowering			Refer to Glyphosate 450 label for use of adjuvant.
Cowvine	From 2 to 5 leaf up to 15cm diameter, prior to flowering			<b>DO NOT</b> plant susceptible crops for up to nine months after application, as specified in General Instructions – Minimum Re-cropping Periods Black cracking Clay soils, NNSW and Qld.
Lucerne (established)	Active growth, 15 to 25cm high, during spring	NSW and Qld only	300 to 500 mL + 1.2L Glyphosate 450 Herbicide + adjuvant	Dry conditions after application will increase the re-cropping interval.

## 2. FALLOW SITUATIONS (Continued)

**Table A: Boom Application**

See **GENERAL INSTRUCTIONS APPLICATION** section for application method details.

FALLOW				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE/100L water	CRITICAL COMMENTS
<i>Ploymeria pusilla</i>	2 to 12-leaf up to 20cm diameter prior to flowering	NSW and Qld only	200 to 400mL + 1.2L Glyphosate 450 Herbicide + adjuvant	Dry conditions after application will increase the re-cropping interval.

**Table B: Blanket Wiper Application**

See **GENERAL INSTRUCTIONS APPLICATION** section for application method details.

FALLOW				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE/100L water	CRITICAL COMMENTS
Biterbark ( <i>Alstonia constricta</i> )	From Summer to end of Autumn	Qld	1+4 (1 part Imtrade Picker Herbicide to 4 parts water) 2% solution for spot spray (e.g. 100 mL Imtrade Picker Herbicide in 5L water)	<p>For use with blanket wipers only. For best results apply in autumn to tall (&gt;60cm) plants using two opposite directional passes (up and back)</p> <p>Follow up "missed" plants with a spot spray application. These will be obvious after 6 weeks.</p> <p>Blanket wiper applications can be made in Summer when plants are smaller but follow up spot spraying may be necessary.</p> <p><b>DO NOT</b> disturb (cultivate) the treated patches for at least 3 months. Best long-term control is achieved when patches are left undisturbed for as long as possible after treatment (at least 6 months).</p> <p><b>Spot spraying "missed" plants:</b> Thoroughly wet all stems and leaves without producing any solution run-off. Avoid any spray reaching the soil surface.</p>

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

## IN TASMANIA FOR BLACKBERRY

**DO NOT** treat bushes carrying mature or near mature fruit.

## FOR NATIVE VEGETATION

Use of Imtrade Picker Herbicide on native vegetation must be done in accordance with STATE and/or LOCAL legislation.

## WITHHOLDING PERIOD

**NOT REQUIRED WHEN USED AS DIRECTED.**

## GENERAL INSTRUCTIONS

**MINIMUM RECROPPING PERIODS – Black Cracking Clay Soils, NNSW & Qld.**

### A. Boom Application

Plant-back periods for crops following the application of Imtrade Picker Herbicide for rates up to 600 mL/ha.				
RATE mL/HA	200	300	400	600
CROP	Months			
Wheat	2	2	4	4
Barley	2	2	4	4
Canola	2	4	4	4
Faba Bean	4	4	6	6
Chickpea	4	6	6	6
Lucerne	6	9	9	9

These plant-back periods are based on normal rainfall pattern. During drought conditions (or when the rainfall is less than 100mm for a period of 4 months or greater) the plant-back period may be significantly longer.

### B. Blanket Wiper Application

**Plant-back periods for crops following blanket wiper application.**

CROP	Months
Broadleaf crops	18
Lucerne	6 This will allow any potential soil residues to dissipate, if any, and allow effective control.

**Note:** Before using Imtrade Picker Herbicide in tank mixes with other herbicides, check the plant-back information on all product labels. The most residual product, i.e. the product with the longest plant-back period, will determine the time between spraying and planting.

## COMPATIBILITY

Follow any regional restrictions and all directions and restrictions on the label, of any chemical mixed with Imtrade Picker Herbicide (e.g. 2,4-D Amine).

Imtrade Picker Herbicide is compatible with the following herbicides: 2,4-D Amine 500, Abound® 300, Dichloprop, Metsulfuron-methyl, Glyphosate 360 & 450, Glyphosate Ammonium and Fluroxypyr.

Imtrade Picker Herbicide is compatible with the following adjuvants, as per Directions for Use: ChemAg Wetter 1000, Pulse® and Uptake® Spraying Oil.

## MIXING

Mix only with water. Half fill the spray unit with water, and add the required amount of Imtrade Picker Herbicide. Add the remaining water with the agitator running. If required, then add spray oils or wetters (surfactants). Maintain mechanical or bypass agitation in the spray tank during spraying. Only mix sufficient solution for immediate daily use and avoid storing.

## APPLICATION

### 1. WOODY WEED SITUATIONS

**Weeds need to be actively growing for herbicides to have optimum effect. Delay treatment until all re-growth has had time to grow to approximately 1 metre in situations which have been bulldozed, slashed, burnt, ploughed or areas having a previous chemical treatment.**

#### A. High Volume Spraying

Thorough coverage of foliage to the point of run-off is essential, however, avoid excess spraying which is wasteful of chemical.

##### Hand Gun

- Apply the recommended mix to give full coverage of leaves and stems through a No. 6 to 8 tip at 700 to 1500 kPa (400 to 500 kPa for St John's Wort).
- A spray volume of 3000 to 4000L per infested hectare of 1 to 2 metre high Blackberry (30 to 40L/100m<sup>2</sup>) should be used.
- Use 2000L of spray mixture/ha for Galenia infestation (i.e. 20L/100m<sup>2</sup> infested area).

##### Knapsack

- Apply the recommended spray mix to give full coverage of leaves and stems. The final volume of application should be similar to the hand gun.
- A spray volume of 3 to 4L/10m<sup>2</sup> infested area should be used.
- A spray volume of 2L/10m<sup>2</sup> should be used for an area infested with Galenia.

#### B. Aerial Application

Apply 200L of water/ha using an aircraft to apply 100L per pass on a double overlap pattern using nozzle configuration to produce droplets 250 to 350 micron diameter. The potential for damage from drift can be greatly reduced by avoiding unsuitable spraying conditions and using spray pressure and nozzles to minimise the production of small droplets.

**DO NOT** spray when wind exceeds 15km/hr and/or air temperature reaches 35°C.

### **C. Controlled Droplet Application (C.D.A.)**

Results similar to high volume spraying can be obtained using Micron Herbi or similar equipment. Select a nozzle to give a flow rate of 2 mL/sec and sweeping action of approximately 1m/sec to ensure a droplet density of 20/cm<sup>2</sup>. Use a marking agent, as recommended by the equipment manufacturers, to check spray coverage. Also, consult directions provided with C.D.A. unit.

### **D. Low Volume High Concentrate Application Techniques**

Good control will be achieved, similar to high volume application, where bush size enables good coverage of entire bush. Use a marking agent, as recommended by the equipment manufacturers, to check spray coverage.

**Gas Powered Gun:** Apply 50 mL shots to obtain uniform coverage to 4 to 5m<sup>2</sup> of surface area of bush. This relates to 20 droplets/cm<sup>2</sup> of leaf surface.

**Sprinkler Sprayer:** This technique involves using a micro sprinkler that is connected to a hollow fibreglass rod attached to a pneumatic knapsack sprayer. Use at low pressures (50 to 200kPa) and apply with a slow sweeping action over the top of the plants ensuring even coverage of the leaves.

### **E. Boom Application**

Application in a minimum spray volume of 200L/ha for Galenia and St. John's Wort and 600L of water/ha for Sicklepod. Flat fan nozzles are recommended, using pressure in the range of 200-300 kPa. Boom height must be set to ensure double overlap of nozzle patterns.

## **2. FALLOW SITUATIONS**

### **A. Boom Application**

Application of Imtrade Picker Herbicide in a minimum spray volume of 50L/ha is recommended. Flat fan nozzles are recommended, using pressure in the range of 200-300 kPa. Boom height must be set to ensure double overlap of nozzle patterns.

### **B. Blanket Wiper Application**

Blanket needs to be made from durable wettable material with a rigid backing. Blanket should be rigidly mounted behind motorised vehicle (e.g. tractor, 4-wheel drive vehicle) and set low but never touching the ground. The chemical solution should be fed to the blanket at a flow rate sufficient to keep the blanket wet but not dripping. In thick patches the blanket may require more frequent solution recharge (re-wetting). Ideally, a scraper bar should be mounted in front of the blanket in order to scrape or damage the bark (but not sever the stems) prior to the blanket wiping the stems. This scraper may be mounted at the front of the vehicle. Two passes (in opposite direction) with the blanket increases the contact with the plant. Ground speeds of 10-15 kph are ideal for blanket wiping application.

## CLEANING SPRAY EQUIPMENT

### Rinsing

After using Imtrade Picker Herbicide, empty the spray unit completely and drain the whole system. Thoroughly wash inside the unit using a pressure hose. Drain the spray unit, and clean any filters in the tank, pump, lines, hoses and nozzles. After cleaning the spray unit as above, quarter fill with clean water and circulate through the pump, lines, hoses and nozzles. Drain and repeat the rinsing procedure twice.

### Decontamination

Before spraying cotton and other sensitive crops with equipment that has been used to apply Imtrade Picker Herbicide, see PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS section. Wash the tank and rinse the system as above. Then quarter fill the tank and add an alkali detergent (e.g. Liquid SURF®, OMO®, OMOMATIC®, DRIVE at 500 mL/100L of water or the powder equivalent at 500g/100L water) and circulate throughout the system for at least 15 minutes. Drain the whole system. Remove filters and nozzles and clean them separately. Finally flush the system with clean water and allow to drain.

**Rinse water should be discharged onto a designated disposal area or, if this is unavailable, onto unused land away from desirable plants and watercourses.**

## RESISTANT WEEDS WARNING

### GROUP 4 HERBICIDE

Imtrade Picker Herbicide is a member of the Pyridines group of herbicides. Imtrade Picker Herbicide has the disruptors of plant cell growth mode of action. For weed resistance management Imtrade Picker Herbicide is a Group 4 Herbicide. Some naturally occurring weed biotypes resistant to Imtrade Picker Herbicide and other disruptors of plant cell growth herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Imtrade Picker Herbicide or other disruptors of plant cell growth. Since the occurrence of resistant weeds is difficult to detect prior to use, Imtrade Australia Pty Ltd accepts no liability for any loss that may result from the failure of Imtrade Picker Herbicide to control resistant weeds. Strategies to minimise the risk of herbicide resistance are available. Contact your farm chemical supplier, consultant, local Department of Agriculture, or local Imtrade Australia Pty Ltd representative.

## **PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS**

Crops susceptible to Imtrade Picker Herbicide include, but are not limited to: Peas, Lupins, Lucerne, Navy Beans, Peanuts, Soybeans and other Legumes, Cotton, Flowers, Fruit, Hops, Ornamentals, Shade Trees and Pinus spp., Potatoes, Safflower, Sugar Beet, Sunflowers, Tobacco, Tomatoes, Vegetables and Vines. Imtrade Picker Herbicide is damaging to susceptible crops during both growing and dormant periods. Grasses are normally unaffected and establish quickly after treatment. Picloram, one of the active constituents in this product can remain active in the soil for extended periods depending on the soil type and application rate, rainfall, temperature, humidity, soil moisture and soil organic matter.

**DO NOT** apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands, pastures, waterways or native vegetation.

**DO NOT** allow physical spray drift onto waterways, native vegetation and susceptible crops.

**DO NOT** apply close to, or in areas, containing roots of desirable vegetation, where treated soil may be washed onto areas growing (or areas to be planted with) desirable plants.

**DO NOT** apply on sites where surface water from heavy rain can be expected to run off to areas containing, or to be planted with susceptible crops or plants. **DO NOT** move soil, which may have been treated to areas where desirable plants are to be grown.

## **PROTECTION OF LIVESTOCK**

Poisonous plants may become more palatable after spraying and stock should be kept away from these plants until they have died down. Many plants remain poisonous after death, and stock should not be allowed to access it, as there is a likelihood that they may graze the dead material. Such material should be burnt if possible.

## **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

**DO NOT** contaminate streams, rivers or waterways with the chemical or used containers.

## **STORAGE AND DISPOSAL (1L only)**

Store product in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. Rinse containers before disposal. Add rinsings to the spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers and product.



### **STORAGE AND DISPOSAL (5L, 10L, 20L, 200L only)**

Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. **DO NOT** store near food, fertilizers or seed. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Empty containers and product should **NOT** be burnt.

### **Envirodrum - Micro Matic Valve (110L only)**

Store the closed, original sealed Envirodrum in a well-ventilated area, as cool as possible and locked away from children, animals, food, foodstuffs, seed and fertilisers. **DO NOT** store for prolonged periods in direct sunlight. **DO NOT** tamper with the Micro Matic valve or the security seal. **DO NOT** contaminate the Envirodrum with water or any other foreign matter. After each use of the product, please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase.

### **Refillable containers (1000L only)**

Store in the closed, original container in a well-ventilated area, as cool as possible and locked away from children, animals, food, foodstuffs, seed and fertilisers. **DO NOT** store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

### **SMALL SPILL MANAGEMENT**

Wear protective equipment (See Safety Directions). Apply absorbent material such as earth, sand, clay granules or cat litter to the spill. Sweep up material for disposal when absorption is completed and contain in a refuse vessel for disposal. (See Storage and Disposal section). If necessary, wash the spill area with an alkali detergent and water and absorb as above the wash liquid for disposal.

### **SAFETY DIRECTIONS**

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When preparing the spray, wear cotton overalls buttoned to the neck and wrists, a washable hat, elbow-length PVC gloves and a face shield or goggles. If the product is in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

## FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre (*Phone Australia 13 11 26*).

### **Additional GHS Hazard & Precautionary Statements**

•Harmful if inhaled; •May cause respiratory irritation; •Avoid breathing fumes, mists, vapours or spray; •**DO NOT** get on clothing; •Wash contacted areas thoroughly after handling; •**DO NOT** eat, drink or smoke when using this product; •Use only outdoors or in a well-ventilated area; •**IF SWALLOWED:** Call a POISON CENTRE or doctor if you feel unwell; •**IF SWALLOWED:** Rinse mouth. Do **NOT** induce vomiting; •**IF ON SKIN:** Wash with plenty of soap and water; •**IF INHALED:** Remove victim to fresh air and keep at rest in a position comfortable for breathing; •If skin irritation occurs: Get medical advice; •If eye irritation persists: Get medical advice; •Not combustible. Use extinguishing media suited to burning materials; •Store locked up; •Store in a dry place.

## SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet which can be obtained from your supplier or Imtrade Australia Pty Ltd website at [www.imtrade.com.au](http://www.imtrade.com.au)

## CONDITIONS OF SALE

Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Imtrade Australia Pty Ltd has any authority to add to or alter these conditions.

® Other Registered Trademarks

**In a Transport Emergency  
Dial 000  
Police or Fire Brigade**

**APVMA Approval No: 56453/54458**