

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight.

All Pack Sizes unless otherwise stated

Triple-rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemical on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers or product. **DO NOT** re-use empty container for any other purpose.

Envirodrum 110L Mini Bulk Returnable Container only

DO NOT tamper with the Micro Matic valve or the security seal. **DO NOT** contaminate the Envirodrum with water or any foreign matter. After each use of the product, please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase.

Refillable Containers (1000L only)

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SMALL SPILL MANAGEMENT

Wear protective equipment (see SAFETY DIRECTIONS). Apply absorbent material such as earth, sand, clay granules or cat litter to the spill. Sweep up material for disposal when absorption is completed and contain in refuse vessel for disposal. (see STORAGE AND DISPOSAL section). If necessary, wash the spill area with an alkali detergent and water and absorb the wash liquid for disposal as above.

SAFETY DIRECTIONS

Will irritate the skin. Will damage the eyes. May irritate the nose and throat. Avoid contact with eyes and skin. **DO NOT** inhale spray mist. When opening the container and preparing product for use and using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length chemical resistant gloves, goggles, disposable mist face mask covering mouth and nose. In addition, when opening the container and preparing product for use, if applying by aerial spraying equipment, wear cotton overalls, over normal clothing, buttoned to the neck and wrist, elbow-length chemical resistant gloves, goggles and disposable mist face mask covering mouth and nose. In addition, if applying by spraying equipment carried on the back of the user, wear cotton overalls, over normal clothing, buttoned to the neck and wrist and a washable hat, elbow-length chemical resistant gloves, goggles and disposable mist face mask covering mouth and nose. If product on skin, immediately wash area with soap and water. If product or spray in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

IMTRADE

RALLY 300 SL

HERBICIDE

ACTIVE CONSTITUENT: 300 g/L CLOPYRALID present as the triisopropanolamine salt

GROUP 4 HERBICIDE

For the control of a wide range of broadleaf weeds in wheat, barley, oats, triticale, canola, pastures, fallow land, forests and industrial situations as specified in the Directions for Use table.

IMPORTANT: Read the attached Leaflet before use.

GROUP

4

FORM

SL

CONTENTS

1L-1000L



Batch No.
DOM:

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (*Phone Australia 13 11 26; New Zealand 0800 764 766*).

Refer to attached Leaflet for additional GHS Hazard & Precautionary Statements.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet which can be obtained from your supplier or Imtrade Australia Pty Ltd website at www.imtrade.com.au

CONDITIONS OF SALE

Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever, or howsoever arising through negligence or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express conditions that the purchaser does not rely on Imtrade Australia's skill or judgement in purchasing or using the same and every person dealing with this product does so at his/her own risk absolutely. No representative of Imtrade Australia Pty Ltd has any authority to alter these conditions.

In a Transport Emergency Dial 000
Police or Fire Brigade

APVMA Approval No: 69632/114212

 **IMTRADE**
CropScience

17 Ocean St, Kwinana Beach WA 6167
08 9419 0333 | www.imtrade.com.au

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



IMTRADE RALLY 300 SL HERBICIDE

ACTIVE CONSTITUENT: 300 g/L CLOPYRALID
present as the triisopropanolamine salt

GROUP

4

HERBICIDE

For the control of a wide range of broadleaf weeds in wheat, barley, oats, triticale, canola, pastures, fallow land, forests and industrial situations as specified in the Directions for Use table.

**IMPORTANT: READ THIS LEAFLET THOROUGHLY
BEFORE OPENING OR USING THIS PRODUCT**

APVMA Approval No: 69632/114212

Imtrade Australia Pty Ltd
ABN 13 090 151 134
17 Ocean Street, Kwinana WA 6167
Tel: 1800 171 799 Fax: 1800 171 788
Web: www.imtrade.com.au

DIRECTIONS FOR USE:

IT IS ESSENTIAL to select a rate appropriate to weed size. Best results will be obtained when weeds are actively growing at treatment.

RESTRAINTS

DO NOT apply to weeds which may be stressed (inactive growth) due to prolonged periods of extreme heat or cold, moisture stress (water logging or drought) or previous herbicide treatment as reduced levels of control may result.

DO NOT sow susceptible crops (see PROTECTION OF CROPS) for nine months following any application up to 300mL/ha or for twelve months following an application in excess of 300 mL/ha up to 500 mL/ha. Field peas should not be sown the season following an application of 500 mL/ha. Where rates in excess of 500 mL/ha have been used, susceptible crops including field peas should not be sown for at least two years.

DO NOT spray if rain is likely within 3 hours.

CROP	CROP STAGE	WEED	WEED STAGE	RATE mL/ha	STATE	CRITICAL COMMENTS
Barley, Oats, Triticale, Wheat.	Pre-sowing	Capeweed	Up to 8 leaf and maximum 10cm diameter	150 plus Knockdown herbicide	WA only	Pre-sowing: This rate should only be used in tank mixtures with formulations of paraquat/diquat or glyphosate.
	Post-sowing pre-emergence through to 3 leaf			150 plus Diuron at label rates		Post-sowing pre-emergent to 3 leaf: This rate should only be used in tank mixtures with Diuron for control of transplants. Refer to Diuron label for specific rates.
	Early post-emergence (2 leaf to jointing)		Cotyledons to 6 leaf and maximum 5cm diameter	150		TANK MIXTURES: Read and follow all label directions including restraints, spray drift restraints, mandatory no-spray zones, critical comments, withholding periods, regional use restrictions and safety directions for the tank mix products. Early post-emergent: Weeds should be growing actively and not larger than 5cm in diameter.

CROP	CROP STAGE	WEED	WEED STAGE	RATE mL/ha	STATE	CRITICAL COMMENTS
Barley, Oats, Triticale, Wheat.	4 to 5 leaf onwards	Capeweed, Soldier Thistle	Up to 10cm diameter (4 to 8 leaf)	300	NSW, ACT, Vic, SA, Tas and WA only	Weeds should be young and actively growing. Weeds will become stunted and not be competitive soon after application although final results may not show for some weeks. Faba Beans and Lupins will only be suppressed.
		Volunteer Chickpeas, Lentils and Safflower	Up to 6 leaf	250		
		Volunteer Faba Beans and Lupins	Up to 4 leaf			
		Volunteer Field Peas	Maximum 10cm high or 6 nodes	150	NSW, ACT, Vic, SA, Tas and WA only	Weeds should be young and actively growing. Weeds will become stunted and not be competitive soon after application although final results may not show for some weeks. Use 75 mL/ha rate only in combination with MCPA LVE.
				75 plus 700 mL/ha MCPA LVE		
		Volunteer Medics and seedling Lucerne	Up to 8 leaf	150		
		Volunteer Sub-clover	Up to 6 leaf			
		Volunteer Vetch	Runners up to 10cm maximum 16 leaf	100		
				75 plus 700 mL/ha MCPA LVE		
Prickly Lettuce	4 to 6 leaf and maximum 8cm diameter	150 plus 700 mL/ha MCPA LVE				
Thistles including: Nodding, Saffron, Scotch, Slender, Spear, Stemless, Variegated	Rosettes up to 10cm maximum diameter	50 plus 1L/ha MCPA Amine (500g/L) or 50 plus 700 mL/ha MCPA LVE	NSW, ACT, Vic, SA, Tas, WA and Qld only			

CROP	CROP STAGE	WEED	WEED STAGE	RATE mL/ha	STATE	CRITICAL COMMENTS
Barley, Oats, Triticale, Wheat.	5 leaf to late tillering	Skeleton Weed	5 to 15cm rosettes	500 plus 1L/ha MCPA Amine (500g/L)	NSW, ACT, Vic and SA only	Weeds should be a minimum 5cm in diameter, and growing actively. This rate will give control until harvest and substantially reduce weed numbers the following seasons.
Canola	2 to 8 leaf	Capeweed, Saffron Thistle, Skeleton Weed, Soldier Thistle	Up to 10cm diameter (4 to 8 leaf)	300	NSW, ACT, Vic, SA, Tas, WA and Qld only	Weeds should be young and actively growing. Weeds will become stunted and not be competitive soon after application although final results may not show for some weeks. Skeleton Weed will only be controlled until harvest. Faba Beans and Lupins will only be Suppressed. For the control of annual grasses, Imtrade Rally 300 SL Herbicide may be tank mixed with Imtrade Haloxyfop 520 Herbicide.
		Volunteer Chickpeas, Lentils and Safflower	Up to 6 leaf	250		
		Volunteer Faba Beans and Lupins	Up to 4 leaf			
		Volunteer Field Peas	Maximum 10cm high or 6 nodes	150		
		Volunteer Medics and seedling Lucerne	Up to 8 leaf			
		Volunteer Sub-clover	Up to 6 leaf			
		Volunteer Vetch	Runners up to 10cm maximum 16 leaf	100		

CROP	CROP STAGE	WEED	WEED STAGE	RATE mL/ha	STATE	CRITICAL COMMENTS
Pastures and Fallow land	Post-emergence	Hardhead Thistle (Creeping Knapweed, Russian Knapweed)	Actively growing plants	Handgun: 500 mL/100L of water Boom Spray: 2 or 4L/ha	Vic only	NOTE: DO NOT USE ON LUCERNE, CLOVERS AND MEDICS WILL BE ELIMINATED FOR AT LEAST ONE YEAR. Victoria only: Use the lower rate only on light soils (sand and sandy loam) where a slightly lower degree of control is acceptable. Use the higher rate on all soil types where complete control is required. Addition of a wetting agent at label rates is recommended for treatment of Hardhead Thistle. Spray between September and April on actively growing plants for effective control. Thorough coverage is essential. Apply in 200 to 250L of water/ha. BOOM SPRAYING: Use the higher rates of Imtrade Rally 300 SL Herbicide plus MCPA on multi-crowned plants or rosettes larger than 30cm in diameter. Spraying may be done at any time during active growth, usually in early winter or spring. Avoid spraying during the dormant winter period or at any time when thistles are not actively growing. DO NOT spray flowering thistles. PRE-SPRAY MANAGEMENT: The pasture should be slightly grazed prior to spraying to reduce clover and grass cover and expose the smaller thistles to the spray. The grazed pasture should be left seven days to allow thistles to freshen prior to treatment.
				Handgun: 500 mL/100L of water Boom Spray: 2 or 4L/ha	Qld only	
		Thistles including: Nodding, Variegated, Scotch, Spear, Slender, Saffron	Rosette stage prior to stem elongation. Treat rosette stage prior to stem elongation	50 or 70 mL/ha plus 1 to 1.5L MCPA Amine (500g/L) Drench Gun: 50 mL/1L water Handgun: 250 mL/100L of water	NSW, ACT, Vic, Tas, SA and Qld only	

Continued next page...

CROP	CROP STAGE	WEED	WEED STAGE	RATE mL/ha	STATE	CRITICAL COMMENTS
Pastures and Fallow land	Post-emergence	Hardhead Thistle (Creeping Knapweed, Russian Knapweed)	Actively growing plants	Handgun: 500 mL/ 100L of water Boom Spray: 2 or 4L/ha	Vic only	<i>...Continued from previous page</i> POST-TREATMENT MANAGEMENT: Response of thistles to treatment with Imtrade Rally 300 SL Herbicide plus MCPA mixture will be slowed compared to the standard treatments with 2,4-D or MCPA.. If possible delay grazing of sprayed thistles for 14 days after treatment. CLOVER DAMAGE: The Imtrade Rally 300 SL Herbicide plus MCPA mixture can be very damaging to Subterranean Clover. The lower rate is no more damaging than label rates of 2,4-D or MCPA. The higher rate of the Imtrade Rally 300 SL Herbicide plus MCPA mixture will reduce the clover component of the pasture for about two months. Clover recovery will be quicker during periods of active growth. HANDGUN (Spot Spray): Treat from rosette stage to early flowering. Thorough spraying is necessary. DRENCH GUN: Apply 10 mL to rosette crown. To multi-crown plants, apply 10mL to each crown.
				Handgun: 500 mL/ 100L of water Boom Spray: 2 or 4L/ha	Qld only	
		Thistles including: Nodding, Variegated, Scotch, Spear, Slender, Saffron	Rosette stage prior to stem elongation. Treat rosette stage prior to stem elongation	50 or 70 mL/ha plus 1 to 1.5L MCPA Amine (500g/L) Drench Gun: 50 mL/1L water Handgun: 250 mL/100L of water	NSW, ACT, Vic, Tas, SA and Qld only	

CROP	CROP STAGE	WEED	WEED STAGE	RATE mL/ha	STATE	CRITICAL COMMENTS
Pastures and Fallow land	Post-emergence	Nodding Thistle	Rosettes up to 20cm diameter	100	NSW, ACT only	Apply the spray from September to October. Apply by boom spray only. DO NOT apply to thistles over 20cm in diameter. When thistles are over 20cm in diameter use Imtrade Rally 300 SL Herbicide plus MCPA (referred to above). Clover Damage: Damage to White Clover will be no greater than damage with MCPA alone and less than damage from Imtrade Rally 300 SL Herbicide plus MCPA or 2,4-D alone. DO NOT use for spot treatment.
		California Thistle	From early buds to flowering (December to February)	Hand Gun: 250 mL/100L Water Boom Spray: 2L/ha	Vic and Tas only	Addition of a wetting agent at label rates is recommended. Re-treatment of regrowth in the year following treatment will usually be necessary to achieve a high level of control. NOTE: Clovers and Medics will be eliminated for at least one year.
Pasture, Forests, Rights-of-Way, Industrial situations	Post-emergence	Groundsel Bush	Young seedlings to mature plants	Hand gun: 330 or 500 mL/100L of water	Qld, NSW and ACT only	Spray foliage when growth is active. Use the lower rate on young seedlings and the higher rate on plants more than 2cm tall or when growth is slow.

CROP	CROP STAGE	WEED	WEED STAGE	RATE mL/ha	STATE	CRITICAL COMMENTS
Pastures and <i>Pinus radiata</i> plantations	Post-emergence	Silver Wattle	Active growth spring to summer	<p>Handgun: 500 mL/100L water</p> <p>Boom and Aerial spray: 5L/ha (weeds 30cm to 2m) 7.0L/ha (weeds 2 to 4m) 8.5L/ha (weeds 4 to 8m)</p>	NSW, ACT, Vic and Tas only	<p>For effective control apply when bushes are growing actively. Large trees will not show complete necrosis.</p> <p>HANDGUN: Means high volume NOT low volume knapsack. Spray to give full coverage of leaves and stems. Add organosilicone surfactant at 200 mL/100L for optimum results.</p> <p>See General Instructions for handgun spraying.</p> <p>BOOM AND AERIAL SPRAYING: For boom spraying apply in 150 to 200L of water/ha. For aerial treatment apply in a minimum of 50L/ha of water containing 25 to 50% by volume of anti-evaporant oil such as Throttle*. Maintain continuous agitation. Clovers and other legumes will be eliminated for at least one year.</p>
Forests		Cape Ivy	Any growth stage	<p>Handgun: 3.3L/ha</p>	Vic and Tas only	<p>Apply by hand held weed wiper or C.D.A., at dilutions with water at 1:3. Application may be made at any time of the year provided foliage is dry at the time. Avoid spraying non-target plants.</p>

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

PASTURES and FALLOW LAND: DO NOT GRAZE OR CUT FOR STOCK FEED FOR 7 DAYS AFTER APPLICATION.

CEREALS: DO NOT HARVEST FOR 10 WEEKS AFTER APPLICATION.

CANOLA: DO NOT HARVEST FOR 12 WEEKS AFTER APPLICATION.

CEREALS and CANOLA: DO NOT GRAZE OR CUT TREATED CEREALS FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.

FORESTS, except *Pinus radiata* PLANTATIONS: DO NOT GRAZE FOR 7 DAYS AFTER APPLICATION.

***Pinus radiata* PLANTATIONS: DO NOT GRAZE FOR 14 DAYS AFTER APPLICATION.**

GENERAL INSTRUCTIONS

MIXING

Half fill the spray tank with water and add the required quantity of Imtrade Rally 300 SL Herbicide and complete filling. Agitate continuously to ensure thorough mixing before and during application.

Only mix sufficient chemical for each days work.

Tank mixtures: Wettable powder or dry flowable formulations should be added to the spray tank first, followed by suspension concentrates (flowables), aqueous concentrates (Imtrade Rally 300 SL Herbicide) and the emulsifiable concentrate formulations (e.g. MCPA LVE).

APPLICATION

Boom Spraying Crop and Pastures: Apply Imtrade Rally 300 SL Herbicide in sufficient water to obtain good coverage. It should be applied by an accurately calibrated ground rig or aircraft, delivering 200 to 300 micron droplets and not less than 50L/ha water volume for boom sprayers or not less than 20L/ha for aerial applications.

Hardhead Thistle – Use a spray volume of 200 to 250L/ha of water.

Silver Wattle – Use a spray volume of 150 to 200L/ha of water by ground boom spray and a minimum spray volume of 50L/ha by aircraft.

High Volume Handgun: Apply the recommended mix to give full coverage of leaves and stems through a No. 6-8 tip at 700 to 1500 kPa. Spray volume for effective coverage of dense two metre high Silver Wattle should be 30 to 40L of spray per 100m² (10m x 10m) of infestation. For larger areas an equivalent would be 3000 to 4000L per infested hectare.

CLEANING SPRAY EQUIPMENT

Rinse water should be discharged onto a designated disposal area or, if this is unavailable, onto unused land away from desirable plants and watercourses.

Partial cleaning (before spraying other labeled tolerant crops):

- After using Imtrade Rally 300 SL Herbicide, empty the tank completely and drain the whole system. Thoroughly wash inside the tank using pressure hose. Quarter fill the tank with clean water and circulate through the pump, line, hoses and nozzles. Drain and repeat procedure twice.

Complete cleaning (before spraying susceptible crops):

- After using Imtrade Rally 300 SL Herbicide, empty the tank completely and drain the whole system. Thoroughly wash inside the tank using a pressure hose. Quarter fill the tank with clean water and circulate as above then drain.
- Quarter fill the tank again and add a liquid alkali detergent (e.g. Surf, Omo, Drive) at 500 mL/100L water and circulate throughout the system for at least fifteen minutes.
- Drain, remove filters and nozzles and clean separately. Rinse inside the tank thoroughly using a pressure hose and flush system with clean water.

COMPATIBILITY

Imtrade Rally 300 SL Herbicide is compatible with the following:

Broadleaf Herbicides: ChemAg Rockstar 200 Herbicide, Chlorsulfuron, Bromoxynil, Metsulfuron, Diuron, Glyphosate, MCPA Amine, MCPA LVE, Imtrade Paraquat 250 Herbicide, Imtrade Spraykill 250 Herbicide, Terbutryn, 2,4-D Amine.

Grass Herbicides in cereal crops: Diclofop-methyl 375g/L, Tralkoxydim 400g/kg, Fenoxaprop-p-ethyl 110g/L (Tralkoxydim and Fenoxaprop-p-ethyl for Wild Oat control only).

Grass Herbicides on broadleaf crops: Imtrade Haloxyfop 520 Herbicide.

RESISTANT WEEDS WARNING

GROUP 4 HERBICIDE

Imtrade Rally 300 SL Herbicide is a member of the Pyridines group of herbicides. The product has the disruptors of plant cell growth mode of action. For weed resistance management, the product is a Group 4 Herbicide. Some naturally-occurring weed biotypes resistant to Imtrade Rally 300 SL Herbicide and other Group 4 Herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Imtrade Rally 300 SL Herbicide or other Group 4 Herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Imtrade Australia Pty Ltd accepts no liability for any losses that may result from the failure of Imtrade Rally 300 SL Herbicide to control resistant weeds.

Strategies to minimize the risk of herbicide resistance are available. Contact your farm chemical supplier, consultant, local Department of Agriculture or local Imtrade Australia Pty Ltd representative.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment that may cause spray drift onto nearby susceptible plants/crops, cropping lands or pastures.

DO NOT apply Imtrade Rally 300 SL Herbicide to crops or pastures, which are to be used for the production of compost or mulches for use with susceptible crops or plants. The use of straw, hay or other plant material treated with Imtrade Rally 300 SL Herbicide for composting or mulching susceptible crops may damage these crops. Susceptible crops include Chickpeas, Cotton, Faba Beans, Field Peas, Fruit Trees, Lentils, Lupins, Lucerne, Medics, Ornamentals, Potatoes, Safflower, Sub-clover, Tomatoes, Vegetables, Vines, Wattles and White Clover.

DO NOT allow spray drift onto susceptible plants.

Plant back periods

Residues in the straw of treated crops can affect subsequent susceptible crops. Susceptible crops listed above should not be sown for:

Nine months following any application up to 300 mL/ha.

Twelve months following an application in excess of 300 mL/ha and up to 500 mL/ha.

PROTECTION OF LIVESTOCK

DO NOT graze or cut treated crops for stock food except as specified under Withholding Periods.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT

Imtrade Rally 300 SL Herbicide has low toxicity to fish, birds, honeybees, livestock, earthworks and aquatic organisms.

DO NOT contaminate streams, rivers or waterways with this product or used container.

STORAGE AND DISPOSAL

The method of disposal of the container depends on the container type. Read the Storage and Disposal instructions on the label that is attached to the container.

Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight.

All Pack Sizes unless otherwise stated

Triple-rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemical on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers or product. **DO NOT** re-use empty container for any other purpose.

Envirodrum 110L Mini Bulk Returnable Container only

DO NOT tamper with the Micro Matic valve or the security seal. **DO NOT** contaminate the Envirodrum with water or any foreign matter. After each use of the product, please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase.

Refillable Containers (1000L only)

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SMALL SPILL MANAGEMENT

Wear protective equipment (see SAFETY DIRECTIONS). Apply absorbent material such as earth, sand, clay granules or cat litter to the spill. Sweep up material for disposal when absorption is completed and contain in refuse vessel for disposal. (see STORAGE AND DISPOSAL section). If necessary, wash the spill area with an alkali detergent and water and absorb the wash liquid for disposal as above.

SAFETY DIRECTIONS

Will irritate the skin. Will damage the eyes. May irritate the nose and throat. Avoid contact with eyes and skin. **DO NOT** inhale spray mist. When opening the container and preparing product for use and using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length chemical resistant gloves, goggles, disposable mist face mask covering mouth and nose. In addition, when opening the container and preparing product for use, if applying by aerial spraying equipment, wear cotton overalls, over normal clothing, buttoned to the neck and wrist, elbow-length chemical resistant gloves, goggles and disposable mist face mask covering mouth and nose. In addition, if applying by spraying equipment carried on the back of the user, wear cotton overalls, over normal clothing, buttoned to the neck and wrist and a washable hat, elbow-length chemical resistant gloves, goggles and disposable mist face mask covering mouth and nose. If product on skin, immediately wash area with soap and water. If product or spray in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26; New Zealand 0800 764 766).

Additional GHS Hazard & Precautionary Statements

•**IF SWALLOWED:** Rinse mouth. Do **NOT** induce vomiting; •If skin irritation occurs: Get medical advice; •If eye irritation persists: Get medical advice; •Collect spillage; •In case of fire: Alcohol resistant foam is the preferred firefighting medium but, if it is not available, normal foam can be used; •Store in a dry place.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet which can be obtained from your supplier or Imtrade Australia Pty Ltd website at www.imtrade.com.au

CONDITIONS OF SALE

Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever, or howsoever arising through negligence or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express conditions that the purchaser does not rely on Imtrade Australia's skill or judgement in purchasing or using the same and every person dealing with this product does so at his/her own risk absolutely. No representative of Imtrade Australia Pty Ltd has any authority to alter these conditions.

* Other Company Registered Products

In a Transport Emergency
Dial 000
Police or Fire Brigade

APVMA Approval No: 69632/114212