

**STORAGE AND DISPOSAL (5L, 10L, 20L & 200L only)**

Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. **DO NOT** store tank-mixes of Imtrade Tiger Gold 250 Herbicide with other materials. When storing, keep from contact with fertilisers, fungicides and seeds. Triple or (preferably) pressure rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers or product.

**Envirodrum - Micro Matic Valve (110L)**

Store the original sealed Envirodrum in a cool, well ventilated area. **DO NOT** store for prolonged periods in direct sunlight. **DO NOT** tamper with the Micro Matic valve or the security seal. **DO NOT** contaminate the Envirodrum with water or any other foreign matter. After each use of the product please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase.

**Refillable containers (1000L only)**

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

**SAFETY DIRECTIONS**

Poisonous if inhaled. Will irritate the skin. Will damage the eyes. Repeated exposure may cause allergic disorders. **DO NOT** inhale vapour. Avoid contact with eyes and skin. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length chemical resistant gloves, goggles and half facepiece respirator. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length chemical resistant gloves.

If applying by hand, wear cotton overalls, over normal clothing, buttoned to the neck and wrist, and a washable hat, elbow-length chemical resistant gloves. If product on skin, immediately wash area with soap and water. If product in eyes wash out immediately with water. Wash hands after use. After each day's use, wash gloves, goggles, respirator and if rubber wash with detergent and warm water and contaminated clothing.

**RE-ENTRY PERIOD**

**DO NOT** allow entry into treated areas for 8 days when applying to summer growing weeds, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

**DO NOT** allow entry into treated areas until the spray has dried when applying to winter growing weeds, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

**DO NOT APPLY USING SPRAYING EQUIPMENT CARRIED ON THE BACK OF THE USER.**

**FIRST AID**

If swallowed, **DO NOT** induce vomiting. Give a glass of water. If poisoning occurs, consult a doctor or Poisons Information Centre (Phone Australia 13 11 26; New Zealand 0800 764 766). If in eyes, wash out immediately with water.

Refer to attached Leaflet for additional GHS Hazard & Precautionary Statements

**SAFETY DATA SHEET**

Additional information is listed in the Safety Data Sheet which can be obtained from your supplier or Imtrade Australia Pty Ltd website at [www.imtrade.com.au](http://www.imtrade.com.au)

**CONDITIONS OF SALE**

Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever, or howsoever arising through negligence or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express conditions that the purchaser does not rely on Imtrade Australia's skill or judgement in purchasing or using the same and every person dealing with this product does so at his/her own risk absolutely. No representative of Imtrade Australia Pty Ltd has any authority to alter these conditions.

# POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

## IMTRADE TIGER GOLD 250 HERBICIDE

ACTIVE CONSTITUENT: 250 g/L QUIZALOFOP-P-ETHYL  
SOLVENT: 493 g/L HYDROCARBON LIQUID  
145 g/L N-METHYL-2-PYRROLIDONE

**GROUP 1 HERBICIDE**

For the control of certain grass weeds in broadleaf crops as specified  
in the Directions For Use table.

**Important: Read the attached Leaflet before use**

<b>UN 3082</b>	ENVIRONMENTALLY HAZARDOUS SUBSTANCES, LIQUID, N.O.S. (contains QUIZALOFOP-P-ETHYL)
In a Transport Emergency Dial 000 Police or Fire Brigade	
PG III	HAZCHEM +3Z

APVMA Approval No. 65807/106313



**GROUP  
1**

**FORM  
EC**

**CONTENTS  
1L-1000L**



Batch No.  
DOM:

**IMTRADE**  
CropScience  
17 Ocean St, Kwinana Beach WA 6167  
08 9419 0333 | [www.imtrade.com.au](http://www.imtrade.com.au)

**POISON**

**KEEP OUT OF REACH OF CHILDREN**

**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**



# **IMTRADE TIGER GOLD 250 HERBICIDE**

**ACTIVE CONSTITUENT: 250 g/L QUIZALOFOP-P-ETHYL**

**SOLVENT: 493 g/L HYDROCARBON LIQUID**

**145 g/L N-METHYL-2-PYRROLIDONE**

<b>GROUP</b>	<b>1</b>	<b>HERBICIDE</b>
--------------	----------	------------------

**For the control of certain grass weeds in broadleaf crops  
as specified in the Directions For Use table.**

**IMPORTANT: READ THIS LEAFLET THOROUGHLY  
BEFORE OPENING OR USING THIS PRODUCT**

**APVMA Approval No: 65807/106313**

Imtrade Australia Pty Ltd

ABN 13 090 151 134

17 Ocean Street, Kwinana, Western Australia 6167

Tel: 1800 171 799 Fax: 1800 171 788

Web: [www.imtrade.com.au](http://www.imtrade.com.au)

## **RESTRAINTS**

**DO NOT** apply to crops that are stressed by any cause (such as severe cold weather, disease, drought or waterlogging).

**DO NOT** apply to weeds that are stressed by any cause (such as weather conditions, drought, waterlogging) or not actively growing. Grass weeds under stress frequently become less susceptible to herbicide activity. This may result in incomplete kill or suppression of weeds only.

**DO NOT** apply under conditions of high temperature and low humidity.

**DO NOT** apply if rainfall is expected within 3 hours.

**DO NOT** apply less than 50L prepared spray/ha.

**DO NOT** apply using spraying equipment carried on the back of the user.

**DIRECTIONS FOR USE**  
**BOOM SPRAYING: WINTER GROWING WEEDS**

Crop	Weeds Controlled	Weeds Stage	Rate/ha	WHP		Critical Comments
				Harvest	Grazing	
Beetroot, Cabbage, Canola, Carrots, Cauliflower, Chickpeas, Clover & Sub-Clover	Annual Ryegrass ( <i>Lolium rigidum</i> ), Brome Grasses ( <i>Bromus</i> spp.)	3-leaf to early tillering	120 mL or 140 mL	14 days 9 wks 11 wks 10 wks 14 days 12 wks	- - 4 wks - - 4 wks	Apply when weeds are actively growing. Use higher rate under heavy weed pressure and/or when weeds have commenced tillering.*
Pastures, Cucumbers*, Faba Beans, Field Peas, Honey Dew Melons*, Lentils, Lucerne, Lupins, Medic Pasture, Mung Beans, Navy Beans, Onions, Potatoes, Pumpkins*, Radish, Tomatoes, Vetch	Barley Grass ( <i>Hordeum</i> spp.), Volunteer Barley, Volunteer Wheat Wild Oats ( <i>Avena</i> spp.)	3-leaf to early tillering   3-leaf to early tillering	100 mL    50 mL or 100 mL	- 14 days 12 wks 9 wks 9 wks 12 wks - 6 wks - 12 wks 12 wks 18 wks 10 wks 9 wks 21 days 4 wks 12 wks	- - 4 wks 4 wks - - 4 wks 14 days 4 wks 4 wks - - - - - 4 wks	Apply when weeds are actively growing.*    Apply when weeds are actively growing. Use the lower rate only under light weed pressure and/or when weeds are at the 1-3 leaf stage. Use higher rate under heavy weed pressure and/or when weeds are tillering. The higher rate will give better or more reliable control.*

\* ALWAYS ADD A SURFACTANT/WETTING AGENT AT THE RECOMMENDED RATE – SEE SECTION ON “USE OF SURFACTANT/WETTING AGENT” FOR DETAILS.

• IMPORTANT DO NOT APPLY INTRADE TIGER GOLD 250 HERBICIDE BEFORE THE FIVE TRUE LEAF STAGE OF THE CROP.

# BOOM SPRAYING: SUMMER GROWING WEEDS

Crop	Weeds Controlled	Weeds Stage	Rate/ha	WHP		Critical Comments
				Harvest	Grazing	
Beetroot, Cabbage, Cauliflower, Carrots, Cucumbers*, Green Beans, Honey Dew Melons*, Lucerne, Mung Beans, Navy Beans, Onions, Peanuts, Potatoes, Pineapples, Pumpkins*, Radish, Soybeans, Sunflowers, Tomatoes	Awnless Barnyard Grass ( <i>Echinochloa colona</i> ), Barnyard Grass ( <i>Echinochloa crus-galli</i> ),	3-leaf to early tillering	200 mL or 300 mL	14 days	-	Apply when weeds are actively growing. Use higher rate under heavy weed pressure and/or when weeds have commenced tillering.*
				9 wks	-	
				14 days	-	
				10 wks	-	
				14 days	-	
	Couch Grass ( <i>Cynodon dactylon</i> )	2-4 leaf seedling	200 mL	5 wks	4 wks	Apply when weeds are actively growing. This rate will only control seedling Couch Grass which is less than 2 months old and established from seed.*
				9 wks	-	
				-	4 wks	
				12 wks	4 wks	
				12 wks	4 wks	
	Dinebra ( <i>Dinebra retroflexa</i> )	3-leaf to early tillering	100 mL or 150 mL	18 wks	-	Apply when weeds are actively growing. Use higher rate under heavy weed pressure and/or when weeds have commenced tillering.*
				11 wks	11 wks	
				10 wks	-	
				7 days	-	
				9 wks	-	
	Crowsfoot Grass ( <i>Eleusine indica</i> )	2 leaf to early tillering	200 mL	21 days	-	Apply when weeds are actively growing. Use higher rate under heavy weed pressure and/or when weeds have commenced tillering.*
				12 wks	4 wks	
				9 wks	11 wks	
				4 wks	-	
				-	-	
	Foxtail Millet ( <i>Setaria italica</i> )	2 leaf to early tillering	100 mL or 200 mL			Apply when weeds are actively growing. Use higher rate under heavy weed pressure and/or when weeds have commenced tillering.*

# BOOM SPRAYING: SUMMER GROWING WEEDS (continued)

Crop	Weeds Controlled	Weeds Stage	Rate/ha	WHP		Critical Comments
				Harvest	Grazing	
Beetroot, Cabbage, Cauliflower, Carrots, Cucumbers*, Green Beans, Honey Dew Melons*, Lucerne, Mung Beans, Navy Beans, Onions, Peanuts, Potatoes, Pineapples, Pumpkins*, Radish, Soybeans, Sunflowers, Tomatoes	Columbus Grass ( <i>Sorghum xalmum</i> ), Johnson Grass ( <i>Sorghum halepense</i> ), Liverseed Grass ( <i>Urochloa panicoides</i> )	2-leaf to 3 tiller	200 mL or 300 mL	14 days	-	Apply when weeds are actively growing. Use higher rate under heavy weed pressure and/or when weeds have commenced tillering.*
				9 wks	-	
				14 days	-	
				10 wks	-	
	Rhodes Grass ( <i>Chloris gayana</i> ), Stink Grass ( <i>Eragrostis cilianensis</i> ),	3-leaf to early tillering	100 mL or 200 mL	14 days	-	
				9 wks	4 wks	
				12 wks	4 wks	
				12 wks	4 wks	
	Queensland Blue Grass ( <i>Dichanthium sericeum</i> )	3-leaf to early tillering	200 mL	18 wks	-	Apply when weeds are actively growing.*
				11 wks	11 wks	
				10 wks	-	
				7 days	-	
	Summer Grass ( <i>Digitaria ciliaris</i> )	3-leaf to early tillering	200 mL or 400 mL	9 wks	-	Apply when weeds are actively growing. Use higher rate under heavy weed pressure and/or when weeds have commenced tillering.*
				21 days	-	
				12 wks	4 wks	
				9 wks	11 wks	
				4 wks	-	

- \* ALWAYS ADD A SURFACTANT/WETTING AGENT AT THE RECOMMENDED RATE – SEE SECTION ON “USE OF SURFACTANT/WETTING AGENT” FOR DETAILS.
- IMPORTANT DO NOT APPLY IMTRADE TIGER GOLD 250 HERBICIDE BEFORE THE FIVE TRUE LEAF STAGE OF THE CROP.

## HAND GUN SPRAYING

Crop	Weeds Controlled	Weeds Stage	Rate/ha	WHP		Critical Comments
				Harvest	Grazing	
Beetroot,	Awnless Barnyard Grass ( <i>Echinochloa colona</i> ), Cauliflower, Canola, Carrots, Chickpeas, Clover & Sub-Clover	3-leaf to early tillering	50 mL	14 days	-	Apply when weeds are actively growing. Use a minimum of 800L of prepared spray/ha.
Cabbage,				9 wks	-	
Pasture,				14 days	-	
Cucumbers*,				11 wks	4 wks	
Faba Beans,				10 wks	-	
Field Peas,	Crowsfoot Grass ( <i>Eleusine indica</i> )	Seedling to established clumps	50 mL	12 wks	4 wks	Thoroughly wet target weeds (especially well established clumps) without causing run-off. Repeated sprays may be necessary for perennial grass species.
Grapes,				-	14 days	
Green Beans,				14 days	-	
Honey Dew Melons*,				12 wks	4wks	
Lentils,				9 wks	4 wks	
Lucerne,	Johnson Grass ( <i>Sorghum halepense</i> )	Seedling 2-leaf to early tillering	50 mL	-	4 wks	Always add a surfactant/wetting agent at the recommended rate – see section on “Use of Surfactant/ Wetting Agent” for details.
Lupins,				5 wks	4 wks	
Medic Pasture,				9 wks	-	
Mung Beans,				12 wks	4 wks	
Navy Beans,				-	4 wks	
Non-crop,	Rhizome 2-leaf to early tillering	Rhizome 2-leaf to early tillering	100 mL	6 wks	4 wks	
Onions,				-	14 days	
Peanuts,				12 wks	4 wks	
Pineapples,				12 wks	4 wks	
Potatoes,				-	-	
Pumpkins*,	Kikuyu ( <i>Pennisetum clandestinum</i> )	Seedling to established plants	100 mL	18 wks	-	Always add a surfactant/wetting agent at the recommended rate – see section on “Use of Surfactant/ Wetting Agent” for details.
Radish,				11 wks	11 wks	
Soybeans,				7 days	-	
Sunflowers,				10 wks	-	
Tomatoes,				9 wks	-	
Vetch				21 days	-	
				12 wks	4 wks	
				9 wks	11 wks	
				4 wks	-	
				12 wks	4 wks	

\* ALWAYS ADD A SURFACTANT/WETTING AGENT AT THE RECOMMENDED RATE – SEE SECTION ON “USE OF SURFACTANT/WETTING AGENT” FOR DETAILS.

• IMPORTANT DO NOT APPLY IMTRADE TIGER GOLD 250 HERBICIDE BEFORE THE FIVE TRUE LEAF STAGE OF THE CROP.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL  
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

**WITHOLDING PERIODS:**

**HARVEST:**

BEETROOT, CAULIFLOWER,  
CUCUMBERS:

DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION.

CABBAGE, FIELD PEAS, HONEY DUE  
MELON, PUMPKINS, SUNFLOWERS:  
CARROTS, POTATOES:

DO NOT HARVEST FOR 9 WEEKS AFTER APPLICATION.  
DO NOT HARVEST FOR 10 WEEKS AFTER  
APPLICATION.

CHICKPEAS, FABA BEANS, LENTILS,  
MUNG BEANS, NAVY BEANS,  
SOYBEANS, VETCH:

DO NOT HARVEST FOR 12 WEEKS AFTER  
APPLICATION.

CANOLA, PEANUTS:

DO NOT HARVEST FOR 11 WEEKS AFTER  
APPLICATION.

GREEN BEANS:

DO NOT HARVEST FOR 5 WEEKS AFTER APPLICATION.  
NOT REQUIRED WHEN USED AS DIRECTED.

GRAPES:

DO NOT HARVEST FOR 6 WEEKS AFTER APPLICATION.

LUPINS:

DO NOT HARVEST FOR 18 WEEKS AFTER APPLICATION.

ONIONS:

DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION.

PINEAPPLES:

DO NOT HARVEST FOR 21 DAYS AFTER APPLICATION.

RADISH:

DO NOT HARVEST FOR 4 WEEKS AFTER APPLICATION.

TOMATOES:

**GRAZING:**

CANOLA, CHICKPEAS, FABA BEANS,  
FIELD PEAS, GREEN BEANS, LENTILS,  
LUCERNE, LUPINS, MUNG BEANS,  
NAVY BEANS, SOYBEANS, VETCH:

DO NOT GRAZE OR CUT FOR STOCK FEED FOR 4  
WEEKS AFTER APPLICATION.

CLOVER, MEDICS, PASTURES (EXCEPT  
LUCERNE), SUB-CLOVER:

DO NOT GRAZE OR CUT FOR STOCK FEED FOR 14  
DAYS AFTER APPLICATION.

PEANUTS, SUNFLOWERS:

DO NOT GRAZE OR CUT FOR STOCK FEED FOR 11  
WEEKS AFTER APPLICATION.

## GENERAL INSTRUCTIONS

Imtrade Tiger Gold 250 Herbicide is an emulsifiable concentrate herbicide used post emergence for the control of certain grass weeds in specified broadleaf crops. Imtrade Tiger Gold 250 Herbicide is a systemic herbicide, which is rapidly absorbed and readily translocates from the treated foliage to the root system and growing points of the plant. Applications made to grass weeds stressed due to insufficient moisture or cold temperatures may result in unsatisfactory control.

## RESISTANT WEEDS WARNING

GROUP	<b>1</b>	HERBICIDE
-------	----------	-----------

Imtrade Tiger Gold 250 Herbicide is a member of the 'Fop' group of herbicides. Imtrade Tiger Gold 250 Herbicide has the inhibitors of acetyl coA carboxylase mode of action. For weed resistance management Imtrade Tiger Gold 250 Herbicide is a Group 1 Herbicide. Some naturally occurring weed biotypes resistant to Imtrade Tiger Gold 250 Herbicide and other Group 1 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Imtrade Tiger Gold 250 Herbicide or other Group 1 Herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Imtrade Australia Pty Ltd accepts no liability for any losses that may result from the failure of Imtrade Tiger Gold 250 Herbicide to control resistant weeds.

Strategies to minimise the risk of herbicide resistance are available. Consult your farm chemical supplier, consultant, local Department of Agriculture or Primary Industries, or local Imtrade Australia Pty Ltd Representative.

## TIME OF APPLICATION

For maximum benefit, weeds should be sprayed when young and actively growing. Application may be made at any stage of crop growth (except to cucurbit crops), but sufficient time must be allowed for the prescribed withholding period to expire before harvest and/or grazing or cutting for stockfeed.

**IMPORTANT: DO NOT** apply Imtrade Tiger Gold 250 Herbicide to cucurbit crops before the five true leaf stage.

## SPRAY PREPARATION

Imtrade Tiger Gold 250 Herbicide is an emulsifiable concentrate to be mixed with water and applied as a spray. Add the amount of Imtrade Tiger Gold 250 Herbicide directly to the spray tank and mix well. If tank mixing with Mancozeb, add this to the spray tank and mix thoroughly prior to adding Imtrade Tiger Gold 250 Herbicide.

## USE OF A SURFACTANT/WETTING AGENT

Imtrade Tiger Gold 250 Herbicide may be used with either of the following adjuvant systems:

1) A non-ionic surfactant (1000 g ac/L strength) at 200 mL/100L of final spray volume (i.e. 0.2% volume/volume).

2) A non-ionic surfactant (1000 g ac/L strength) at 0.1% v/v and a mineral spray oil at 1% v/v.

For Imtrade Tiger Gold 250 Herbicide and Imtrade Tiger Gold 250 Herbicide tank-mixes with Sethoxydim 186 g/L and Imtrade Clethodim 240 EC Herbicide the recommended mixture is 100 mL non-ionic surfactant + 1L DC-Trate® Spray Oil per 100L of final spray volume. This adjuvant system provides a more robust treatment than a surfactant only treatment and is strongly recommended.

**DO NOT** use the surfactant + mineral oil adjuvant system on vegetable crops, use a surfactant only.

**DO NOT** use a spray oil alone, a surfactant must always be added.

**DO NOT** use spray oils of crop origin (e.g. Codacide oil, Synertril oil) as spray adjuvants.

## COMPATIBILITIES

Imtrade Tiger Gold 250 Herbicide is compatible with the insecticides Imtrade Delta-Duo Insecticide, Imtrade Omen 290 Insecticide, Dimethoate, the fungicide Mancozeb and the herbicides Difen Options Selective Herbicide and Rally Herbicide when using the surfactant only adjuvant system.

Imtrade Tiger Gold 250 Herbicide is compatible with Sethoxydim 186 g/L and Imtrade Clethodim 240 EC Herbicide when using the surfactant plus spray oil adjuvant system.

When tank mixing Imtrade Tiger Gold 250 Herbicide and broadleaf herbicides **DO NOT** use a spray oil, use a surfactant only.

Application of the Imtrade Tiger Gold 250 Herbicide tank mix may result in crop effects. The effect is temporary and usually appears as yellow or white bleaching on the leaves. New growth and subsequent yield, however, will not be affected.

**DO NOT** add crop oil to tank mixtures of Imtrade Tiger Gold 250 Herbicide/Difen Options Selective Herbicide.

To minimise crop effects; Apply Imtrade Tiger Gold 250 Herbicide a minimum of 5 days prior to Difen Options Selective Herbicide application or a minimum 10 days after Difen Options Selective Herbicide application (14-21 days if weeds have been stressed).

## GROUND SPRAYING

Use sprayers with flat fan or hollow cone nozzles at 210-350 kPa to give an even distribution of droplets with no less than 50% of the spray volume in the 150-300 micron range. 110° flat fan nozzles are preferred.

Use between 50 and 100L of prepared spray/ha, the higher spray volume gives more reliable results. If weeds are large and the crop canopy is dense, increase the volume to 150L prepared spray/ha.

For proper spray coverage adjust the boom and nozzle height according to the specifications listed by the nozzle manufacturer. Ensure the spray is evenly distributed and a double overlap spray pattern is obtained. Avoid overlapping boom spray widths and shut off spray booms while starting, turning, slowing or stopping.

## **PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS**

**DO NOT** apply under weather conditions or from spraying equipment that may cause spray to drift onto nearby or susceptible plants/crops, cropping lands or pastures.

**DO NOT** plant cereal crops into the treated area for a period of 18 weeks after application of Imtrade Tiger Gold 250 Herbicide.

## **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

**DO NOT** contaminate streams, rivers or waterways with the chemical or used containers.

## **SPRAYER CLEANUP**

Thoroughly clean all traces of Imtrade Tiger Gold 250 Herbicide from application equipment immediately after use. Flush tank, pump, hoses and boom with several changes of water after removing nozzle tips and screens (clean those parts separately). Failure to thoroughly clean the equipment may result in injury to subsequently sprayed cereal and grass crops. If spray equipment has previously been used for the application of sulfonylurea herbicides, ensure that the recommended sprayer clean-up procedure for the product previously sprayed has been followed before using the equipment for the application of Imtrade Tiger Gold 250 Herbicide.

## **STORAGE AND DISPOSAL (1L & 2L only)**

Store product in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. Rinse containers before disposal. Add rinsings to the spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers and product.

## **STORAGE AND DISPOSAL (5L, 10L, 20L & 200L only)**

Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. **DO NOT** store tank-mixes of Imtrade Tiger Gold 250 Herbicide with other materials. When storing, keep from contact with fertilisers, fungicides and seeds. Triple or (preferably) pressure rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers or product.

### **Envirodrum - Micro Matic Valve (110L)**

Store the original sealed Envirodrum in a cool, well ventilated area. **DO NOT** store for prolonged periods in direct sunlight. **DO NOT** tamper with the Micro Matic valve or the security seal. **DO NOT** contaminate the Envirodrum with water or any other foreign matter. After each use of the product please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase.

### **Refillable containers (1000L only)**

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

### **SAFETY DIRECTIONS**

Poisonous if inhaled. Will irritate the skin. Will damage the eyes. Repeated exposure may cause allergic disorders. **DO NOT** inhale vapour. Avoid contact with eyes and skin. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length chemical resistant gloves, goggles and half facepiece respirator. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length chemical resistant gloves.

If applying by hand, wear cotton overalls, over normal clothing, buttoned to the neck and wrist, and a washable hat, elbow-length chemical resistant gloves. If product on skin, immediately wash area with soap and water. If product in eyes wash out immediately with water. Wash hands after use. After each day's use, wash gloves, goggles, respirator and if rubber wash with detergent and warm water and contaminated clothing.

### **RE-ENTRY PERIOD**

**DO NOT** allow entry into treated areas for 8 days when applying to summer growing weeds, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

**DO NOT** allow entry into treated areas until the spray has dried when applying to winter growing weeds, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

**DO NOT APPLY USING SPRAYING EQUIPMENT CARRIED ON THE BACK OF THE USER.**

### **FIRST AID**

If swallowed, **DO NOT** induce vomiting. Give a glass of water. If poisoning occurs, consult a doctor or Poisons Information Centre (*Phone Australia 13 11 26; New Zealand 0800 764 766*). If in eyes, wash out immediately with water.

### Additional GHS Hazard & Precautionary Statements

●Combustible liquid; ●Repeated exposure may cause skin dryness or cracking; ●Harmful if swallowed; ●May cause respiratory irritation; ●May cause drowsiness or dizziness; ●May damage fertility or the unborn child; ●Toxic to aquatic life with long lasting effects; ●Avoid breathing fumes, mists or spray; ●**DO NOT** eat, drink or smoke when using this product; ●Use only outdoors or in a well-ventilated area; ●**IF ON SKIN (or hair):** Remove immediately all contaminated clothing. Rinse skin with water; ●**IF INHALED:** Remove victim to fresh air and keep at rest in a position comfortable for breathing; ●If exposed or concerned: Get medical advice; ●If skin irritation occurs: Get medical advice; ●If eye irritation persists: Get medical advice; ●Collect spillage; ●In case of fire, use carbon dioxide, dry chemical, foam, water fog. Alcohol resistant foam is the preferred firefighting medium but, if it is not available, normal foam can be used; ●Store locked up; ●Store in a dry place.

### **SAFETY DATA SHEET**

Additional information is listed in the Safety Data Sheet which can be obtained from your supplier or Imtrade Australia Pty Ltd website at [www.imtrade.com.au](http://www.imtrade.com.au)

### **CONDITIONS OF SALE**

Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever, or howsoever arising through negligence or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express conditions that the purchaser does not rely on Imtrade Australia's skill or judgement in purchasing or using the same and every person dealing with this product does so at his/her own risk

<b>UN 3082</b>	<b>ENVIRONMENTALLY HAZARDOUS SUBSTANCES, SOLID, N.O.S. (contains QUIZALOFOP-P-ETHYL)</b>
<b>IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE</b>	
<b>PG III</b>	<b>HAZCHEM •3Z</b>



**APVMA Approval No: 65807/106313**