

STORAGE AND DISPOSAL

Keep out of reach of children. Store in the closed, original container in a secure, cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. Protect from frost. Triple rinse container for disposal. Dispose of rinsate by adding it to the spray tank. **DO NOT** dispose of undiluted chemical on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any container management program site. The cap should not be replaced but may be taken separately. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers and product.

For Refillable containers (110L): Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

May irritate eyes. Avoid contact with eyes and skin. When opening the container and using the product wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length PVC gloves and face shield. If product comes in contact with skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia. 13 11 26).

Additional GHS Hazard & Precautionary Statements

•Suspected of causing cancer; •Very toxic to aquatic life with long lasting effects; •**DO NOT** get on clothing; •**IF SWALLOWED:** Rinse mouth. **DO NOT** induce vomiting; •If exposed or concerned: Get medical advice; •Collect spillage; •Not combustible. Use extinguishing media suited to burning materials. Alcohol resistant foam is the preferred firefighting medium but, if it is not available, normal foam can be used; •Store in a dry place.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from your supplier or Imtrade Australia Pty Ltd website at www.imtrade.com.au


READ SAFETY DIRECTIONS BEFORE OPENING OR USING



IMTRADE IPRODIONE 500SC FUNGICIDE

ACTIVE CONSTITUENT: 500 g/L IPRODIONE

GROUP **2** FUNGICIDE



For the control of certain fungal diseases in various crops and situations as specified in the Directions for Use table

IMPORTANT: READ THIS LABEL AND THE ATTACHED LEAFLET BEFORE THOROUGHLY OPENING OR USING THIS PRODUCT

Imtrade Australia Pty Ltd
ABN 13 090 151 134
17 Ocean Street, Kwinana WA 6167
Tel: 1800 171 799 Fax: 1800 171 788
Web: www.imtrade.com.au

**NET CONTENTS:
5L-1000L**

CONDITIONS OF SALE

Imtrade Australia Pty Ltd warrants that this product conforms to the chemical description on the label. The use of Imtrade Iprodione 500SC Fungicide being beyond the control of the manufacturer, Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Imtrade Australia Pty Ltd has any authority to add to or alter these conditions.

UN 3082	ENVIRONMENTALLY HAZARDOUS SUBSTANCES, LIQUID, N.O.S. (contains IPRODIONE)
IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE	
PG III	HAZCHEM •3Z

APVMA Approval No: 70433/63451



Batch No:
Date of Manufacture:

READ SAFETY DIRECTIONS BEFORE OPENING OR USING



IMTRADE IPRODIONE 500SC FUNGICIDE

ACTIVE CONSTITUENT: 500 g/L IPRODIONE

GROUP	2	FUNGICIDE
--------------	----------	------------------

For the control of certain fungal diseases
in various crops and situations as
specified in the Directions for Use table

**IMPORTANT: READ THIS LEAFLET THOROUGHLY
BEFORE OPENING OR USING THIS PRODUCT**

APVMA APPROVAL NO: 70433/63451

Imtrade Australia Pty Ltd
ABN 13 090 151 134
17 Ocean Street, Kwinana WA 6167
Tel: 1800 171 799 Fax: 1800 171 788
Web: www.imtrade.com.au

DIRECTIONS FOR USE:**Tree Crops/Vines**

RATE					CRITICAL COMMENTS
In the following table, all rates are given for dilute spraying. For concentrate spraying, refer to the Special Instructions for Tree Crops/Vines section					
CROP	DISEASE	STATE	RATE	WHP	Refer to the Special Instructions for Tree Crops/Vines section.
Almonds	Blossom Blight, Brown Rot (<i>Monilinia</i> spp., <i>Sclerotinia</i> spp.)	All states	50 mL/100L water	Nil	Apply first at full bloom and, if conditions are favourable for disease development, up to two subsequent applications can be made; at petal fall and up to four weeks after petal fall.
Boysenberries	Grey Mould (<i>Botrytis cinerea</i>)		100 mL/100L water	1 day	Spray at 10% blossom and full bloom. For fruit protection, apply at 2 to 3 weeks pre-harvest.
Grapes				7 days	Good crop hygiene will aid in the control of disease. This use is subject to a CropLife fungicide resistance management strategy: The number of consecutive applications and the total number of applications of Group 2 Fungicides permitted is limited. Refer to CropLife Australia Resistance Management Guidelines. See under “ General Instructions – Resistance Management ” for details of where these guidelines can be obtained.
Kiwifruit	Botrytis Blight (<i>Botrytis</i> spp.)	NSW, Vic, WA only			Apply the spray to vines every 10 to 14 days ensuring that all fruit is thoroughly wet. Apply 3 applications at 10 to 14 day intervals from 10% bloom to petal fall for protection of flowers and young fruit. Apply a further 2 applications of Imtrade Iprodione 500SC Fungicide to control late season Botrytis.

Tree Crops/Vines (continued)

RATE					CRITICAL COMMENTS
In the following table, all rates are given for dilute spraying. For concentrate spraying, refer to the Special Instructions for Tree Crops/Vines section					
CROP	DISEASE	STATE	RATE	WHP	For all uses in this table: Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. Refer to the Special Instructions for Tree Crops/Vines section.
Macadamias	Botrytis Blight (<i>Botrytis</i> spp.)	All states	50 mL/100L water	Nil	Apply as a thorough cover spray to flower racemes when they open. A follow up spray may be needed one week later if wet conditions persist during flowering. Remove nuts under trees prior to spraying.
Mandarins (non-bearing)	Alternaria Leaf Spot (Brown Spot) (<i>Alternaria alternate</i>)	QLD, WA, NT only	100 mL/100L water		Apply to non-bearing trees of Murcott variety monthly from first flush in spring until flushing ceases in the autumn. Reduce intervals to fortnightly during periods of wet weather.
Passionfruit	Alternata spot (Brown Spot) (<i>Alternaria</i> spp., <i>Alternaria passiflorae</i>)	QLD, NSW, WA, NT only	100 mL/100L water	7 days	This use is subject to an CropLife fungicide resistance management strategy: 1. Maintain a protective cover with protectant fungicide such as mancozeb. 2. Limit the use of Imtrade Iprodione 500SC Fungicide to strategic periods, i.e. before, during and after extended wet periods. 3. Always tank mix Imtrade Iprodione 500SC Fungicide with a protectant such as mancozeb. 4. DO NOT apply more than four Imtrade Iprodione 500SC Fungicide (or other Group 2 Fungicide) sprays in a season.
Raspberries	Grey Mould (<i>Botrytis cinerea</i>)	All States	100 mL/100L water	1 day	Spray at 10% blossom and full bloom. For fruit protection, apply at 2 to 3 weeks pre-harvest.

Tree Crops/Vines (continued)

RATE					CRITICAL COMMENTS
In the following table, all rates are given for dilute spraying. For concentrate spraying, refer to the Special Instructions for Tree Crops/Vines section					For all uses in this table: Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. Refer to the Special Instructions for Tree Crops/Vines section.
CROP	DISEASE	STATE	RATE	WHP	
Stone Fruit: Apricots, Cherries, Nectarines, Peaches, Plums	Orchard Spraying Blossom Blight (<i>Monilinia fructicola</i> , <i>Monilinia laxa</i>), Brown Rot (<i>Monilinia fructicola</i> , <i>Monilinia laxa</i>)	QLD, NSW, Vic, Tas, SA, WA only	50 to 75mL/ 100L water	Nil	Critical timings for control of blossom blight, are 10% blossom, full bloom and petal/shuck fall and for control of subsequent brown rot in fruit, spray at 3 weeks and 1 week pre-harvest. Use the higher rate under severe conditions of challenge, or for single applications of Imtrade Iprodione 500SC Fungicide in the spray program. This use is subject to an CropLife fungicide resistance management strategy: The number of consecutive applications and the total number of applications of Group 2 fungicides permitted is limited. Refer to CropLife Australia Resistance Management Guidelines. See under “ General Instructions – Resistance Management ” for details of where these guidelines can be obtained.
Youngberries	Grey Mould (<i>Botrytis cinerea</i>)	All States	100 mL/100L water	1 day	Spray at 10% blossom and full bloom. For fruit protection, apply at 2 to 3 weeks pre-harvest.

Berries:

(See **Tree Crops/Vines** for Boysenberries, Raspberries and Youngberries)

CROP	DISEASE	STATE	RATE	WHP	CRITICAL COMMENTS
Strawberries	Grey Mould (<i>Botrytis cinerea</i>)	All States	1.0L/ha where spray volume is less than 1000L/ha OR 100 mL/100L water where spray volume equals or exceeds 1000L/ha	1 day	This use is subject to an CropLife fungicide resistance management strategy: 1. Apply a program of protectant fungicides during flowering. If conditions favour disease development during this period use Imtrade Iprodione 500SC Fungicide. 2. DO NOT apply more than two successive sprays of Imtrade Iprodione 500SC Fungicide (or other Group 2 Fungicides).

FRUIT – POST-HARVEST DIPPING:

CROP	DISEASE	STATE	RATE	WHP	CRITICAL COMMENTS
Pome Fruit: Apples, Pears	Post-Harvest Dipping Storage Rots (<i>Penicillium</i> spp.) (<i>Botrytis</i> spp.), (<i>Gleosporium</i> spp.)	All States	100mL/100L water	Nil	<p>To minimise the development of post harvest rots handle fruit carefully to avoid fruit injury and dip promptly after harvest. Remove any infected fruit from the packing house immediately and destroy. When dipping, allow sufficient time to thoroughly wet the fruit. Top up the dip with 10 mL Imtrade Iprodione 500SC Fungicide in 100L of water.</p> <p>When using Imtrade Iprodione 500SC Fungicide as a post-harvest dip, DO NOT use a Group 2 Fungicide as a final pre-harvest spray. Rotation between different modes of action in post-harvest dips is advised.</p> <p>This use is subject to an CropLife fungicide resistance management strategy: See under “General Instructions – Resistance Management” for details of where these guidelines can be obtained.</p>
Stone Fruit: Apricots, Cherries, Nectarines, Peaches, Plums	Post-Harvest Dipping Brown Rot (<i>Monilinia fructicola</i> , <i>Monilinia laxa</i>), *Transit Rot (<i>Rhizopus</i> spp.)	NSW, QLD, VIC, TAS, SA, WA only			<p>To minimize the development of post-harvest rots, handle fruit carefully to avoid fruit injury and dip promptly after harvest. Remove any infected fruit from the packing house immediately and destroy.</p> <p>When dipping, allow sufficient time to thoroughly wet the fruit. Top-up dip with 200 mL Imtrade Iprodione 500SC Fungicide in 100L water. A non-ionic wetting agent should be used.</p> <p>*Transit Rot is suppressed at this rate.</p> <p>When using Imtrade Iprodione 500SC Fungicide as a post-harvest dip, DO NOT use a Group 2 Fungicide as the final pre-harvest field spray. Rotation between different modes of action in post-harvest dips is advised.</p> <p>This use is subject to an CropLife fungicide resistance management strategy: See under “General Instructions – Resistance Management” for details of where these guidelines can be obtained.</p>

VEGETABLES:

CROP	DISEASE	STATE	RATE	WHP	CRITICAL COMMENTS
Celery	Sclerotinia Rot (Pink Rot) (<i>Sclerotinia sclerotiorum</i>)	All States	1.0L/ha where spray volume is less than 1000L/ha OR 100 mL/100L water where spray volume equals or exceeds 1000L/ha	1 day	Commence spraying 1 to 2 weeks post-transplanting and then every 2 to 3 weeks. Use only five sprays.
Lettuces	Sclerotinia Rot (drop) (<i>Sclerotinia sclerotiorum</i> , <i>Sclerotinia minor</i>)			7 days	<p>Spray should be directed to the stems at ground level and to the underside of lower leaves.</p> <p>This use is subject to an CropLife fungicide resistance management strategy:</p> <ol style="list-style-type: none"> 1. Apply Imtrade Iprodione 500SC Fungicide as a seedling drench soon after emergence. 2. Apply a protectant fungicide as a high volume foliar spray before planting out, then Imtrade Iprodione 500SC Fungicide immediately after planting. 3. Maintain cover with protectant fungicide sprays at 7-10 day intervals. 4. If weather conditions favour Botrytis infection, tank mix the protectant with Imtrade Iprodione 500SC Fungicide. <p>DO NOT apply Imtrade Iprodione 500SC Fungicide (or other Group 2 Fungicides) more than four times per season, irrespective of the target disease.</p>
	Grey Mould (<i>Botrytis</i> spp.)	TAS, WA only			

VEGETABLES (Continued):

CROP	DISEASE	STATE	RATE	WHP	CRITICAL COMMENTS
Potatoes	Sclerotinia Rot (<i>Sclerotinia sclerotiorum</i>)	All States	500 mL to 1.0L/ha where spray volume is less than 1000L/ha	1 day	Apply 2 sprays, once immediately before and once immediately after hilling-up. For most effective treatment, concentrate the spray at the base of the stems and surrounding soil surface, where the fungus is active. Use the higher rate where disease is severe.
	Target Spot (Early Blight) (<i>Alternaria solani</i>)		OR 50-100 mL/100L water where spray volume equals or exceeds 1000L/ha		Ensure thorough coverage to the whole plant. Treatment is generally not required until after flowering. Use the higher rate where disease is severe. Limit the use of Imtrade Iprodione 500SC Fungicide to periods when conditions favour disease development. This use is subject to an CropLife fungicide resistance management strategy: The number of consecutive applications of Group 2 Fungicides permitted is limited. Refer to CropLife Australia Resistance Management Guidelines. See under “ General Instructions – Resistance Management ” for details of where these guidelines can be obtained.
	Hypocotyl Rot (Black Scurf) (<i>Rhizoctonia solani</i>)		400 mL/tonne seed material		Imtrade Iprodione 500SC Fungicide will protect emerging shoots from hypocotyl rot, improving overall germination. Imtrade Iprodione 500SC Fungicide may also reduce occurrence of black scurf on the harvested potatoes. Ensure good coverage of seed material and planting furrow. This can be achieved by applying Imtrade Iprodione 500SC Fungicide as a fine spray to the seed at the time of planting using spray equipment mounted on the planter, and nozzles located at three points on each planter row to ensure uniform coating of the seed. DO NOT plant into waterlogged soil. A minimum water volume of 80L/tonne seed should be used.

VEGETABLES (Continued):

CROP	DISEASE	STATE	RATE	WHP	CRITICAL COMMENTS
Tomatoes	Sclerotinia Rot (<i>Sclerotinia sclerotiorum</i>)	QLD, NSW, TAS, SA, WA only	1.0L/ha where spray volume is less than 1000L/ha OR 100 mL/100L water where spray volume equals or exceeds 1000L/ha	7 days	Spray at 14-day intervals from transplanting and throughout the period of disease pressure.
	Grey Mould (<i>Botrytis</i> spp.)	All States			Commence spraying 3 to 4 weeks after transplanting or at the onset of disease. Repeat treatment at 14-day intervals or when conditions favour spread of the disease, i.e. at trimming or de-leafing. This use is subject to an CropLife fungicide resistance management strategy: 1. Alternate or tank mix Imtrade Iprodione 500SC Fungicide with a protectant such as chlorothalonil. Avoid applying two Imtrade Iprodione 500SC Fungicide (or other Group 2 Fungicide) sprays in succession, unless tank mixed with a protectant. DO NOT apply more than four Imtrade Iprodione 500SC Fungicide (or other Group 2 Fungicide) sprays in a season.
	Target Spot (Early Blight) (<i>Alternaria solani</i>)	QLD, TAS, WA, NT only			Commence spraying 1 week post-transplanting. Use adequate water to give thorough coverage of the plants. Use high volume spray equipment. This use is subject to an CropLife fungicide resistance management strategy: 1. Limit the use of Imtrade Iprodione 500SC Fungicide to periods when conditions favour disease development. 2. DO NOT apply more than four Imtrade Iprodione 500SC Fungicide (or other Group 2 Fungicide) sprays in one season. Apply no more than two consecutive sprays of a Group 2 Fungicide.

FIELD CROPS:

CROP	DISEASE	STATE	RATE	WHP	CRITICAL COMMENTS
Lucerne	Lucerne Leaf Spot (<i>Stemphylium botryosum</i>)	QLD, WA only	250 mL to 500L/ha where spray volume is less than 1000L/ha	7 days	Spray every 10 to 14 days when cool, damp weather favours the disease. Use the higher rate under conditions of high disease pressure.
	Leptosphaerulina Leaf Spot (<i>Leptosphaerulina trifolii</i>)		OR 25 to 50 mL/100L water where spray volume equals or exceeds 1000L/ha		Apply in at least 300L water/ha every 10 to 14 days when cool, damp weather favours the disease. Use the higher rate under conditions of high disease pressure.
Peanuts	Sclerotinia Rot (<i>Sclerotinia sclerotiorum</i> , <i>Sclerotinia minor</i>)		1.0L/ha OR 220 mL/100L water (spot application)	12 days	Apply when disease first appears. Repeat if necessary. Use a high water volume to ensure good coverage of foliage and stem at ground level. DO NOT mix Imtrade Iprodione 500SC Fungicide with a foliar fungicide due to the different target positions on the plant.
Soybeans	Black Leaf Blight (<i>Arkoala nigra</i>)	NSW, WA only	1.0L/200 to 400L water/ha	7 days	If disease is present on leaves apply an initial spray at early pod set (pods approximately 5mm long). An additional spray 14 days later may be required if wet seasonal conditions prevail.

CROP	DISEASE	STATE	RATE	WHP	CRITICAL COMMENTS
Ornamentals	Grey Mould (<i>Botrytis cinerea</i>)	All States	100 mL/100L water	Nil	Spray at 14 day intervals commencing when the disease first becomes apparent and continuing until conditions no longer favour the disease. Spraying saintpaulia and poinsettia flowers may result in some petal scorch. Tepid water should be used and wet plants protected from direct sunlight. This use is subject to an CropLife fungicide resistance management strategy: DO NOT apply more than two consecutive sprays of a Group 2 Fungicide.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS - HARVEST

ALMONDS, MACADAMIAS, MANDARINS, POTATOES, STONE FRUIT:
BOYSENBERRIES, CELERY, RASPBERRIES, STRAWBERRIES, YOUNGBERRIES:
GRAPES, KIWIFRUIT, LETTUCE, TOMATOES, PASSIONFRUIT:
PEANUTS:
SOYBEANS:

NOT REQUIRED WHEN USED AS DIRECTED.
DO NOT HARVEST FOR 1 DAY AFTER APPLICATION.
DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION.
DO NOT HARVEST FOR 12 DAYS AFTER APPLICATION.
DO NOT HARVEST FOR 7 WEEKS AFTER APPLICATION.

WITHHOLDING PERIODS - GRAZING

LUCERNE:

DO NOT GRAZE OR CUT FOR STOCK FOOD WITHIN
7 DAYS OF TREATMENT.

FUNGICIDE RESISTANCE WARNING

GROUP	2	FUNGICIDE
--------------	----------	------------------

Imtrade Iprodione 500SC Fungicide is a member of the dicarboximide group of fungicides. For fungicide resistance management the product is a Group 2 Fungicide. Some naturally occurring individual fungi resistant to the product and other Group 2 Fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product or other Group 2 Fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Imtrade Australia Pty Limited accepts no liability for any losses that may result from the failure of this product to control resistant fungi.

Resistance Management

Rotate use of this product with as many different modes of action as possible to reduce the possibility of resistance development where alternatives are available.

Where specific resistance management strategies are established these are detailed in the CropLife Australia website (www.croplifeaustralia.org.au) or from your local agronomist.

Since the occurrence of resistant fungi is difficult to detect prior to use, Imtrade Australia Pty Ltd accepts no liability for any losses that may result from the failure of Imtrade Iprodione 500SC Fungicide to control resistant fungi.

EXPORT OF TREATED PRODUCE

Growers should note that suitable MRLs or import tolerances may not be established in all markets for produce treated with Imtrade Iprodione 500SC Fungicide. If you are growing produce for export, please check with Imtrade Australia Pty Ltd for the latest information on MRLs and import tolerances BEFORE using Imtrade Iprodione 500SC Fungicide.

MIXING

Note: Imtrade Iprodione 500SC Fungicide may be unstable in conditions where the pH is 7 or higher. It is therefore essential to check the pH of the spray solution before adding Imtrade Iprodione 500SC Fungicide. A suitable registered buffering agent may have to be added to bring the pH down below 7.

Add the required amount of Imtrade Iprodione 500SC Fungicide to the spray tank containing half the required volume of water. Mix thoroughly and make up to the required volume.

APPLICATION

Good disease control requires even, thorough coverage of the target area. Application should be made using appropriate spray equipment and sufficient water to provide adequate penetration and coverage. Equipment settings and water volume may need to vary, depending on the growth stage of the crop.

Special Instructions for Tree Crops/Vines

Dilute Spraying

Use a sprayer designed to apply high spray volumes, up to the point of run-off and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient spray solution to cover the crop to the point of run-off. Avoid excessive run-off. The required spray volume to achieve point of run off may be determined by applying different test volumes, using different settings on the sprayer, or from industry guidelines or other expert advice. Add the amount of product specified in the Directions for Use Table for each 100L of water. Spray to the point of run-off. The required dilute spray volume to achieve point of run-off will change and the sprayer set up and operation may also need to be changed, as the crop grows.

Concentrate Spraying

Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies spray volumes less than those required to reach the point of run-off) and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen spray volume. Determine an appropriate dilute spray volume (See *Dilute Spraying* above) for the crop canopy. This is needed to calculate the concentrate mixing rate. The mixing rate for concentrate spraying can then be calculated in the following way:

EXAMPLE ONLY

1. Dilute spray volume as determine above: for example 1500 L/ha
2. Your chosen concentrate spray volume: for example 500 L/ha
3. The concentration factor in this example is: $3x$ (i.e. $1500L \div 500L = 3$)
4. If the dilute label rate is 10mL/100L, then the concentrate rate becomes 3×10 , that is 30 mL of product per 100L water for concentrate spraying.

The chosen spray volume, amount of product per 100L of water, and the sprayer set up and operation may need to be changed as the crop grows. For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

COMPATIBILITY

Imtrade Iprodione 500SC Fungicide is compatible with the following products:

*Aliette WG# (see NOTE below), Bugmaster Flowable#, calcium chloride, chlorpyrifos 500 EC, copper oxychloride, dimethoate, Dithane, Dithane M-45#, endosulfan, Kelthane EC#, maldison, metalaxyl, methomyl or parathion-methyl.

When tank mixing products the order of mixing is determined by formulation type. As a guide the following mixing sequence is recommended:

1. Wettable powders
2. Suspension concentrates
3. Water Dispersible Granules
4. Suspo-emulsions (e.g. Imtrade Iprodione 500SC Fungicide)
5. Soluble powders
6. Solutions
7. Emulsifiable concentrates
8. Soluble concentrates
9. Wetting agents and oils

With any mixture, thoroughly agitate immediately before applying. It is not recommended to mix this product with more than one of the above chemicals in the tank. The use of a surfactant or spray oil is not recommended with Imtrade Iprodione 500SC Fungicide as it may result in crop damage to sensitive plants. **DO NOT** mix with fertilisers.

NOTE: Mixing Imtrade Iprodione 500SC Fungicide with Aliette WG# may result in some settling out.

As formulations of other manufacturers' products are beyond the control of Imtrade Australia Pty Limited, all mixtures should be tested prior to mixing commercial quantities.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Toxic to aquatic organisms.

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

STORAGE AND DISPOSAL

Keep out of reach of children. Store in the closed, original container in a secure, cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. Protect from frost. Triple rinse container for disposal. Dispose of rinsate by adding it to the spray tank. **DO NOT** dispose of undiluted chemical on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any container management program site. The cap should not be replaced but may be taken separately. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers and product.

For Refillable containers (110L): Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

May irritate eyes. Avoid contact with eyes and skin. When opening the container and using the product wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length PVC gloves and face shield. If product comes in contact with skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia. 13 11 26).

Additional GHS Hazard & Precautionary Statements

●Suspected of causing cancer; ●Very toxic to aquatic life with long lasting effects; ●**DO NOT** get on clothing; ●**IF SWALLOWED:** Rinse mouth. **DO NOT** induce vomiting;; ●If exposed or concerned: Get medical advice; ●Collect spillage; ●Not combustible. Use extinguishing media suited to burning materials. Alcohol resistant foam is the preferred firefighting medium but, if it is not available, normal foam can be used; ●Store in a dry place.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from your supplier or Imtrade Australia Pty Ltd website at www.imtrade.com.au

CONDITIONS OF SALE

Imtrade Australia Pty Ltd warrants that this product conforms to the chemical description on the label. The use of Imtrade Iprodione 500SC Fungicide being beyond the control of the manufacturer, Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Imtrade Australia Pty Ltd has any authority to add to or alter these conditions

#Other Company Registered Products

UN 3082	ENVIRONMENTALLY HAZARDOUS SUBSTANCES, LIQUID, N.O.S. (contains IPRDIONE)
IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE	
PG III	HAZCHEM •3Z

APVMA Approval No: 70433/63451

