

## GENERAL INSTRUCTIONS

Imtrade Metakill Snail & Slug Bait kills snails and slugs by irreversibly destroying the mucus producing cells that are essential for movement. Where pressure is heavy, it may be necessary to repeat the application. Best results are achieved by application during mild, damp weather when slugs and snails are most active, but not when rain is imminent. Snails need to be active to come into contact with the bait.

Scatter baits evenly using an appropriate applicator by ground or air, or direct drill with seed for slug control. **DO NOT** place in heaps. Clean up any spilled baits, including after filling applicator.

## TIMING

Apply baits when crops are most vulnerable and when the pest is evident and active. For broadacre crops, start when environmental conditions trigger snail activity in autumn and before significant egg laying.

For other crops, time the application to protect the sensitive growth stage. For seedlings best results may be achieved by applying 5-10 days prior to planting. See your distributor for further advice.

## PRECAUTIONS

**DO NOT** use human flaggers/markers unless they are protected by engineering controls.

## PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

**DO NOT** allow chemical containers or bait to get into drains, sewers, streams or ponds.

## PROTECTION OF LIVESTOCK

**DO NOT** place baits in locations which are accessible to domestic animals, livestock or birds.

## STORAGE AND DISPOSAL

Store in the closed, original container in a cool, dry place out of direct sunlight and out of the reach of children, animals, poultry or ducks.

- Keep away from domestic pets.
- Keep away from dogs. Dogs find this bait attractive and it may kill them.
- If pets are poisoned, contact a Veterinary Surgeon.

Shake container contents until the container is empty. **DO NOT** dispose of chemicals on site. Puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

# POISON

**KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**



# IMTRADE METAKILL SNAIL & SLUG BAIT

**ACTIVE CONSTITUENT: 50 g/kg METALDEHYDE**



**For the control of Snails and Slugs  
in agricultural and horticultural  
crops, vineyards and other  
agricultural, industrial and  
commercial situations where  
snails and slugs congregate  
as specified in the  
Directions for Use table**

**IMPORTANT: READ THIS LABEL  
AND THE ATTACHED LEAFLET  
THOROUGHLY BEFORE OPENING  
OR USING THIS PRODUCT**

**NET CONTENTS:  
500g-1000kg**

Imtrade Australia Pty Ltd  
ABN 13 090 151 134  
17 Ocean Street, Kwinana WA 6167  
Tel: 1800 171 799 Fax: 1800 171 788  
Web: [www.imtrade.com.au](http://www.imtrade.com.au)

## SAFETY DIRECTIONS

Harmful if swallowed. Harmful if inhaled. **DO NOT** inhale dust. Wash hands after use. When opening the container, preparing the product for use and if applying by hand, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow length rubber gloves and disposable dust mask covering mouth and nose. After each day's use, wash gloves and contaminated clothing.

## FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26; New Zealand 0800 764 766).

### Additional GHS Hazard & Precautionary Statements

• **DO NOT** get in eyes, on skin or on clothing; • **DO NOT** eat, drink or smoke when using this product; • Use only outdoors or in a well-ventilated area; • **IF SWALLOWED:** Rinse mouth. Do **NOT** induce vomiting; • **IF INHALED:** Remove victim to fresh air and keep at rest in a position comfortable for breathing; • In case of fire, use carbon dioxide, dry chemical, foam, water fog.

## SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from your supplier or Imtrade Australia Pty Ltd website at [www.imtrade.com.au](http://www.imtrade.com.au)

## CONDITIONS OF SALE

Imtrade Australia Pty Ltd warrants that this product conforms to the chemical description on the label. The use of Imtrade Metakill Snail & Slug Bait being beyond the control of the manufacturer, Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Imtrade Australia Pty Ltd has any authority to add to or alter these conditions.

**APVMA Approval No: 64990/117488**

**Batch No:**

**Date of Manufacture:**

**POISON**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**



# **IMTRADE METAKILL SNAIL & SLUG BAIT**

**ACTIVE CONSTITUENT: 50 g/kg METALDEHYDE**

**For the control of Snails and Slugs in agricultural and horticultural crops, vineyards and other agricultural, industrial and commercial situations where snails and slugs congregate as specified in the Directions for Use table**

**IMPORTANT: READ THIS LEAFLET THOROUGHLY  
BEFORE OPENING OR USING THIS PRODUCT**

**APVMA Approval No. 64990/117488**

Imtrade Australia Pty Ltd  
ABN 13 090 151 134  
17 Ocean Street, Kwinana WA 6167  
Tel: 1800 171 799 Fax: 1800 171 788  
Web: [www.imtrade.com.au](http://www.imtrade.com.au)

**DIRECTIONS FOR USE****RESTRAINTS:****DO NOT** apply to windrowed crops.**DO NOT** apply to edible plant parts or to crops where baits may be collected with harvested commodities.**DO NOT** heap pellets. Clean up spilled pellets immediately.

<b>CROPS/SITUATIONS</b>	<b>PEST</b>	<b>RATE</b>	<b>HOW TO APPLY</b>
<b>Agricultural Crops</b> Cereals, Oilseeds (inc. Poppies), Pulses	Snails and Slugs (except Small Brown Snails)	5-8 kg/ha	Broadcast evenly over the surface of the ground where snails and slugs are active, or incorporate with seeding equipment when direct drilling. Use low rate only when snail density is low. Use higher rate for moderate to high infestations
	Small Brown Snails	10 kg/ha	
<b>Horticultural Crops</b> Fruit (inc. Grapevines) Vegetables			For best results on small brown snails, use 10 kg/ha rate in a full integrated Program with careful monitoring.
<b>Direct Drilled Pasture</b>	Slugs	5-8 kg/ha	Incorporate with seed when direct drilling. Use low rate only when slug density is low. Use higher rate for moderate to high infestations
Other Agricultural, Industrial & Commercial situations	Snails and Slugs (except Small Brown Snails)	5-8 kg/ha	Broadcast evenly over the surface of the ground where snails and slugs are active, or incorporate with seeding equipment when direct drilling. Use low rate only when snail density is low. Use higher rate for moderate to high infestations
	Small Brown Snail	10 kg/ha	
Seedlings and mature crops of vegetables and flowers	Snails, Slugs	4-8 kg/ha	Water area well. Scatter pellets on garden beds and in shady situations around the base of plants.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

## **WITHHOLDING PERIOD**

### **HARVEST**

**HORTICULTURAL CROPS, VINEYARDS: DO NOT APPLY LATER THAN 7 DAYS (1 WEEK) BEFORE HARVEST.**

**AGRICULTURAL CROPS: DO NOT APPLY LATER THAN 8 WEEKS BEFORE HARVEST.**

### **GRAZING**

**DO NOT ALLOW STOCK OR POULTRY TO GRAZE TREATED CROPS, VINEYARDS OR OTHER AREAS.**

**DO NOT ALLOW STOCK TO GRAZE TREATED PASTURE FOR 8 WEEKS.**

## **GENERAL INSTRUCTIONS**

Imtrade Metakill Snail & Slug Bait kills snails and slugs by irreversibly destroying the mucus producing cells that are essential for movement. Where pressure is heavy, it may be necessary to repeat the application. Best results are achieved by application during mild, damp weather when slugs and snails are most active, but not when rain is imminent. Snails need to be active to come into contact with the bait.

Scatter baits evenly using an appropriate applicator by ground or air, or direct drill with seed for slug control. **DO NOT** place in heaps. Clean up any spilled baits, including after filling applicator.

## **TIMING**

Apply baits when crops are most vulnerable and when the pest is evident and active. For broadacre crops, start when environmental conditions trigger snail activity in autumn and before significant egg laying.

For other crops, time the application to protect the sensitive growth stage. For seedlings best results may be achieved by applying 5-10 days prior to planting. See your distributor for further advice.

## **PRECAUTIONS**

**DO NOT** use human flaggers/markers unless they are protected by engineering controls.

## **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

**DO NOT** allow chemical containers or bait to get into drains, sewers, streams or ponds.

## **PROTECTION OF LIVESTOCK**

**DO NOT** place baits in locations which are accessible to domestic animals, livestock or birds.

## **STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, dry place out of direct sunlight and out of the reach of children, animals, poultry or ducks.

- Keep away from domestic pets.
- Keep away from dogs. Dogs find this bait attractive and it may kill them.
- If pets are poisoned, contact a Veterinary Surgeon.

Shake container contents until the container is empty. **DO NOT** dispose of chemicals on site. Puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

## **SAFETY DIRECTIONS**

Harmful if swallowed. Harmful if inhaled. **DO NOT** inhale dust. Wash hands after use. When opening the container, preparing the product for use and if applying by hand, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow length rubber gloves and disposable dust mask covering mouth and nose. After each day's use, wash gloves and contaminated clothing.

## **FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre (*Phone Australia 13 11 26; New Zealand 0800 764 766*).

### **Additional GHS Hazard & Precautionary Statements**

•**DO NOT** get in eyes, on skin or on clothing; •**DO NOT** eat, drink or smoke when using this product; •Use only outdoors or in a well-ventilated area; •**IF SWALLOWED:** Rinse mouth. Do **NOT** induce vomiting; •**IF INHALED:** Remove victim to fresh air and keep at rest in a position comfortable for breathing; •In case of fire, use carbon dioxide, dry chemical, foam, water fog.

### **SAFETY DATA SHEET**

Additional information is listed in the Safety Data Sheet, which can be obtained from your supplier or Imtrade Australia Pty Ltd website at [www.imtrade.com.au](http://www.imtrade.com.au)

### **CONDITIONS OF SALE**

Imtrade Australia Pty Ltd warrants that this product conforms to the chemical description on the label. The use of Imtrade Metakill Snail & Slug Bait being beyond the control of the manufacturer, Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Imtrade Australia Pty Ltd has any authority to add to or alter these conditions.

**APVMA Approval No: 64990/117488**