

#### STORAGE AND DISPOSAL

Keep out of reach of children. Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. **DO NOT** store for prolonged periods in direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers.

Triple rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers or product.

#### SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. **DO NOT** inhale dust or spray mist. When opening the container, preparing spray, wear cotton overalls, over normal clothing, buttoned to the neck and wrist (or equivalent clothing) and chemical-resistant gloves and a half face respirator. If applying by low pressure hand wand and high pressure hand wand, wear cotton overalls, over normal clothing, buttoned to the neck and wrist (or equivalent clothing) and chemical-resistant gloves and a half face respirator. If product on skin, immediately wash area with soap and water. Wash hands after use. After each day's use wash gloves, respirator and if rubber wash with detergent and warm water and contaminated clothing.

#### FIRST AID

First aid is not generally required. If in doubt, contact a Poisons Information Centre (Phone Australia 13 11 26; New Zealand 0800 764 766) or a doctor.

Refer to attached Leaflet for additional GHS  
Hazard & Precautionary Statements

#### SAFETY DATA SHEET

For further information refer to the Safety Data Sheet, which can be obtained from your supplier or Imtrade Australia Pty Ltd website at [www.imtrade.com.au](http://www.imtrade.com.au)

#### CONDITIONS OF SALE

Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Imtrade Australia Pty Ltd has any authority to add to or alter these conditions.

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

# IMTRADE LINURON 800 WG HERBICIDE

ACTIVE CONSTITUENT: 800g/kg LINURON

**GROUP 5 HERBICIDE**

For the selective weed control in Carrots & Parsnips, Celeriac, Celery, Coriander, Jojoba, Onions (including Bulb Onions, Leek, Shallots and Spring Onions), Oil Tea Tree, Parsnips, Potatoes, Pyrethrum, Soybeans, Sweetcorn & Maize, Wheat, Oats & Barley as specified in the Directions for Use table.

**Important: Read the attached Leaflet before use.**

**GROUP  
5**

**FORM  
WG**

**CONTENTS  
1kg-20kg**

<b>UN 3077</b>	ENVIRONMENTALLY HAZARDOUS SUBSTANCES, SOLID, N.O.S. (contains LINURON)
In a Transport Emergency Dial 000 Police or Fire Brigade	
PG III	HAZCHEM 2Z

APVMA Approval No: 81574/131931



Batch No.  
DOM:

**IMTRADE**  
CropScience

17 Ocean St, Kwinana Beach WA 6167  
08 9419 0333 | [www.imtrade.com.au](http://www.imtrade.com.au)

**KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**



# **IMTRADE LINURON 800 WG HERBICIDE**

**ACTIVE CONSTITUENT: 800g/kg LINURON**

<b>GROUP</b>	<b>5</b>	<b>HERBICIDE</b>
--------------	----------	------------------

**For the selective weed control in Carrots & Parsnips, Celeriac, Celery, Coriander, Jojoba, Onions (including Bulb Onions, Leek, Shallots and Spring Onions), Oil Tea Tree, Parsnips, Potatoes, Pyrethrum, Soybeans, Sweetcorn & Maize, Wheat, Oats & Barley as specified in the Directions for Use table.**

**IMPORTANT: READ THIS LEAFLET THOROUGHLY  
BEFORE OPENING OR USING THIS PRODUCT**

**APVMA Approval No: 81574/131931**

Imtrade Australia Pty Ltd  
ABN 13 090 151 134  
17 Ocean Street, Kwinana, Western Australia 6167  
Tel: 1800 171 799 Fax: 1800 171 788  
Web: [www.imtrade.com.au](http://www.imtrade.com.au)

## DIRECTIONS FOR USE

### RESTRAINTS:

Sufficient moisture, either by irrigation or rainfall is required within 3 – 4 days of application to carry the product into the soil profile.

**DO NOT** apply using low pressure hand wand application, except for application to coriander seed crops and onions.

**DO NOT** apply using spraying equipment carried on the back of the user.

**DO NOT** apply by air, except for application to Wheat, Oats and Barley.

Crop	Weeds Controlled	State	Rate/ha		Critical Comments	
			Light Soils	Heavy Soils		
Carrots & Parsnips	<b>Annual Broadleaved Weeds and Grasses such as:</b> Amaranth, Blackberry Nightshade, Capeweed, Chickweed, Common Sowthistle, Corn Spurry, Deadnettle, Doublegee/ Three-cornered Jack (Spiny Emex), Fat-hen, Hedge Mustards, Heliotrope, Marsh Mallow, Paterson's Curse, Pigweed, Potato Weed, Redshank, Scarlet Pimpernel, Staggerweed, Stinging Nettle, Summer Grass (Summer Grass not controlled in Qld or November to February in other states), Swinecress, Variegated Thistle, Wild Mustard, Wild Radish, Wild Turnip, Winter Grass	All States	1.4kg	2.8kg	<b>Pre-emergence:</b> Apply immediately after planting but before crop emerges. <b>DO NOT</b> use on light sandy soils, low in clay or organic matter.	<b>Carrots and Parsnips:</b> Heavy rains after application may cause crop damage.
Carrots			690g	1.4kg	<b>Post-emergence:</b> <b>DO NOT</b> apply before carrots have reached the 4 true leaf stage, weeds no more than 7cm tall, or within 5 days of any application of any emulsifiable pesticide or wetting agent as damage to crop may result. On soils high in organic matter, use post-emergence rather than pre-emergence.	<b>DO NOT</b> replant treated areas to other crops within 3 months of application.

Crop	Weeds Controlled	State	Rate/ha		Critical Comments
			Light Soils	Heavy Soils	
<b>Celeriac</b> ( <i>Apium graveolens</i> )	<b>Annual Broadleaved Weeds and Grasses such as:</b> Amaranth, Blackberry, Nightshade, Capeweed, Chickweed, Common Sowthistle, Corn Spurry, Deadnettle, Doublegee/ Three-cornered Jack (Spiny Emex), Fat-hen, Hedge Mustards, Heliotrope, Marsh Mallow, Paterson's Curse, Pigweed, Potato Weed, Redshank, Scarlet Pimpernel, Staggerweed, Stinging Nettle, Summer Grass (Summer Grass not controlled in Qld or November to February in other states), Swinecress, Variegated Thistle, Wild Mustard, Wild Radish, Wild Turnip, Winter Grass	All States	700g	1.4kg	<b>Post-emergence at 2-4 true leaf stage:</b> Apply once transplants are established but before the crop reaches 20cm tall. <b>DO NOT</b> apply after the 5-leaf stage of the crop. <b>DO NOT</b> apply more than one (1) application per crop. <b>DO NOT</b> add wetting agents.  Use lower rate for light soils and higher rate for heavy clay soils. <b>DO NOT</b> apply if temperature exceeds 30°C.
<b>Celery</b> ( <i>Apium graveolens</i> )			625g		<b>Post-emergence at 2-4 true leaf stage:</b> Apply once transplants are established but before the crop reaches 20cm tall. <b>DO NOT</b> apply after the 5-leaf stage of the crop. <b>DO NOT</b> apply if temperature exceeds 30°C.  The sensitivity of the crop to be treated has not been fully evaluated. Treat a small area to ascertain the reaction before treating the whole crop.
<b>Coriander Seed Crops</b>	Amsinckia, Capeweed, Charlock, Corn Gromwell (Sheepweed, White Ironweed), Deadnettle, Doublegee/Three-cornered Jack (Spiny Emex), Fat-hen, Hedge Mustards, Indian Hedge Mustard, Saffron Thistle, Stinging Nettle, Wild Radish, Wild Turnip	SA, VIC, NSW, Tas only	470g	625g	<b>DO NOT</b> use Imtrade Linuron WDG Herbicide on fresh coriander production. Use lower rate quoted on light soils and increase up to the higher rate for heavy clay soils or soils high in organic matter.  <b>Pre-emergence:</b> Apply immediately after sowing.  <b>Post-emergence:</b> Apply from 2-leaf onwards. Higher rates give longer residual control. Wet soil and/ or rain is a benefit soon after application.

Crop	Weeds Controlled	State	Rate/ha		Critical Comments
			Light Soils	Heavy Soils	
Jojoba ( <i>Simmondsia chinensis</i> )	<b>Annual Broadleaved Weeds and Grasses such as:</b> Amaranth, Apple-of-Peru, Blackberry Nightshade, Capeweed, Chickweed, Clover, Common Sowthistle, Corn Spurry, Deadnettle, Doublegee/ Three-cornered Jack (Spiny Emex), Fat-hen, Hedge Mustards, Pigweed, Potato Weed, Redshank, Shepherd's Purse, Staggerweed, Stinging Nettle, Wild Mustard, Wild Radish, Wild Turnip, Winter Grass, Winter Weed, Wireweed	All States	700g	2.5kg	<p>Apply one (1) treatment only in sufficient volume to adequately cover target weeds.</p> <p>Avoid spraying of crop. Nozzle shields and targeted application techniques are recommended to minimise overspray and/or incidental spray drift onto crops.</p> <p><b>DO NOT</b> allow spray to contact green bark, stems, suckers or foliage.</p> <p><b>DO NOT</b> apply spray near trees less than 3 years old unless they are effectively shielded from spray and spray drift.</p> <p><b>DO NOT</b> add wetting agents or surfactants.</p> <p>Use lower application rate on lighter soils. Increase the rate on heavier soils. <b>DO NOT</b> apply on lighter sandy soils, low in clay or organic matter without first assessing crop safety.</p> <p>Rotate herbicide Mode of Action groups within and across growing seasons.</p> <p>The sensitivity of the crop to be treated has not been fully evaluated. Treat a small area to ascertain the reaction before treating the whole crop.</p>

Crop	Weeds Controlled	State	Rate/ha		Critical Comments
			Light Soils	Heavy Soils	
Onions including Bulb Onions, Leek, Shallots and Spring Onions	<b>Annual Broadleaved Weeds and Grasses such as:</b> Black Bindweed, Capeweed, Charlock, Chickweed, Clover, Common Sowthistle, Corn Spurry, Deadnettle, Fat-hen, Field Cress, Hedge Mustards, Lesser Swinecress, Medics, Prickly Sowthistle, Shepherd's Purse, Spear Thistle, Stinging Nettle, Wild Mustard, Wild Radish, Wild Turnip	QLD, NSW, VIC, SA, WA only	190g	345g	<b>Post-emergence only:</b> Apply when onions are 15cm or more in height. Re-treat when necessary. <b>Leeks</b> Apply as required (5-10 weeks after transplanting is common), over the top of the crop, using a boomspray. Can be applied over seeded crops at the same stage. Apply in 80-110L water/ha. <b>DO NOT</b> apply more than three (3) applications per crop. If weed control is less than adequate and resistance is suspected, the second application should not be applied. <b>Shallots and Spring Onions</b> Apply after 3 true leaf stage, when onions are 15cm high. Foliage must be waxy. Apply to foliage by aerial or ground boomspray or equivalent technology. Use a spray volume of 300L/ha for ground application and 20-30L/ha for aerial application. <b>DO NOT</b> apply more than two (2) applications per crop. <b>DO NOT</b> apply less than 14-21 days after initial treatment.
		Tas only	440g to 625g		<b>Post-emergence only:</b> Apply lowest rate when onion plants are 2-3 leaf stage and high rate from 3-leaf stage onwards. Up to three applications may be used.
Oil Tea Tree ( <i>Melaleuca alternifolia</i> )	Colombian Waxweed	All States	625g		Apply post-harvest through to canopy closure by directed boomspray application in a spray volume of 120L/ha. <b>DO NOT</b> apply more than two (2) applications per season. <b>DO NOT</b> add wetting agent or surfactant.

Crop	Weeds Controlled	State	Rate/ha		Critical Comments
			Light Soils	Heavy Soils	
Oil Tea Tree ( <i>Melaleuca alternifolia</i> ) (Continued)	Seedling Sedges	All States	690g – 1.4kg		Apply one (1) application to Seedling Sedges as an over-the-top of Oil Tea Tree seedlings spray. Use approximately 80-110L of water per hectare with a boomsprayer, or 40L per hectare when applying by helicopter or 20-30L per hectare when applying by fixed wing aircraft. <b>DO NOT</b> add wetting agent or surfactant.
Parsnips	<b>Annual Broadleaved Weeds and Grasses such as:</b> Amaranth, Blackberry Nightshade, Capeweed, Chickweed, Common Sowthistle, Corn Spurry, Deadnettle, Doublegee/ Three-cornered Jack (Spiny Emex), Fat-hen, Hedge Mustards, Heliotrope, Marsh Mallow, Paterson's Curse, Pigweed, Potato Weed, Redshank, Scarlet Pimpernel, Staggerweed, Stinging Nettle, Summer Grass (Summer Grass not controlled in Qld or November to February in other states), Swinecress, Variegated Thistle, Wild Mustard, Wild Radish, Wild Turnip, Winter Grass	All States	125 - 440g		Early post emergence from the 1-2 true leaf stage. Apply alone or as a tank mix with other approved herbicides. Apply as a foliar application by boomspray in spray volume of 500L/ha. <b>DO NOT</b> apply more than one (1) application per crop. Use low rate on lighter soils or early crop development stage and higher rates on heavier soils or later crop development stages. <b>DO NOT</b> apply if temperature is above 30°C. <b>DO NOT</b> add wetting agents, surfactants or crop oils. <b>DO NOT</b> apply to weeds, under severe stress due to, for example, very dry or windy conditions, or diseased crops. <b>DO NOT</b> allow mixture to stand in vat for prolonged periods, as it is difficult to re-suspend and may lose effectiveness. <b>DO NOT</b> apply with misting machines. Sufficient moisture, by either irrigation or rainfall is required within 3-4 days of application to carry the product into the soil profile. The sensitivity of the crop to be treated has not been fully evaluated. Treat a small area to ascertain the reaction before treating the whole crop.

Crop	Weeds Controlled	State	Rate/ha		Critical Comments
			Light Soils	Heavy Soils	
Potatoes	<b>Annual Broadleaved Weeds and Grasses such as:</b> Amaranth, Apple-of-Peru, Blackberry Nightshade, Capeweed, Chickweed, Clover, Common Sowthistle, Corn Spurry, Deadnettle, Doublegee/ Three-cornered Jack (Spiny Emex), Fat-hen, Hedge Mustards, Pigweed, Potato Weed, Redshank, Shepherd's Purse, Staggerweed, Stinging Nettle, Wild Mustard, Wild Radish, Wild Turnip, Winter Grass, Winter Weed, Wireweed	All States	1.4kg	2.8kg	<b>Pre-emergence:</b> Apply after planting but before emergence of crop. Incorporate by irrigation or rainfall as soon as possible after application. <b>DO NOT</b> cultivate after application. Use higher rate during summer months.
<b>Pyrethrum</b> <i>(Tanacetum cinerariifolium)</i>	Chickweed, Lesser Swinecress, Shepherds Purse, Volunteer Potato, Wild Radish, Wild Turnip	All States	110 - 180g		Apply from 4 true leaf stage of seedling pyrethrum. Avoid application to stressed plants or applications during the hottest part of the day. The <i>sensitivity of pyrethrum</i> has not been fully evaluated. Treat a small area of crop and monitor plant health closely over the following days/weeks. Discontinue applications when any adverse symptoms post-treatment are observed.



Crop	Weeds Controlled	State	Rate/ha		Critical Comments
			Light Soils	Heavy Soils	
Soybeans	<b>Annual Broadleaved Weeds and Grasses such as:</b> Amaranth, Apple-of-Peru, Barmyard Grass, Bathurst Burr, Black Bindweed, Blackberry Nightshade, Common Sowthistle, Fat-hen, Hedge Mustards, Noogoora Burr, Pigweed, Potato Weed, Redshank, Stinking Roger, Summer Grass, Turnip Weed, Wild Radish	QLD, NSW, VIC, WA only	1.4kg	3.4kg	<b>Pre-emergence:</b> Apply within 2 – 3 days after sowing. Sow beans to a depth of at least 2.5cm. Heavy rains after application may cause crop damage. <b>DO NOT</b> replant treated area to other crops within 3 months of application.
Sweetcorn & Maize	Refer to weeds under Soybeans plus Stinging Nettle, Wild Mustard, Wild Turnip. Apply where grasses are not a problem.	All States	—	1.4kg to 2.2kg	<b>Pre-emergence:</b> Apply as a band of broadcast treatment immediately after planting. <b>DO NOT</b> use on light sandy soils low in clay or organic matter. Heavy rains after application may cause crop damage.
	Refer to weeds above. Apply where Barnyard Grass or Summer Grass are a problem.			2.8kg to 3.4kg	
	Refer to weeds under Soybeans plus Stinging Nettle, Wild Mustard, Wild Turnip. Apply where grasses are not a problem.		690g to 1.4kg	1.4kg to 2.8kg	<b>Post-emergence:</b> Apply as directed band spray to each side row when crop is 30 – 40cm high. <b>DO NOT</b> use misting machines. Avoid spray contact with crop foliage as injury may result. <b>DO NOT</b> replant treated areas to other crops within 3 months of application. If weeds are present add a wetting agent at the use rate on its label.
	Refer to weeds above. Apply where Barnyard Grass or Summer Grass are a problem.		690g to 2.2kg	1.4kg to 3.4kg	

Crop	Weeds Controlled	State	Rate/ha		Critical Comments
			Light Soils	Heavy Soils	
<b>Wheat, Oats &amp; Barley</b>	Capeweed, Charlock, Corn Gromwell (Sheepweed, White Ironweed), Deadnettle, Doublegee/Three-cornered Jack (Spiny Emex), Fat-hen, Hedge Mustards, Indian Hedge Mustard, Stinging Nettle, Wild Radish, Wild Turnip	All States	315g		<b>All Weeds:</b> For best results apply early. <b>DO NOT</b> apply Imtrade Linuron 800 WDG Herbicide if weeds are greater than 15cm in height or diameter, or crop is beyond early tillering or crop and weeds are under drought stress. Apply when weeds are less than 5cm in height or diameter, crop is in 2-4 leaf stage and soil is moist enough for good growth of crop and weed seedlings.
	As above plus Amsinckia and Safron Thistle		530g		Apply if weeds are 5-15cm in height or diameter and crop is in 4-leaf to early tillering stage.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

#### **WITHHOLDING PERIOD**

***Celeriac:* DO NOT HARVEST FOR 4 WEEKS AFTER APPLICATION**

***Coriander, Carrots, Leeks and Parsnips* - DO NOT HARVEST FOR 10 WEEKS AFTER APPLICATION.**

***Pyrethrum (Tanacetum cinerariifolium)* – DO NOT HARVEST FOR 25 WEEKS AFTER APPLICATION.**

***Celery, Onions, Potatoes, Wheat, Barley, Oats, Maize, Shallots, Spring Onions Sweetcorn, Soybeans* - NOT REQUIRED WHEN USED AS DIRECTED.**

***Oil Tea Tree* – DO NOT GRAZE TREATED AREAS OR CUT FOR STOCKFEED FOR 10 WEEKS AFTER APPLICATION.**

***Jojoba (Simmondsia chinensis)* – DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 10 WEEKS AFTER SPRAY APPLICATION.**

***Parsnips, Shallots and Spring Onions* – DO NOT GRAZE OR CUT FOR STOCK FOOD.**

#### **EXPORT OF TREATED PRODUCE**

Growers should note that Maximum Residue Limits (MRLs) or import tolerances may not exist in all markets for edible produce treated with Imtrade Linuron 800 WG Herbicide. If you are growing edible produce for export, please check with Imtrade Australia Pty Ltd for the latest information on MRLs and import tolerances before using Imtrade Linuron 800 WG Herbicide.

## GENERAL INSTRUCTIONS

Imtrade Linuron 800 WG Herbicide is a foliar absorbed and residual herbicide i.e. it controls weeds present at application by absorption through the foliage and subsequent germination of weeds through root absorption. For best results, apply to well cultivated, weed-free surface which is free of trash and clods. **DO NOT** cultivate after application of Imtrade Linuron 800 WG Herbicide as weed control may be reduced and/or crop injury may result.

## EQUIPMENT & MIXING INSTRUCTIONS

Imtrade Linuron 800 WG Herbicide is a WG formulation to be mixed with water and applied as a spray. Partially fill the spray tank with water. Measure the amount of Imtrade Linuron 800 WG Herbicide required as indicated in the Directions for Use table. Add this to the spray tank with the agitation system engaged. Top up tank to the correct water volume. To prevent nozzle blockage, strainer and nozzle screens should be 50 mesh or coarser.

### THE MATERIAL MUST BE KEPT IN SUSPENSION AT ALL TIMES BY CONTINUOUS AGITATION.

#### Water Volume & Spray Pressure

- **Pre-emergence:** 300 – 500L/ha and 200 – 300 kPa pressure.
- **Post-emergence:** Use minimum of 300L/ha and 140 – 170 kPa pressure.
- **Band Treatments:** Use proportionately less.
- **Wheat, Barley, Oats only:**
  - **Aerial Application:** Use a minimum of 20L/ha spray under optimum conditions otherwise a minimum of 30L/ha.
  - **Ground Application:** Apply in a minimum of 90L/ha (spray).

## SPRAY EQUIPMENT HYGIENE

Before and after spraying Imtrade Linuron 800 WG Herbicide, spray equipment including tanks, nozzles, filters and spraylines should be cleaned thoroughly with a commercially available tank and equipment cleaner such as All Clear®.

## RESISTANT WEEDS WARNING

GROUP	<b>5</b>	HERBICIDE
-------	----------	-----------

Imtrade Linuron 800 WG Herbicide is a member of the Urea-group of herbicides. Imtrade Linuron 800 WG Herbicide has the inhibitors of photosynthesis at photosystem II mode of action. For weed resistance management Imtrade Linuron 800 WG Herbicide is a Group 5 Herbicide. Some naturally occurring weed biotypes resistant to Imtrade Linuron 800 WG Herbicide and other inhibitors of photosynthesis at photosystem II Herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Imtrade Linuron 800 WG Herbicide or other inhibitors of photosynthesis at photosystem II Herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Imtrade Australia Pty Ltd accepts no liability for any losses that may result from the failure of Imtrade Linuron 800 WG Herbicide to control resistant weeds.

Strategies to minimise the risk of herbicide resistance are available. Consult your farm chemical supplier, consultants, local Department of Agriculture or Primary Industries or local Imtrade Australia Pty Ltd representative.

### RE-ENTRY

**DO NOT** enter treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

### PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Imtrade Linuron 800 WG Herbicide is slightly toxic to fish. **DO NOT** contaminate streams, rivers or watercourses with this product or the used containers.

### STORAGE AND DISPOSAL

Keep out of reach of children. Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. **DO NOT** store for prolonged periods in direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers.

Triple rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers or product.

## SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. **DO NOT** inhale dust or spray mist. When opening the container, preparing spray, wear cotton overalls, over normal clothing, buttoned to the neck and wrist (or equivalent clothing) and chemical-resistant gloves and a half face respirator. If applying by low pressure hand wand and high pressure hand wand, wear cotton overalls, over normal clothing, buttoned to the neck and wrist (or equivalent clothing) and chemical-resistant gloves and a half face respirator. If product on skin, immediately wash area with soap and water. Wash hands after use. After each day's use wash gloves, respirator and if rubber wash with detergent and warm water and contaminated clothing.

## FIRST AID

First aid is not generally required. If in doubt, contact a Poisons Information Centre (*Phone Australia 13 11 26; New Zealand 0800 764 766*) or a doctor).

### Additional GHS Hazard & Precautionary Statements

•Suspected of causing cancer; •May damage fertility or the unborn child; •May cause damage to organs through prolonged or repeated exposure; •Very toxic to aquatic life with long-lasting effects; •**DO NOT** get on clothing; •**DO NOT** eat, drink or smoke when using this product; •Rinse mouth; •If eye irritation persists: Get medical advice; •**IF SWALLOWED:** Rinse mouth. Do **NOT** induce vomiting; •If exposed or concerned: Get medical advice; •In case of fire, use carbon dioxide, dry chemical, foam, water fog; •Store locked up.

## SAFETY DATA SHEET

For further information refer to the Safety Data Sheet, which can be obtained from your supplier or Imtrade Australia Pty Ltd website at [www.imtrade.com.au](http://www.imtrade.com.au)

## CONDITIONS OF SALE

Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Imtrade Australia Pty Ltd has any authority to add to or alter these conditions.

® Other Registered Trademarks

<b>UN 3077</b>	<b>ENVIRONMENTALLY HAZARDOUS SUBSTANCES, SOLID, N.O.S. (contains LINURON)</b>
<b>IN A TRANSPORT EMERGENCY, DIAL 000 POLICE OR FIRE BRIGADE</b>	
<b>PG III</b>	<b>HAZCHEM 2Z</b>

APVMA Approval No: 81574/131931