

#### STORAGE AND DISPOSAL

Keep out of reach of Children. Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. Triple-rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemical on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers or product. **DO NOT** re-use empty container for any other purpose.

#### Bags

Store in the closed, original container in a dry, cool, well-ventilated area, out of direct sunlight. Single-rinse or shake reminder into spray tank/water/dip/drench etc. **DO NOT** dispose of undiluted chemicals on-site. Puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers or product.

#### SAFETY DIRECTIONS

May irritate the eyes, nose, throat and skin. Avoid contact with eyes and skin. **DO NOT** inhale dust or spray mist. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

#### FIRST AID

First aid is not generally required. If in doubt, contact a Poisons Information Centre (*Phone Australia 13 11 26; New Zealand 0800 764 766*) or a doctor.

Refer to attached Leaflet for additional GHS Hazard & Precautionary Statements.

#### SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet which can be obtained from your supplier or Imtrade Australia Pty Ltd website at [www.imtrade.com.au](http://www.imtrade.com.au)

#### CONDITIONS OF SALE

Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Imtrade Australia Pty Ltd has any authority to add to or alter these conditions.

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

# IMTRADE BROMACIL 800 WG HERBICIDE

ACTIVE CONSTITUENT: 800g/kg BROMACIL

**GROUP 5 HERBICIDE**

For the control of certain annual broadleaved weeds and grasses in Citrus, Pineapple, Asparagus and in Commercial and Industrial areas, Rights of Way and around Agricultural Buildings as per the Directions for Use table

**IMPORTANT: Read the attached Leaflet before use.**

**GROUP  
5**

**FORM  
WG**

**CONTENTS  
2kg-15kg**

<b>UN 3077</b>	ENVIRONMENTALLY HAZARDOUS SUBSTANCE, SOLID, N.O.S. (contains BROMACIL)
In a Transport Emergency Dial 000 Police or Fire Brigade	
PG III	HAZCHEM Z2

APVMA Approval No: 87247/134347



Batch No.  
DOM:

**IMTRADE**  
CropScience  
17 Ocean St, Kwinana Beach WA 6167  
08 9419 0333 | [www.imtrade.com.au](http://www.imtrade.com.au)

**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**



# **IMTRADE BROMACIL 800 WG HERBICIDE**

**ACTIVE CONSTITUENT: 800g/kg BROMACIL**

**GROUP 5 HERBICIDE**

For the control of certain annual broadleaved weeds and grasses in Citrus, Pineapple, Asparagus and in Commercial and Industrial areas, Rights of Way and around Agricultural Buildings as per the Directions for Use table.

**IMPORTANT: READ THIS LEAFLET THOROUGHLY  
BEFORE OPENING OR USING THIS PRODUCT**

**APVMA APPROVAL No: 87247/134347**

Imtrade Australia Pty Ltd  
ABN 13 090 151 134  
17 Ocean Street, Kwinana WA 6167  
Tel: 1800 171 799 Fax: 1800 171 788  
Web: [www.imtrade.com.au](http://www.imtrade.com.au)

## RESTRAINTS

**DO NOT allow spray to contact foliage or fruit of desirable plants.**

**For best results sufficient moisture, either by rainfall or irrigation, is required within 3-4 days of treatment to carry BROMACIL 800 WG into the soil.**

## DIRECTIONS FOR USE

CROP	WEEDS CONTROLLED	STATE	RATE/HA**	CRITICAL COMMENTS
Asparagus (older than 1 year)	Annual Weeds – Annual Ryegrass, Barley Grass, Bathurst Burr, Brome Grass, Caltrop, Capeweed, Cobbler's-pegs, Doublegee (Spiny Emex, Three-cornered Jack), Fat Hen, Khaki Weed (seedling), Noogoora Burr, Paterson's Curse, Prairie Grass, Annual Ragweed, Saffron Thistle, Summer Grass, Thornapple, Wild Mustard,	All States	1.1 to 2.5kg	Apply after cultivation to established plantings only as damage may occur to young plantings. Apply 3-4 weeks prior to spear emergence in Spring or before the early cutting period. If cultivation occurs during cutting period apply immediately after last cultivation. If needed, apply a second spray after harvest to keep fields weed-free. Irrigate before or as soon as possible after spraying. (See Crop Safety Section).  <b>Note:</b> For Wild Oat control, use no less than 2.5kg/ha.
			1kg/500L water	<b>Spot Spray</b> – wet plant thoroughly to point of run-off (See Crop Safety Section).
Citrus – established groves only	Wild Oats, Wild Radish, Wild Turnip, Winter Grass.		2.2 to 4.5kg	* Apply to weed-free ground in Autumn and Spring. After 1-2 years of application, once per year may be adequate. <b>DO NOT</b> apply more than 9kg per hectare per year. (See Crop Safety Section).  <b>Note:</b> For Wild Oat control, use no less than 3kg
				<b>Spot Treatment</b> – spray to point of run-off. (See Crop Safety Section).

CROP	WEEDS CONTROLLED	STATE	RATE/HA**	CRITICAL COMMENTS
Pineapples	Barnyard Grass, Bilygoat Weed/Bluetop, Cobbler's-pegs, Crowsfoot, Guinea Grass, Red Natal Grass, Sourgrass, Stinking Roger, Summer Grass, Thickhead	Qld, NT only	2.2 to 4.5kg	Apply as a blanket spray prior to weed emergence immediately after planting or before planting material begins to grow.
			2.2kg	* If needed apply as a directed inter-row spray prior to flower differentiation. <b>DO NOT</b> spray ovetop of plants. <b>DO NOT</b> apply more than 9kg per hectare per year.
Industrial Weed Control Commercial & Industrial areas; Rights of Way; around Agricultural Buildings	<b>Annual Weeds –</b> Annual Ryegrass, Barley Grass, Bathurst Burr, Brome Grass, Caltrop, Capeweed, Cobbler's-pegs, Doublegee (Spiny Emex, Three-cornered Jack), Fat Hen, Khaki Weed (seedling), Noogoora Burr, Paterson's Curse, Prairie Grass, Annual Ragweed, Saffron Thistle, Summer Grass, Thornapple, Wild Mustard, Wild Oats, Wild Radish, Wild Turnip, Winter Grass.	All States	3.5 to 6.5kg	* The lowest rate will provide control in low rainfall areas (250mm or less).
			<b>Retreatment</b> – All the above weeds	2 to 6.5kg
	<b>Small Areas</b> – Annual and Perennial weeds		20g/10m <sup>2</sup> (20kg/ha)	*

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

\* If weeds are present add non-ionic surfactant at 100 mL/100L (0.1% v/v) of spray solution.

\*\* Use lower rates on lighter soils low in clay or organic matter and higher rates on heavier soils high in clay or organic matter.

## **WITHHOLDING PERIODS: NOT REQUIRED WHEN USED AS DIRECTED.**

### **GENERAL INSTRUCTIONS**

For best results apply shortly before weed growth occurs to moist well prepared trash and clod free soil. Well-established weeds should be removed before applying BROMACIL 800 WG. Actively growing weeds up to about 10cm in height may be controlled, but a wetting agent must be used.

**DO NOT** cultivate soil surface after application as efficacy may be reduced and/or crop injury may result.

For best results sufficient moisture, either by rainfall or irrigation is required within 3-4 days of treatment to carry BROMACIL 800 WG into the soil. With irrigation thoroughly wet the entire area. Where rainfall is limited during the growth period, control of deep-rooted hard-to-kill perennial weeds will not be satisfactory.

### **EQUIPMENT AND APPLICATION**

Fill spray tank no more than 50% full with clean water. Before adding BROMACIL 800 WG begin agitating vigorously and continue agitation during the entire mixing and spraying operation. Pour required amount of BROMACIL 800 WG steadily into the spray tank. Allow vigorous bypass agitation to completely disperse product.

**DO NOT** dump BROMACIL 800 WG into spray tank all at once.

After adding required quantity of BROMACIL 800 WG and obtaining complete dispersion, continue to fill tank to desired level for spraying. Thorough agitation of the spray liquid should continue during the entire spraying operation. Spray solution should not be left standing in the tank overnight.

#### ***Large Areas***

Use a fixed boom sprayer or handgun. To prevent nozzle blockage, strainer and nozzle screens should be 50 mesh or coarser. Always agitate in spray tank to keep material in suspension.

##### **a) Fixed Boom Sprayer**

Use 1000-2000L/ha water for industrial weed control. For selective weed control use 250-450L/ha at 200-300 kPa pressure.

##### **b) Handgun**

Use 1000-2000L/ha and 600-800 kPa pressure.

#### ***Small Areas***

Use knapsack or pressurized hand sprayer or watering can. Shake or stir frequently.

### **SPRAYER CLEANUP**

Thoroughly clean all traces of BROMACIL 800 WG from application equipment immediately after use. Flush tank, pump, hoses and boom with several changes of water after removing spray nozzles and screen (clean these parts separately).

## RESISTANT WEEDS WARNING

### GROUP 5 HERBICIDE

BROMACIL 800 WG is a member of the uracils group of herbicides. BROMACIL 800 WG has the inhibitor of photosynthesis at photosystem II mode of action. For weed resistance management, BROMACIL 800 WG is a Group 5 Herbicide. Some naturally occurring weed biotypes resistant to BROMACIL 800 WG and other inhibitors of photosynthesis at photosystem II herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by BROMACIL 800 WG or other inhibitors of photosynthesis at photosystem II herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Imtrade CropScience accepts no liability for any loss that may result from the failure of BROMACIL 800 WG to control resistant weeds.

Advice as to strategies and alternative treatments that can be used should be obtained from your local supplier, consultant, local Department of Agriculture, Primary Industries Department or an Imtrade CropScience representative.

### RE-ENTRY PERIODS

After treating Asparagus or Pineapple, **DO NOT** allow entry into treated areas for 14 days unless wearing cotton overalls buttoned to the neck and wrist over normal clothing and chemical resistant gloves. Clothing must be laundered after each days use.

After treating Citrus, **DO NOT** allow entry into treated areas for 3 days unless wearing cotton overalls buttoned to the neck and wrist over normal clothing and chemical resistant gloves. Clothing must be laundered after each days use.

### PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

**DO NOT** apply or drain or flush equipment on or near desirable trees or other plants or where their roots may extend.

Movement of soil from treated areas should be avoided because damage to off-target species may occur.

**DO NOT** spray under weather conditions or from spraying equipment which may cause spray to drift onto nearby susceptible plants or crops, cropping lands or pastures.

**Asparagus:** **DO NOT** apply to newly seeded asparagus, or to young plants during the first growing season after setting or to plants whose roots are exposed.

**Citrus:** Use with caution on light sandy soils as yellowing may occur.

**Treated areas for Asparagus and Citrus - DO NOT** re-plant any crop within three (3) years of last application as injury may occur.

**Pineapples:** Only pineapples can be re-planted into the same area within the 3 year plan.

## **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT**

**DO NOT** contaminate streams, rivers or waterways with the chemical or used containers.

### **STORAGE AND DISPOSAL**

Keep out of reach of Children. Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight.

Triple-rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemical on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers or product. **DO NOT** re-use empty container for any other purpose.

### **Bags**

Store in the closed, original container in a dry, cool, well-ventilated area, out of direct sunlight. Single-rinse or shake remainder into spray tank/water/dip/drench etc. **DO NOT** dispose of undiluted chemicals on-site. Puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers or product.

### **SAFETY DIRECTIONS**

May irritate the eyes, nose, throat and skin. Avoid contact with eyes and skin. **DO NOT** inhale dust or spray mist. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

### **FIRST AID**

First aid is not generally required. If in doubt, contact a Poisons Information Centre (*Phone Australia 13 11 26; New Zealand 0800 764 766*) or a doctor.

### **Additional GHS Hazard & Precautionary Statements**

•Harmful if swallowed; •Very toxic to aquatic life with long-lasting effects; •**DO NOT** get on clothing; •Wash contacted areas thoroughly after handling; •Avoid release to the environment; •Wear protective gloves, protective clothing and eye or face protection; •Get medical advice or attention if you feel unwell; •Wash contaminated clothing before reuse; •**IF SWALLOWED:** Call a POISON CENTRE or doctor if you feel unwell; •**IF SWALLOWED:** Rinse mouth. Do **NOT** induce vomiting; •In case of fire: Use carbon dioxide, dry chemical, foam, to extinguish; •Store in a dry place.

### **SAFETY DATA SHEET**

Additional information is listed in the Safety Data Sheet which can be obtained from your supplier or Imtrade CropScience website at [www.imtrade.com.au](http://www.imtrade.com.au)

### **CONDITIONS OF SALE**

Imtrade CropScience shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Imtrade CropScience has any authority to add to or alter these conditions.

<b>UN 3077</b>	<b>ENVIRONMENTALLY HAZARDOUS SUBSTANCES, SOLID, N.O.S. (contains BROMACIL)</b>
<b>IN A TRANSPORT EMERGENCY, DIAL 000 POLICE OR FIRE BRIGADE</b>	
<b>PG III</b>	<b>HAZCHEM 2Z</b>



**APVMA Approval No: 87247/134347**