

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight.

All other non-refillable containers

Triple-rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemical on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers or product. **DO NOT** re-use empty container for any other purpose.

Bags

Store in the closed, original container in a dry, cool, well-ventilated area, out of direct sunlight. Single-rinse or shake remainder into spray tank/water/dip/drench, etc. **DO NOT** dispose of undiluted chemicals on-site. Puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers or product.

SAFETY DIRECTIONS

Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. If the product in eyes, wash it out immediately with water. If product on skin, immediately wash area with soap and water. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and a washable hat, elbow-length chemical resistant gloves, and face shield or goggles. Wash hands after use. After each day's use wash gloves, face shield or goggles and contaminated clothing.

ADDITIONAL USER SAFETY INFORMATION

WARNING: May cause birth defects.

FIRST AID

First aid is not generally required. If in doubt, contact a Poisons Information Centre (Phone Australia 13 11 26; New Zealand 0800 764 766) or a doctor.

Refer to attached Leaflet for additional GHS Hazard & Precautionary Statements.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet which can be obtained from your supplier or Imtrade Australia Pty Ltd website at www.imtrade.com.au

CONDITIONS OF SALE

Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Imtrade Australia Pty Ltd has any authority to add to or alter these conditions.

DALBIE® is a Registered Trademark of Imtrade Australia Pty Ltd

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

IMTRADE

DALBIE®

800 WG FUNGICIDE

ACTIVE CONSTITUENT: 800g/kg PROTHIOCONAZOLE

GROUP 3 FUNGICIDE

For the control of various diseases in Peanuts, Cereals and Canola
as specified in the Directions for Use table.

Important: Read the attached Leaflet before use.

FORM
WG

CONTENTS
1KG-25KG

UN 3077

ENVIRONMENTALLY
HAZARDOUS
SUBSTANCE, SOLID,
N.O.S. (contains
PROTHIOCONAZOLE)

In a Transport Emergency
Dial 000 Police or Fire Brigade

PG III

HAZCHEM 2Z

APVMA APPROVAL NO: 87492/128572



Batch No.
DOM:

IMTRADE
CropScience

17 Ocean St, Kwinana Beach WA 6167
08 9419 0333 | www.imtrade.com.au

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

IMTRADE

 **DALBIE**®

800 WG FUNGICIDE

ACTIVE CONSTITUENT: 800g/kg PROTHIOCONAZOLE

GROUP 3 FUNGICIDE

For the control of various diseases in Peanuts, Cereals and Canola
as specified in the Directions for Use table

**IMPORTANT: READ THIS LEAFLET THOROUGHLY
BEFORE OPENING OR USING THIS PRODUCT**

APVMA APPROVAL No: 87492/128572

Imtrade Australia Pty Ltd
ABN 13 090 151 134
17 Ocean Street, Kwinana WA 6167
Tel: 1800 171 799 Fax: 1800 171 788
Web: www.imtrade.com.au

DIRECTIONS FOR USE

RESTRAINTS:

CEREALS AND CANOLA

A maximum of two applications may be made per cereal or canola crop.

PEANUTS

DO NOT apply where the slope exceeds 7%.

DO NOT apply by aircraft to Peanut crops.

DO NOT apply if heavy rains or storms that are likely to cause runoff are forecast within 48 hours.

DO NOT apply to waterlogged soils

DO NOT irrigate to the point of runoff for at least 48 hours after application.

SPRAY DRIFT RESTRAINTS

Specific definitions for terms used in this selection of the label can be found at apvma.gov.au/spraydrift

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. The buffer zones in the relevant buffer zone table/s below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

DO NOT apply unless the wind speed is between three and 20 kilometres per hour at the application site during the time of application.

DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

DO NOT apply by a boom sprayer unless the following requirements are met:

- spray droplets not smaller than a (MEDIUM) spray droplet size category
- minimum distances between the application site and downwind sensitive areas are observed
(see Mandatory Zone Table).

MANDATORY BUFFER ZONE

DO NOT apply if there are aquatic and wetland areas, including aquacultural ponds, surface streams and rivers downwind from the application area and within the **mandatory buffer zones** shown in Tables A, B and C below.

Cereals

Table A – Buffer Zones for Protection of the Aquatic Environment	
FOR AERIAL APPLICATION	
Wind Speed Range at Time of Application	Mandatory Downwind Buffer Zone
Fixed-Wing	
From 3 to 20 kilometres per hour	80 metres
Helicopter	
From 3 to 20 kilometres per hour	60 metres
FOR GROUND APPLICATION	
From 3 to 20 kilometres per hour	10 metres

Canola

Table B – Buffer Zones for Protection of the Aquatic Environment	
FOR AERIAL APPLICATION	
Wind Speed Range at Time of Application	Mandatory Downwind Buffer Zone
Fixed-Wing	
From 3 to 20 kilometres per hour	180 metres
Helicopter	
From 3 to 20 kilometres per hour	120 metres
FOR GROUND APPLICATION	
From 3 to 20 kilometres per hour	5 metres

Peanuts

Table C – Buffer Zones for Protection of the Aquatic Environment	
FOR GROUND APPLICATION	
Wind Speed Range at Time of Application	Mandatory Downwind Buffer Zone
From 3 to 20 kilometres per hour	5 metres

CROP	DISEASE	STATE	RATE	CRITICAL COMMENTS
Barley	Net Form Net Blotch (<i>Pyrenophora teres f. teres</i>)	All States	40-80g/ha + 40-80g/ha Imtrade Turbulence 800 WG Fungicide	Monitor crops from mid tillering. On susceptible varieties apply at the first sign of disease development. Monitor and reapply within 14 to 21 days if conditions favour disease development. Use the higher rates where conditions favour severe disease. Where lower rates are used apply with a suitable adjuvant (refer to Use of Adjuvant).
	Spot form Net Blotch (<i>Pyrenophora teres f. maculata</i>)			Monitor crops from mid tillering. Use the higher rate in higher yielding crops where conditions favour disease development or susceptible varieties are grown.
	Powdery Mildew (<i>Blumeria graminis f.sp. hordei</i>)			Monitor crops from mid tillering (earlier if no effective seed treatment has been applied). On susceptible varieties apply at the first sign of disease development. Monitor and reapply within 14 to 21 days if conditions favour disease development. Use the higher rates where conditions favour severe disease. Where lower rates are used apply with a suitable adjuvant (refer to Use of Adjuvant).
	Leaf Scald (<i>Rhynchosporium secalis</i>)			Monitor crops from late tillering. Apply at the first sign of disease development. Monitor and reapply within 14 to 21 days if conditions favour disease development. Use the higher rates where conditions favour severe disease, or disease is established in the lower canopy. Where lower rates are used apply with a suitable adjuvant (refer to Use of Adjuvant).
	Leaf Rust (<i>Puccinia hordei</i>)			

CROP	DISEASE	STATE	RATE	CRITICAL COMMENTS
Oats	Stem Rust (<i>Puccinia graminis</i> f.sp. <i>avenae</i>)	All States	80 g/ha + 80 g/ha Imtrade Turbulence 800 WG Fungicide	Monitor crops from early stem elongation, and on susceptible varieties apply at the first sign of infection. Refer to General Instructions – Disease control in Oats , for potential risks associated with application to oats
	Leaf Rust (<i>Puccinia coronata</i> f.sp. <i>avenae</i>)		+ Adjuvant (refer to Use of Adjuvant)	Monitor crops from early stem elongation, and on susceptible varieties apply at the first sign of infection. Refer to General Instructions – Disease control in Oats , for potential risks associated with application to oats.
	Septoria Blotch (<i>Phaeosphaeria avenaria</i>)		40-80g/ha + 40-80g/ha Imtrade Turbulence 800 WG Fungicide	Monitor crops from early tillering and on susceptible varieties apply at the first sign of infection. Use the higher rate in higher yielding crops where conditions favour disease development or susceptible varieties are grown. Continue to monitor crops after application. Re-application may be required if conditions favour disease development. Where lower rates are used, apply with a suitable adjuvant (refer to Use of Adjuvant). Refer to General Instructions – Disease control in Oats , for potential risks associated with application to oats.

CROP	DISEASE	STATE	RATE	CRITICAL COMMENTS
Wheat	Stripe Rust (<i>Puccinia striiformis</i>)	All States	40-80g/ha +	Monitor crops from early stem elongation, and on susceptible varieties apply at the first sign of infection. Use the higher rate in higher yielding crops where conditions favour disease development or susceptible varieties are grown. Continue to monitor crops after application, re-application may be required if conditions favour disease development and initial application is made before the flag leaf has emerged.
	Stem Rust (<i>Puccinia graminis tritici</i>)		40-80g/ha Imtrade Turbulence 800 WG Fungicide	
	Leaf Rust (<i>Puccinia recondita</i> f.sp. <i>tritici</i> , <i>Puccinia triticina</i>)		+	
	Fusarium Head Blight/Head Scab (<i>Fusarium graminearum</i>)		Adjuvant (refer to Use of Adjuvant)	
	Yellow Leaf Spot (<i>Pyrenophora tritici-repentis</i>)		40-80g/ha +	
	Septoria Nodorum -Glume Blotch (<i>Phaeosphaeria nodorum</i>)		40-80g/ha Imtrade Turbulence 800 WG Fungicide	
Powdery Mildew (<i>Blumeria graminis</i> f.sp. <i>tritici</i>)		Monitor crops from late tillering and spray before disease has infected any of the top three leaves of the crop. Aim to protect the three top leaves of the plant from disease.		
		Monitor crops from late tillering. Aim to protect the three top leaves of the plant from disease. Where lower rates are used apply with a suitable adjuvant (refer to Use of Adjuvant).		
		Monitor crops from mid tillering. Apply at the first sign of disease development. Monitor and reapply within 14 to 21 days if conditions favour disease development. Use the higher rates where conditions favour severe disease, or disease is established in the lower canopy. Where lower rates are used apply with a suitable adjuvant (refer to Use of Adjuvant).		

CROP	DISEASE	STATE	RATE	CRITICAL COMMENTS
Triticale	Stripe Rust (<i>Puccinia striiformis</i>)	All States	40-80g/ha + 40-80g/ha Imtrade Turbulence 800 WG Fungicide + Adjuvant (refer to Use of Adjuvant)	Monitor crops from early stem elongation, and on susceptible varieties apply at the first sign of infection. Use the higher rate in higher yielding crops where conditions favour disease development or susceptible varieties are grown. Continue to monitor crops after application. Re-application may be required if conditions favour disease development and initial application is made before the flag leaf has emerged.
Canola	Blackleg (<i>Leptosphaeria maculans</i>)	All States	100 - 120g/ha + 100-120g/ha Imtrade Turbulence 800 WG Fungicide	Apply at the 4 to 6 leaf crop stage of blackleg susceptible varieties (blackleg ratings of MS or lower) or in situations of high blackleg risk (refer to General Instructions – Disease control in Canola). Will reduce lodging and stem canker from blackleg. A follow up application may be required at green bud stage in high disease risk situations or where an effective blackleg seed treatment has not been used.
	Sclerotinia stem Rot (<i>Sclerotinia sclerotiorum</i>)			Apply between 20 and 50% (full bloom) flowering. For best results apply as a preventative application at 20-30% flowering prior to significant disease expression (refer to General Instructions – Disease control in Canola). Good coverage throughout the entire canopy is essential. Using a water rate at the higher end of the range (i.e. 100 L/ha for ground application and 30 L/ha for aerial application) will improve spray coverage. Apply the higher rate under high disease pressure.

CROP	DISEASE	STATE	RATE	CRITICAL COMMENTS
Peanuts	Early Leaf Spot, Late Leaf Spot, Rust	Qld, NSW, WA, NT	150 - 240 g/ha	Begin spraying no later than 3-4 weeks after emergence. Repeat applications at 10-14 day intervals. Under conditions conducive to high disease pressure, prolonged wet weather or heavy rains, use the high rate and shortest spray interval. In localities with frequent and persistent rainfall, or in irrigated crops, it is recommended that DALBIE® 800 WG usage should be integrated with a base program of protectant fungicide applications for best control of leaf diseases. Apply no more than four applications of DALBIE® 800 WG per season.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

A MANDATORY NO-SPRAY ZONE IS REQUIRED FOR PROTECTION OF THE ENVIRONMENT. REFER TO RESTRAINTS.

WITHHOLDING PERIODS

Harvest

Canola: NOT REQUIRED WHEN USED AS DIRECTED.

Peanuts: DO NOT HARVEST FOR 4 WEEKS AFTER APPLICATION.

Cereals: DO NOT HARVEST FOR 5 WEEKS AFTER APPLICATION.

Grazing

Canola and Cereals: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION

Peanuts: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER APPLICATION.

EXPORT OF TREATED PRODUCE

Growers should note that MRLs or import tolerances do not exist in all markets for produce treated with DALBIE® 800 WG. If you are growing produce for export, please check with Intrade Australia for the latest information on MRLs and import tolerances.

GENERAL INSTRUCTIONS

FOLIAR DISEASES ON CEREAL CROPS

Monitor the crop regularly for symptoms of disease. Generally spray at the first sign of disease, although this will depend on factors such as expected weather conditions and the particular crop variety resistance. Refer to Directions for Use for particular disease recommendations. Up to two sprays of DALBIE® 800 WG may be applied per season to the crop. Ensure good coverage of all susceptible plant parts.

DISEASE CONTROL IN OATS

Caution: Application of tebuconazole (present in Imtrade Turbulence 800 WG Fungicide) to some varieties of Oats may result in early senescing and bronzing of leaves.

Varieties most at risk may also exhibit this trait under various stress conditions not related to fungicide sprays. Mitika variety of Oats has been identified as being susceptible to this condition when tebuconazole is applied, although other varieties may also be susceptible.

The potential disease control to be achieved by using DALBIE® 800 WG + Imtrade Turbulence 800 WG Fungicide in Mitika Oats should be weighed against the risk of crop damage.

DISEASE CONTROL IN CANOLA

Blackleg

Higher Blackleg risk can be expected in higher rainfall districts (above 500mm annual rainfall), where crops are grown within 500m of a previous year's stubble and in later sown crops (May to August). Other factors will also increase the risk of Blackleg infection, including the intensity of canola cropping in a district, rainfall before sowing and the frequency of growing the same canola cultivar. Consult industry guidelines for more detailed assessment of blackleg risk in specific situations. Up to two sprays of DALBIE® 800 WG may be applied per season to the crop.

Sclerotinia

DALBIE® 800 WG in combination with Imtrade Turbulence 800 WG Fungicide is most effective when application is made prior to conditions conducive to sclerotinia infection.

Infection and disease development are most conducive in warmer winter or spring conditions with extended periods of leaf wetness due to rainfall, dew and high humidity. Sclerotinia is most likely to develop where day temperatures are warmer coinciding with a saturated soil profile and rainfall events. Refer also to industry guidelines for advice on conditions under which sclerotinia are most likely to develop.

Control of Sclerotinia Stem Rot is more effective in crops which have a uniform flowering. Uneven flowering (e.g. caused by staggered germinations) makes optimum spray timing difficult and two sprays may be required in these crops.

Generally a single application of DALBIE® 800 WG + Imtrade Turbulence 800 WG Fungicide at 20 to 30% flowering will control sclerotinia in crops with a short flowering interval. Crops with an extended flowering period may require a second application prior to 50% flowering (full-bloom) to adequately control sclerotinia if conditions late in the season are conducive to development of disease.

Length of protection may be reduced in bulky crops where coverage is difficult and where there is growth dilution of the fungicide. For optimum protection, application should be directed to obtain coverage on petals, leaves and stems.

FOLIAR DISEASES ON PEANUTS

Monitor the crop regularly for symptoms of disease. Generally spray at the first sign of disease, although this will depend on factors such as expected weather conditions and the particular crop variety resistance. Refer to Directions for Use for particular disease recommendations. Up to four sprays of DALBIE® 800 WG per season may be applied to the crop. Ensure good coverage of all susceptible plant parts.

MIXING

Add the required quantity of DALBIE® 800 WG to water in the spray vat while under agitation. Allow to fully disperse before adding products with other formulation types.

Ground:

Wheat, Barley, Oats and Triticale: Apply product using a spray volume of 70 – 100L/ha and a MEDIUM spray quality as defined by the ASABE S572 Standard.

Canola: Apply product using a spray volume of 60 – 100L/ha and a MEDIUM spray quality as defined by the ASABE S572 Standard.

Peanuts: Apply using a spray volume of 150 to 450L/ha and a nozzle that provides spray droplets not smaller than a (MEDIUM) spray droplet size category. Ensure thorough coverage of foliage.

Aerial: (not Pyrethrum or Peanut)

Apply product using a minimum spray volume of 20L/ha and a MEDIUM spray quality as defined by the ASABE S572 Standard.

USE OF ADJUVANT

Depending on the disease that is to be treated in the crop, some benefit in efficacy may be gained from addition of an appropriate adjuvant to the spray mixture. Follow these guides when deciding on the addition of an adjuvant to the tank mixture prior to spraying.

Disease	Addition of Adjuvant	
	DALBIE® 800 WG + Imtrade Turbulence 800 WG Fungicide	
	40 + 40 g/ha	80 + 80 g/ha
Barley		
Net Form Net Blotch	Yes	Not required
Spot Form Net Blotch	Yes	Not required
Powdery Mildew	Not required	Not required
Leaf Scald	Yes	Not required
Leaf Rust	Yes	Not required
Oats		
Stem Rust	N/A	Yes (BS 1000 only)
Leaf Rust	N/A	Yes (BS 1000 only)
Septoria Blotch	Yes	Not required
Wheat		
Stripe Rust	Yes	Yes (BS 1000 only)
Stem Rust	Yes	Yes (BS 1000 only)
Leaf Rust	Yes	Yes (BS 1000 only)
Yellow Leaf Spot	Not required	Not required
Septoria Nodorum – Glume Blotch	Yes	Not required
Powdery Mildew	Yes	Not required
Fusarium Head Blight/Head Scab	Yes	Yes (BS 1000 only)
Triticale		
Stripe Rust	Yes	Yes (BS 1000 only)
Canola		
	100 + 100 g/ha	120 + 120 g/ha
All Diseases	Not required	Not required
Peanut		
	260 + 260 g/ha	
All Diseases	Not required	

Suitable Adjuvants	Rate (%v/v)	Comments
BS 1000	0.25%	Can be used at all rates of DALBIE® 800 WG + Imtrade Turbulence 800 WG Fungicide for ground and aerial application.
Hasten®	1%	For use with DALBIE® 800 WG + Imt Turbulence 800 WG Fungicide at 40 g/ha only. DO NOT use at rates above 40 g/ha. DO NOT use for aerial application.
Rocket®	1%	
Kwickin®	1%	
D-C-Trate® Advance	1%	
D-C-Trate	1%	
Uptake®	0.5%	

COMPATABILITY

DALBIE® 800 WG is compatible with Imtrade Whack® 720 Fungicide. For the latest information on product compatibilities, contact your local Imtrade Australia Pty Ltd representative.

FUNGICIDE RESISTANCE WARNING

GROUP **3** FUNGICIDE

DALBIE® 800 WG is a member of the DMI group of fungicides. For fungicide resistance management, DALBIE® 800 WG is a Group **3** Fungicide. Some naturally occurring individual fungi resistant to DALBIE® 800 WG and other Group **3** Fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by DALBIE® 800 WG and other Group **3** Fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, Imtrade Australia Pty Ltd accepts no liability for any loss that may result from the failure of DALBIE® 800 WG to control resistant pathogens.

Advice as to strategies and alternative treatments that can be used should be obtained from your local supplier, consultant, local Department of Agriculture, Primary Industries Department or an Imtrade Australia Pty Ltd representative.

PRECAUTIONS

Re-entry Period

DO NOT allow entry into treated areas until spray has dried. If prior entry is necessary wear cotton overalls buttoned to the neck and wrists (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT

Very toxic to aquatic life.

DO NOT contaminate wetlands or watercourses with this product or used containers.

A spray drift minimisation strategy should be employed at all times. Spray drift may occur under adverse meteorological conditions or from certain spraying equipment. **DO NOT** allow spray to drift onto sensitive areas including, but not limited to, susceptible plants/crops, cropping land, pasture, natural streams, rivers, wetlands, waterways or human dwellings.

Integrated Pest Management - where IPM is practiced: DALBIE® 800 WG + Imtrade Turbulence 800 WG Fungicide may have adverse effects on some non-target beneficial invertebrates such as predatory mites.

STORAGE AND DISPOSAL

The method of disposal of the container depends on the container type. Read the Storage and Disposal instructions on the label that is attached to the container.

Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight.

All other non-refillable containers

Triple-rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemical on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers or product. **DO NOT** re-use empty container for any other purpose.

Bags

Store in the closed, original container in a dry, cool, well-ventilated area, out of direct sunlight. Single-rinse or shake remainder into spray tank/water/dip/drench, etc. **DO NOT** dispose of undiluted chemicals on-site. Puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers or product.

SAFETY DIRECTIONS

Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. If the product in eyes, wash it out immediately with water. If product on skin, immediately wash area with soap and water. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and a washable hat, elbow-length chemical resistant gloves, and face shield or goggles. Wash hands after use. After each day's use wash gloves, face shield or goggles and contaminated clothing.

ADDITIONAL USER SAFETY INFORMATION

WARNING: May cause birth defects.

FIRST AID

First aid is not generally required. If in doubt, contact a Poisons Information Centre (Phone Australia 13 11 26; New Zealand 0800 764 766) or a doctor.

Additional GHS Hazard & Precautionary Statements

•Harmful if swallowed; •May cause damage to organs through prolonged or repeated exposure; •Very toxic to aquatic life with long-lasting effects; •**DO NOT** breathe dusts; •**DO NOT** get on clothing; •Wash contacted areas thoroughly after handling; •**DO NOT** eat, drink or smoke when using this product; •Use personal protective equipment as required; •Get medical advice or attention if you feel unwell; •Brush off loose particles from skin; •**IF SWALLOWED:** Call a POISON CENTRE or doctor if you feel unwell; •**IF SWALLOWED:** Rinse mouth. Do **NOT** induce vomiting; •Store in a dry place.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet which can be obtained from your supplier or Imtrade Australia Pty Ltd website at www.imtrade.com.au

CONDITIONS OF SALE

Imtrade Australia Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Imtrade Australia Pty Ltd has any authority to add to or alter these conditions.

DALBIE® is a Registered Trademark of Imtrade Australia Pty Ltd

© Other Company Registered Trademarks

UN 3077	ENVIRONMENTALLY HAZARDOUS SUBSTANCES, SOLID, N.O.S. (contains PROTHIOCONAZOLE)
IN A TRANSPORT EMERGENCY, DIAL 000 POLICE OR FIRE BRIGADE	
PG III	HAZCHEM 2Z



APVMA Approval No: 87492/128572