

STORAGE AND DISPOSAL

Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight.

All other non-refillable containers

Triple rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemical on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers and product. **DO NOT** re-use empty container for any other purpose.

Envirodrum - Micro Matic Valve (110L)

Store the original sealed Envirodrum in a cool, well ventilated area. **DO NOT** store for prolonged periods in direct sunlight. **DO NOT** tamper with the Micro Matic valve or the security seal. **DO NOT** contaminate the Envirodrum with water or any other foreign matter. After each use of the product please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase.

Refillable Containers (1000L only)

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

FOR USE ONLY AS AN AGRICULTURAL AND HORTICULTURAL HERBICIDE, THIS PRODUCT IS TOO HAZARDOUS TO BE USED IN THE HOME GARDEN.

SAFETY DIRECTIONS

GENERAL

Very dangerous. Poisonous if inhaled or swallowed. Harmful if absorbed by skin contact. Will irritate the skin. Will damage the eyes. Avoid contact with eyes and skin. **DO NOT** inhale spray mist.

WHEN MIXING AND USING

When opening the container, preparing spray and using the prepared spray wear cotton overalls, over normal clothing, buttoned to the neck and wrist, elbow-length chemical resistant gloves, goggles and full facepiece respirator with dust cartridge or canister. If applying by hand, when opening the container, preparing spray and using the prepared spray wear cotton overalls, over normal clothing, buttoned to the neck and wrist and a washable hat, elbow-length chemical resistant gloves, goggles and disposable mist face mask covering mouth and nose.

AFTER USE

Wash hands after use. After each day's use wash gloves, goggles, respirator and if rubber wash with detergent and warm water and contaminated clothing.

SPRAY APPLICATION

DO NOT work in spray mist. **DO NOT** continue to use if skin irritation or nosebleed occurs. This may be caused by exposure to spray mist as a result of incorrect use of equipment or adverse climatic conditions. Stop and review handling and spraying techniques before further spraying. If symptoms persist seek medical advice.

Where there is a risk of exposure to spray mist, wear waterproof footwear and waterproof protective clothing, impervious gauntlet length gloves (rubber or PVC), goggles and a face mask and respirator covering nose and mouth and capable of filtering spray droplets. A high efficiency type particulate respirator is recommended by in any event use a respirator which complies with the requirements of AS1716 (Standards Association of Australia). Further advice on safety equipment should be obtained from a safety equipment manufacturer. Avoid contacting vegetation wet with spray but if necessary to do so wear waterproof footwear and waterproof protective clothing and gloves.

FIRST AID

If poisoning occurs get to a doctor or hospital quickly. If in eyes, hold eyes open and flood with water for at least 15 minutes and see a doctor.

NOTE TO PHYSICIANS

For additional advice on the treatment of paraquat poisoning please consult the booklet "The Treatment of Paraquat Poisoning: A Guide for Doctors".

Refer to attached Leaflet for additional GHS Hazard & Precautionary Statements.

DANGEROUS POISON

KEEP OUT OF REACH OF CHILDREN

CAN KILL IF SWALLOWED

DO NOT PUT IN DRINK BOTTLES

KEEP LOCKED UP

READ SAFETY DIRECTIONS BEFORE OPENING OR USING



**NOW REGISTERED
FOR OPTICAL
SPOT SPRAYING
TECHNOLOGY**

IMTRADE



GUERRILLA®

HERBICIDE

ACTIVE CONSTITUENTS: 300g/L PARAQUAT present as paraquat dichloride
12g/L AMITROLE

GROUP 22 | 34 HERBICIDE

For the control of a wide range of grasses and broadleaved
weeds as per Directions for Use table.

Important: Read the attached Leaflet before use.

GROUP

22

GROUP

34

FORM

SL

CONTENTS

5L-1000L

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from your supplier or Imtrade CropScience website at www.imtrade.com.au

CONDITIONS OF SALE

Imtrade CropScience warrants that this product conforms to the chemical description on the label. The use of GUERRILLA® being beyond the control of the manufacturer, Imtrade CropScience shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Imtrade CropScience has any authority to add to or alter these conditions.

GUERRILLA® is a Registered Trademark of Imtrade Australia Pty Ltd

UN 3016	
BIPYRIDILIUM PESTICIDES LIQUID, TOXIC, N.O.S. (contains Paraquat)	
In a Transport Emergency Dial 000 Police or Fire Brigade	
PG III	HAZCHEM 2X

APVMA Approval No: 89484/133061



Batch No.
DOM:



17 Ocean St, Kwinana Beach WA 6167
08 9419 0333 | www.imtrade.com.au

DANGEROUS POISON

KEEP OUT OF REACH OF CHILDREN

CAN KILL IF SWALLOWED

DO NOT PUT IN DRINK BOTTLES

KEEP LOCKED UP

READ SAFETY DIRECTIONS BEFORE OPENING OR USING



IMTRADE GUERRILLA® HERBICIDE

**ACTIVE CONSTITUENTS: 300g/L PARAQUAT present as paraquat dichloride
12g/L AMITROLE**

GROUP 22|34 HERBICIDE

For the control of a wide range of grasses and broadleaf weeds
as per Directions for Use table.

**IMPORTANT: READ THIS LEAFLET THOROUGHLY
BEFORE OPENING OR USING THIS PRODUCT**

APVMA Approval No: 89484/133061

Imtrade Australia Pty Ltd
ABN 13 090 151 134
17 Ocean Street, Kwinana, Western Australia 6167
Tel: (08) 9419 0333 Fax: (08) 9419 7516
Web: www.imtrade.com.au

DIRECTIONS FOR USE

RESTRAINTS:

For **PROFESSIONAL** use only.

DO NOT add wetter unless spraying at high volume. Where GUERRILLA® is mixed with water at less than 400 mL/100L of water, add 60 mL 1000g/L NON-IONIC SURFACTANT per 100L spray.

DO NOT spray plants, which are waterlogged, under stress of any kind or covered with soil or dust.

DO NOT spray plants covered with heavy dew, but rain following spraying will not affect results.

DO NOT sow or cultivate for 1 hour after spraying but operations should commence within 7 days.

DO NOT apply with high pressure handwand for control of annual weeds.

For ground application only – DO NOT use through aircraft, misting machines or hand-held ultra-low volume controlled droplet applications (CDA units).

SPRAY DRIFT RESTRAINTS:

Specific definitions for terms used in this section of the label can be found at apvma.gov.au/spraydrift

DO NOT allow **bystanders** to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an unacceptable impact to **native vegetation, agricultural crops, landscaped gardens** and **aquaculture production**, or cause contamination of plant or livestock commodities, outside the application site from **spray drift**.

Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

DO NOT apply unless **wind speed** is between 3 and 20 kilometres per hour at the **application site** during the time of application

DO NOT apply if there are **hazardous surface temperature inversion** conditions present at the **application site** during the time of application. **Surface temperature inversion conditions** exist most evenings one to two hours before sunset and persist until one to two hours after sunrise

CROP USE OR SITUATION	WEEDS CONTROLLED	STATE	RATE/ha	CRITICAL COMMENTS
Aid to Cultivation to minimise cultivation and prepare a clean bed for sowing	Annual grass and broadleaf weed control	QLD, NSW, VIC, SA, TAS, NT, ACT only	#	Where cultivation follows spraying, it may commence one hour after spraying but should be completed within 7 days. Where heavy weed growth is present at spraying, a better seed bed will result if cultivation is delayed 3-5 days. Use the higher rates for dense, more mature weed stands. Wild Oats must have at least two leaves. If Diquat 200g/L is used, the lower GUERRILLA® rate should be sufficient to control dense mature weeds. Pasture: Remains of old pasture should be reduced by continuous heavy grazing. Remove stock 3-5 days before spraying to allow to freshen up.
	Early Autumn sowing		750 mL – 1L	
	Winter, Spring and early Summer sowing	QLD, NSW, VIC, SA, TAS, NT, ACT only	1 – 1.5L	
	Wild Oats at 2-5 leaf stage in Autumn/Winter	QLD, VIC, SA, TAS, NT only	375mL – 500mL	
		NSW, ACT only	375 mL	

CROP USE OR SITUATION	WEEDS CONTROLLED	STATE	RATE/ha	CRITICAL COMMENTS
Fallows	Flaxleaf Fleabane	All States	1.7 – 2L	Imtrade Wetter 1000, Imtrade Pro 700 Surfactant or Imtrade Pronto Spray Adjuvant may be added at a rate of 0.2% v/v if climatic conditions require. Using water rates of 80 L/ha or greater is recommended.
Wild Oat control in Spring Fallows	Wild Oats at 2-5 leaf stage	QLD, NSW, NT, ACT only	750 mL – 1.25L	Use higher rate for summer growth. Avoid spraying under hot, dry conditions. Best results will be obtained when spraying is carried out in the late evening.
Selective Weed Control: Autumn/early winter - Annual and Perennial Clover	Annual grasses and some broadleaf weed control except Paterson's Curse, Sorrel, Dock, Shepherd's Purse and some Thistles. Control of some broadleaf weeds including Paterson's Curse, Sorrel, Dock, Shepherd's Purse and some Thistles will not be achieved. Alternative methods such as the spray-graze technique with 2,4-D or MCPA should be considered.	All States	# 375 mL to 750mL 750 mL to 1L	Use the higher rates for dense weed stands. Use the higher rate in Winter/early Spring when barley grass is present. All Applications: Graze pastures continuously after the seasonal break to a height of 2-4cm. Remove stock 2-3 days before spraying to allow weeds to freshen up. DO NOT apply until clover has reached the 6 leaf stage. DO NOT spray clovers which are affected by insect attack, disease or moisture stress. DO NOT use on clover pastures growing in water repellent sands or other situations subject to moisture stress at or immediately following treatment. Poor recovery of the clover will result. DO NOT APPLY TO MEDICS

CROP USE OR SITUATION	WEEDS CONTROLLED	STATE	RATE/ha	CRITICAL COMMENTS
Spray topping to reduce seed set	Grasses generally (particularly Annual Ryegrass)	All States	250 mL	Heavily graze paddocks during spring flush to encourage even head development. Remove stock 2-3 weeks before the anticipated maturity date of the target species. However, if this is not feasible through lack of stock it is preferable to allow the pasture to mature ungrazed. Delay spraying until the last seed-heads at the bottom of the plant have emerged and initial signs of haying off appear. Spray with a calibrated boom spray raised to give double overlap at the level of the seed heads.
Pastures	Barley Grass	All States	250 mL	Manage paddocks as above. Spray after head emergence but when all seed heads are green and there is no sign of haying off. Inspect paddocks before returning stock. Provided spraying was carried out before hardening seeds are present, harrow to knock seeds from the heads. DO NOT introduce lambs into paddock until safe from the risk of seed injury. If seasonal conditions favour regeneration, stock should be returned to selectively graze new shoots. Spray with a calibrated boom raised to give double overlap at the level of the seed heads.
	Saffron Thistle	NSW, SA, ACT only		Spray after the plant begins to run to head until flowering.
Prevention of Annual Ryegrass toxicity	Spray top – Graze to destroy seed heads	WA only	250 mL	<p>Grazing management as for spray topping above. Remove stock 3-4 weeks before anticipated maturity date. Spray must be applied within 10 days after emergence of the first Ryegrass seed heads.</p> <p>To ensure adequate control of toxin development, heavy continuous grazing is essential from 1 day after spraying until the pasture has completely hayed off.</p> <p>The required stocking rate will vary but must be sufficient to keep all re-growth after spraying completely eaten off to prevent further growth producing new seed heads which could become toxic.</p>

CROP USE OR SITUATION	WEEDS CONTROLLED	STATE	RATE/ha	CRITICAL COMMENTS
Hay freezing	Maximum retention of protein in standing dry feed.	All States	500 mL	Graze paddocks as for spray topping above. Remove 3-4 weeks before the anticipated maturity date. Apply prior to commencement of haying off regardless of the grass species involved. Spray with a calibrated boom spray raised to give double overlap at the level of the seed heads.
Orchards and Vineyards	Annual Weed control	QLD, VIC, SA, WA, TAS, NT only	# + 1 to 2L/ sprayed ha	Spray as necessary for control of annual weeds. Avoid contact to crop foliage. GUERRILLA® will not harm trees or vines with mature brown bark if this alone is sprayed. Use the higher rate for dense weed growth.
		NSW only	# + 1.1L/ sprayed ha	If Fat Hen <i>Chenopodium album</i> or <i>Portulaca</i> spp. are present and GUERRILLA® rate is less than the ratio 800 mL/100L add 120 mL Imtrade Penetrate Wetter per 100L spray mix.
Orchards and Vineyards	Flaxleaf Fleabane	All States	1.7 – 2L	Flaxleaf Fleabane: Imtrade Wetter 1000, Imtrade Pro 700 Surfactant or Imtrade Pronto Spray Adjuvant may be added at a rate of 0.2% v/v if climatic conditions require. Using water rates of 80 L/ha or greater is recommended. Avoid contact to crop foliage.
Potatoes	Pre-harvest weed control	All States	# 1.75L	Spray 4-5 weeks before digging and after tops have died down.

CROP USE OR SITUATION	WEEDS CONTROLLED	STATE	RATE/ha	CRITICAL COMMENTS
Non-Agricultural situations, around sheds, roadways, paths.	Annual weed control	All States	# + 1 to 2.5L/ha or 125 mL/ 100L	Spray to thoroughly wet weed growth. GUERRILLA® can be combined with soil residual herbicides i.e. Simazine 900 DF Herbicide, to give rapid knockdown and prolonged weed control. Use the higher rate for dense weed growth.
	Columbus Grass	NSW only	+ Spot Spraying 100 mL/ 100L plus 1L Smack Water Soluble Herbicide Boom Spray 1.4 to 2.8L/ha plus 12 to 22L Smack Water Soluble Herbicide	
Non-Agricultural situations, around sheds, roadways, paths.	Flaxleaf Fleabane	All States	1.7 – 2L	Flaxleaf Fleabane: Imtrade Wetter 1000, Imtrade Pro 700 Surfactant or Imtrade Pronto Spray Adjuvant may be added at a rate of 0.2% v/v if climatic conditions require. Using water rates of 80 L/ha or greater is recommended.
Firebreaks	Knock down weed growth to eliminate fire hazard or assist firebreak burn.	All States	1 to 2.5L	Apply mid-Winter to early Summer. Use the higher rate for dense weed growth. After desiccation is complete, the sprayed area may be burnt (normally 7-10 days after spraying). GUERRILLA® can be combined with soil residual herbicides i.e. Simazine 900 DF or Atrazine 900 DF or Flowable Atrazine Herbicides, to give rapid knockdown and prolonged weed control.

Capeweed or *Erodium* spp. present: Add Diquat 200g/L at 750 mL to 1.5L/ha (125 mL to 250 mL/100L for high volume spraying). Use higher rate for plants more than 10cm diameter.
+ If GUERRILLA® rate is less than the ratio 330 mL/100L add 60 mL of Imtrade Penetrate Wetter per 100L of spray mix.

OPTICAL SPOT SPRAY TECHNOLOGIES

Note: Calibrate the sprayer to spray the equivalent of 100L/ha

For weed cover between 0% and 30% only. If percentage weed cover exceeds 30%, use approved boom spray rates.

Follow on cropping should not commence for at least 5 days after application.

SITUATION	WEEDS	RATES	CRITICAL COMMENTS
Fallow	Yellow Vine (Caltrop)	1.9-5.6L/ 100L	Apply to rosette to flowering plants. Use higher rate on late flowering/mature plants or plants under moisture stress.
	Barnyard Grass		Use higher rate on large mature plants
	Bladder Ketmia		
	Fleabane	3.75- 5.6L/ 100L	Apply to rosette to flowering plants. Use higher rate on late flowering/mature plants or plants under moisture stress.
	Sowthistle		Apply to flowering plants. Use higher rate on late flowering/mature plants or plants under moisture stress.
	Turnip Weed		Use higher rate on large mature plants
	Australian Bindweed	5.6L/ 100L	Apply from seedling to 60cm in diameter plants.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

FOR USE ONLY AS AN AGRICULTURAL AND HORTICULTURAL HERBICIDE, THIS PRODUCT IS TOO HAZARDOUS TO BE USED IN THE HOME GARDEN.

WITHHOLDING PERIOD – DO NOT GRAZE OR CUT SPRAYED VEGETATION FOR STOCK FOOD FOR AT LEAST 1 DAY, OR GRAZE HORSES FOR 7 DAYS AFTER APPLICATION.

REMOVE STOCK FROM TREATED AREAS 3 DAYS BEFORE SLAUGHTER.

GENERAL INSTRUCTIONS

FOR USE ONLY AS AN AGRICULTURAL AND HORTICULTURAL HERBICIDE, THIS PRODUCT IS TOO HAZARDOUS TO BE USED IN THE HOME GARDEN.

This product kills annual grasses and most annual broadleaf weeds in specific situations and should not be used for any other purpose. Quickly kills green plant tissue on contact. It is immediately inactivated in the soil or heavy dew. The principle of selective weed control with this product is that annual weeds are killed but perennial plants and clovers recover after an initial scorch. The control of annual weeds by spraying with this product will allow the desirable perennial species to thicken up at the expense of the weeds. Moisture and fertility should not be limiting at spraying and the proportion of desirable species must be great enough for them to fill in the areas previously occupied by weeds. Long-term weed control can be obtained following the quick knockdown given by this product if it is combined with soil residual chemicals.

READ ALL SAFETY DIRECTIONS before commencing work.

DO NOT use hand-held ultra-low volume controlled droplet applicators (CDA units), boomless jets or misting machines.

MIXING

Add the required quantity of product to water in the spray tank and agitate to give even mixing. Agitate again if left standing.

Wetting Agent

This product contains a wetting agent and additional wetter is not required unless high volume spraying results in excessive dilution of the wetter content. This will occur when product rates fall below 400 mL per 100L of spray. Under such circumstances wetter should be added at the rate of 60 mL of 1000g/L NON-IONIC SURFACTANT per 100L of spray mix. Where Fat Hen and Portulaca are present in orchard or vineyard situations, extra wetter should be used when this product ratio is less than 800 mL per 100L. Add wetter at double the above recommendations. **DO NOT** use alkaline or anionic wetting agents.

Clean Water

Mix this product with clean water only. Water should be clean and free from clay, silt and algae. Providing it meets this requirement, saline water, water collected from roofs, bore water, dam water and water from creeks may be used.

APPLICATION

(i) Cereals and Broadacre Spraying

Use only through a properly calibrated boom spray, which should be fitted with flat fan jets and adjusted to a height to give at least double overlap of the spray at the top of the weeds being sprayed. Spraying pressures should be in the range of 200-300 kPa. Speed of travel should be in the range of 6-15 km/hr. It is essential that a good marking system be used. If a disc marked is used, it must be mounted so as to turn the soil back on to the area sprayed. It is essential to obtain good leaf coverage with the spray and volumes of dilute spray must be adjusted according to density of weed growth. 100L/ha may be used for seedlings or well grazed weeds up to 2cm high. For plant height 2-5cm use 150L/ha and up to 6-10cm use 200L/ha. Spray volumes may be as low as 50L/ha (30L/ha in WA) for weed growth below 5cm high, or for spray topping and hay freezing.

Equipment must be appropriate to this volume, properly calibrated and fitted with spraying tips designed to give medium spray droplet size category according to nozzle manufacturer specifications that refer to the ASAE S572 Standard or the BCPC Guideline.

READ ALL SAFETY DIRECTIONS before commencing work.

(ii) High Volume Applications

Higher volumes will generally be required to give good coverage of weed growth in situations other than those specified under cereals and other broadacre crops.

READ ALL SAFETY DIRECTIONS before commencing work.

(iii) Wash spray equipment with clean water immediately after use. This product is highly corrosive to metals particularly galvanised iron and aluminum and should not be left for long periods in tanks or equipment made of these materials.

READ ALL SAFETY DIRECTIONS before commencing work.

Spraying Conditions

Avoid spraying plants under stress from waterlogging, frost, and drought etc. or covered with dust and soil. Results will be better if application is made in dull weather or at the end of the day. Light rain following spraying will not affect results. Avoid drift into neighboring crops.

COMPATIBILITY

GUERRILLA® combines satisfactorily with the soil active herbicides Atrazine 900DF and Flowable Atrazine 600, Simazine 900DF and Flowable Simazine where prolonged weed control is required as well as a quick knockdown. This product is compatible with Imtrade Penetrate Wetter, Diquat, Imtrade Spraykill 250 Herbicide, Spray.Seed® 250, Dicamba 500 g/L, 2,4-D Amine 625, Nufarm Surpass® 475, MCPA 500 (no more than 1L/800 mL GUERRILLA®), Chlorsulfuron 750 g/kg, Oryzalin/Trifluralin 250 g/L, Imtrade Tri-Allate 500 EC Herbicide, Oxyfluorfen 240 g/L and Imtrade Trifluralin 480 EC Herbicide.

RESISTANT WEEDS WARNING

GROUP **22|34** HERBICIDE

GUERRILLA® is a member of the bipyridyl and triazole groups of herbicides. GUERRILLA® has the “photosynthesis at photosystem I inhibitor” and the inhibitor of carotenoid biosynthesis modes of action. For weed resistance management GUERRILLA® is a Group **22** and Group **34** Herbicide. Some naturally occurring weed biotypes resistant to GUERRILLA® and other “photosynthesis at photosystem I inhibitor” and inhibitor of carotenoid biosynthesis herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by GUERRILLA® or other “photosynthesis at photosystem I inhibitor” and inhibitor of carotenoid biosynthesis herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Imtrade CropScience accepts no responsibility for any losses that may result from the failure of GUERRILLA® to control resistant weeds.

PRECAUTIONS

RE-ENTRY PERIOD

DO NOT enter treated areas until the spray has dried. If prior entry is necessary, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves and impervious footwear. Contaminated clothing must be washed after each day's use.

WARNING: DO NOT APPLY WITH HIGH PRESSURE HANDWAND FOR CONTROL OF ANNUAL WEEDS.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT

Domestic pets and poultry – keep away from treated areas. **DO NOT repack. DO NOT** contaminate ponds, waterways or ditches with the chemical or used container. This formulation should not be applied on or near water which is used for human consumption, livestock watering or irrigation purposes or water used for commercial or recreational fishing.

STORAGE AND DISPOSAL

The method of disposal of the container depends on the container type. Read the Storage and Disposal instructions on the label that is attached to the container.

Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight.

All other non-refillable containers

Triple rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemical on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers and product. **DO NOT** re-use empty container for any other purpose.

Envirodrum - Micro Matic Valve (110L)

Store the original sealed Envirodrum in a cool, well ventilated area. **DO NOT** store for prolonged periods in direct sunlight. **DO NOT** tamper with the Micro Matic valve or the security seal. **DO NOT** contaminate the Envirodrum with water or any other foreign matter. After each use of the product please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase.

Refillable Containers (1000L only)

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

FOR USE ONLY AS AN AGRICULTURAL AND HORTICULTURAL HERBICIDE, THIS PRODUCT IS TOO HAZARDOUS TO BE USED IN THE HOME GARDEN.

SAFETY DIRECTIONS

GENERAL

Very dangerous. Poisonous if inhaled or swallowed. Harmful if absorbed by skin contact. Will irritate the skin. Will damage the eyes. Avoid contact with eyes and skin. **DO NOT** inhale spray mist.

WHEN MIXING AND USING

When opening the container, preparing spray and using the prepared spray wear cotton overalls, over normal clothing, buttoned to the neck and wrist, elbow-length chemical resistant gloves, goggles and full facepiece respirator with dust cartridge or canister. If applying by hand, when opening the container, preparing spray and using the prepared spray wear cotton overalls, over normal clothing, buttoned to the neck and wrist and a washable hat, elbow-length chemical resistant gloves, goggles and disposable mist face mask covering mouth and nose.

AFTER USE

Wash hands after use. After each day's use wash gloves, goggles, respirator and if rubber wash with detergent and warm water and contaminated clothing.

SPRAY APPLICATION

DO NOT work in spray mist. **DO NOT** continue to use if skin irritation or nosebleed occurs. This may be caused by exposure to spray mist as a result of incorrect use of equipment or adverse climatic conditions. Stop and review handling and spraying techniques before further spraying. If symptoms persist seek medical advice.

Where there is a risk of exposure to spray mist, wear waterproof footwear and waterproof protective clothing, impervious gauntlet length gloves (rubber or PVC), goggles and a face mask and respirator covering nose and mouth and capable of filtering spray droplets. A high efficiency type particulate respirator is recommended by in any event use a respirator which complies with the requirements of AS1716 (Standards Association of Australia). Further advice on safety equipment should be obtained from a safety equipment manufacturer. Avoid contacting vegetation wet with spray but if necessary to do so wear waterproof footwear and waterproof protective clothing and gloves.

FIRST AID

If poisoning occurs get to a doctor or hospital quickly. If in eyes, hold eyes open and flood with water for at least 15 minutes and see a doctor.

NOTE TO PHYSICIANS

For additional advice on the treatment of paraquat poisoning please consult the booklet "The Treatment of Paraquat Poisoning: A Guide for Doctors".

Additional GHS Hazard & Precautionary Statements

•May cause respiratory irritation; •Suspected of damaging the unborn child; •Causes damage to organs through prolonged or repeated exposure; •Very toxic to aquatic life with long-lasting effects; •**DO NOT** breathe fumes, mists, vapours or spray; •**DO NOT** get in eyes, on skin or on clothing; •Wash contacted areas thoroughly after handling; •**DO NOT** eat, drink or smoke when using this product; •Use only outdoors or in a well-ventilated area; •Remove all contaminated clothing immediately; •**IF SWALLOWED:** Rinse mouth. **DO NOT** induce vomiting; •**IF ON SKIN:** Wash with plenty of soap and water; •**IF INHALED:** Remove victim to fresh air and keep at rest in a position comfortable for breathing; •If skin irritation occurs: Get medical advice; •If eye irritation persists: Get medical advice; •Collect spillage; •Not combustible. Use extinguishing media suited to burning materials. Water fog or fine spray is the preferred medium for large fires; •Store locked up; •Store in a dry place.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from your supplier or Imtrade CropScience website at www.imtrade.com.au

CONDITIONS OF SALE

Imtrade CropScience warrants that this product conforms to the chemical description on the label. The use of GUERRILLA® being beyond the control of the manufacturer, Imtrade CropScience shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Imtrade's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Imtrade CropScience has any authority to add to or alter these conditions.

GUERRILLA® is a Registered Trademark of Imtrade Australia Pty Ltd
® Other Registered Trademarks

UN 3016	BIPYRIDILUM PESTICIDES LIQUID, TOXIC, N.O.S. (contains Paraquat)
IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE	
PG III	HAZCHEM 2X



APVMA Approval No: 89484/133061